



**Republic of Croatia**

**Ministry of Environmental Protection,  
Physical Planning and Construction**

# **NATIONAL INVENTORY REPORT 2007**

**Submission to the United Nations  
Framework Convention on Climate Change**



With the contribution of the LIFE  
Financial instrument of the EC



Ordered by: Ministry of Environmental Protection,  
Physical Planning and Construction

Contract No.: Klasa: 112-04/05-01/47  
Ur. Broj: 531-05/02-VG-05-02

Title:

# **NATIONAL INVENTORY REPORT 2007**

**Croatian greenhouse gas inventory  
for the period 1990-2005**

Project Coordinator: **Andrea Hublin**

List of Authors: **Iva Švedek** (Energy CRF sector 1)  
**Andrea Hublin** (Industrial processes CRF sector 2,  
Solvent and Other Product Use CRF sector 3, Waste  
CRF sector 6)  
**Veljko Vorkapić** (Agriculture CRF sector 4, LULUCF  
CRF sector 5)  
**Jasmina Burek** (CRF manager)  
**Davor Vešligaj** (QA/QC manager)

External Author: **Branko Vuk** (Energy balance, EIHP)

Atmospheric Protection  
Department Manager:

General Manager:

**Davor Vešligaj**

**Zdravko Mužek**

Zagreb, October 2007

## CONTENT:

<b>LIST OF ABBREVIATIONS.....</b>	<b>i</b>
<b>LIST OF TABLES AND FIGURES .....</b>	<b>ii</b>
<b>EXECUTIVE SUMMARY .....</b>	<b>vi</b>
ES.1. BACKGROUND INFORMATION ON GHG INVENTORIES AND CLIMATE CHANGE .....	vi
ES.1.1. INSTITUTIONAL AND ORGANIZATIONAL STRUCTURE OF GREENHOUSE GAS EMISSIONS INVENTORY PREPARATION .....	vii
ES.2. SUMMARY OF NATIONAL EMISSION AND REMOVAL RELATED TRENDS .....	viii
ES.3. OVERVIEW OF SOURCES AND SINK CATEGORY EMISSION ESTIMATES AND TRENDS ..	ix
ES.3.1. CARBON DIOXIDE EMISSION (CO <sub>2</sub> ) .....	xi
ES.3.1.1. Energy sector .....	xi
ES.3.1.2. Industrial processes .....	xiii
ES.3.1.3. CO <sub>2</sub> removals .....	xiv
ES.3.2. METHANE EMISSION (CH <sub>4</sub> ) .....	xv
ES.3.3. NITROUS OXIDE EMISSION (N <sub>2</sub> O) .....	xvi
ES.3.4. HALOGENATED CARBONS (HFCs, PFCs) AND SF <sub>6</sub> EMISSIONS .....	xvii
ES.4. EMISSION OF INDIRECT GREENHOUSE GASES .....	xvii
<b>1. INTRODUCTION.....</b>	<b>1</b>
1.1. BACKGROUND INFORMATION ON GHG INVENTORIES AND CLIMATE CHANGE.....	1
1.2. BRIEF DESCRIPTION OF THE INSTITUTIONAL ARRANGEMENT FOR INVENTORY PREPARATION.....	2
1.3. BRIEF DESCRIPTION OF THE PROCESS OF INVENTORY PREPARATION .....	3
1.4. BRIEF DESCRIPTION OF METHODOLOGIES AND DATA SOURCES USED .....	4
1.5. BRIEF DESCRIPTION OF KEY CATEGORIES.....	6
1.6. INFORMATION ON THE QA/QC PLAN INCLUDING VERIFICATION AND TREATMENT OF CONFIDENTIALITY ISSUES .....	7
1.6.1. QA/QC plan and procedures .....	7
1.6.2. Verification and confidentiality issues .....	8
1.7. GENERAL UNCERTAINTY EVALUATION .....	9
1.8. GENERAL ASSESSMENT OF THE COMPLETENESS .....	9
<b>2 TRENDS IN GREENHOUSE GAS EMISSIONS .....</b>	<b>10</b>
2.1. BRIEF DESCRIPTION AND INTERPRETATION OF EMISSION TRENDS FOR AGGREGATED GREENHOUSE GAS EMISSIONS .....	10
2.2. BRIEF DESCRIPTION AND INTERPRETATION OF EMISSION TRENDS BY GAS .....	11
2.2.1. Carbon dioxide – CO <sub>2</sub> .....	11
2.2.2. Methane – CH <sub>4</sub> .....	12
2.2.3. Nitrous oxide – N <sub>2</sub> O .....	12
2.2.4. Fluorocarbons – HFCs and PFCs.....	12
2.2.5. Sulphur hexafluoride SF <sub>6</sub> .....	12
2.3. BRIEF DESCRIPTION AND INTERPRETATION OF EMISSION TRENDS BY CATEGORY .....	13
2.3.1. Energy.....	13
2.3.2. Industrial Processes .....	14
2.3.3. Solvent and Other Product Use .....	14
2.3.4. Agriculture .....	14
2.3.5. Waste .....	14
2.4. BRIEF DESCRIPTION AND INTERPRETATION OF EMISSION TRENDS FOR INDIRECT GREENHOUSE GASSES AND SO <sub>2</sub> .....	15

<b>3. ENERGY (CRF sector 1)</b>	<b>16</b>
3.1. OVERVIEW OF SECTOR	16
3.1.1. Introduction	16
3.1.2. Energy structure	17
3.1.3. Comparison of the sectoral approach with the reference approach	18
3.1.4. International bunker fuels	19
3.1.5. Feedstocks and non-energy use of fuels	19
3.1.6. CO <sub>2</sub> Capture from flue gases and subsequent CO <sub>2</sub> storage	20
3.1.7. Country-specific issues	20
3.2. FUEL COMBUSTION ACTIVITIES (CRF 1.A.)	21
3.2.1. Source category description	21
3.2.2. Methodological issues	28
3.2.3. Uncertainties and time-series consistency	30
3.2.4. Source-specific QA/QC	32
3.2.5. Source-specific recalculations	33
3.2.6. Source-specific planned improvements	34
3.3. FUGITIVE EMISSIONS FROM FUELS (CRF 1.B.)	35
3.3.1. Source category description	35
3.3.2. Methodological issues	37
3.3.3. Uncertainties and time series consistency	38
3.3.4. Source-specific QA/QC	38
3.3.5. Source-specific recalculations	39
3.3.6. Source-specific planned improvements	39
3.4. OVERVIEW OF GHG EMISSIONS FROM ENERGY SECTOR	40
3.5. REFERENCES	41
<b>4. INDUSTRIAL PROCESSES (CRF sector 2)</b>	<b>42</b>
4.1. OVERVIEW OF SECTOR	42
4.2. MINERAL PRODUCTS (CRF 2.A.)	43
4.2.1. Cement production	43
4.2.2. Lime production	48
4.2.3. Limestone and dolomite use	51
4.2.4. Soda ash production and use	53
4.2.5. Production and use of miscellaneous mineral products	55
4.3. CHEMICAL INDUSTRY (CRF 2.B.)	58
4.3.1. Ammonia production	58
4.3.2. Nitric acid production	60
4.3.3. Production of other chemicals	63
4.4. METAL PRODUCTION (CRF 2.C.)	66
4.4.1. Iron and steel production	66
4.4.2. Ferroalloys production	68
4.4.3. Aluminium production	70
4.5. OTHER PRODUCTION (CRF 2.D.)	72
4.5.1. Pulp and paper	72
4.5.2. Food and drink	72
4.6. CONSUMPTION OF HALOCARBONS AND SF <sub>6</sub> (CRF 2.F.)	74
4.6.1. Refrigeration and air conditioning equipment	74
4.6.2. Other consumption of HFCs, PFCs and SF <sub>6</sub>	76
4.7. NON - ENERGY USE (CRF 2.G.)	77
4.8. EMISSION OVERVIEW	78

4.8.1. GHG emissions .....	78
4.8.2. Indirect GHG emissions .....	82
4.9. REFERENCES .....	84
<b>5. SOLVENT AND OTHER PRODUCT USE (CRF sector 3).....</b>	<b>85</b>
5.1. SOLVENT AND OTHER PRODUCT USE .....	85
5.1.1. Source category description .....	85
5.1.2. Methodological issues .....	85
5.1.3. Uncertainties and time-series consistency .....	88
5.1.4. Source-specific QA/QC and verification .....	88
5.1.5. Source-specific recalculations .....	89
5.1.6. Source-specific planned improvements.....	89
5.2. REFERENCES .....	90
<b>6. AGRICULTURE (CRF sector 4) .....</b>	<b>91</b>
6.1. OVERVIEW OF SECTOR .....	91
6.2. CH <sub>4</sub> EMISSIONS FROM ENTERIC FERMENTATION (CRF 4.A.) .....	93
6.2.1. Source category description .....	93
6.2.2. Methodological issues .....	93
6.2.3. Uncertainties and time-series consistency .....	94
6.2.4. Source specific recalculations .....	94
6.3. MANURE MANAGEMENT – CH <sub>4</sub> EMISSIONS (CRF 4.B.) .....	95
6.3.1. Source category description .....	95
6.3.2. Methodological issues .....	95
6.3.3. Uncertainties and time-series consistency .....	95
6.3.4. Source specific recalculations .....	96
6.4. N <sub>2</sub> O EMISSIONS FROM MANURE MANAGEMENT (CRF 4.B.).....	96
6.4.1. Source category description .....	96
6.4.2. Methodological issues .....	96
6.4.3. Uncertainties and time-series consistency .....	97
6.4.4. Source specific recalculations .....	97
6.5. AGRICULTURAL SOILS (CRF 4.D.) .....	98
6.5.1. Direct emission from agricultural soils .....	98
6.5.2. Direct emission of N <sub>2</sub> O from animals.....	101
6.5.3. Indirect N <sub>2</sub> O emissions from nitrogen used in agriculture .....	102
6.6. SOURCE SPECIFIC QA/QC AND VERIFICATION .....	104
6.7. SOURCE SPECIFIC PLANNED IMPROVEMENT .....	104
6.8. REFERENCES .....	104
<b>7. LAND-USE, LAND USE CHANGE AND FORESTRY (CRF sector 5) .....</b>	<b>106</b>
7.1. OVERVIEW OF SECTOR .....	106
7.2. SOURCE CATEGORY .....	108
7.2.1. Source category description .....	108
7.2.2. Methodological issues .....	108
7.2.3. Uncertainties and time-series consistency .....	111
7.2.4. Source-specific QA/QC and verification .....	111
7.2.5. Source-specific recalculations .....	111
7.2.6. Source-specific planned improvements.....	111
7.3. REFERENCES .....	112

<b>8. WASTE (CRF sector 6).....</b>	<b>113</b>
8.1. OVERVIEW OF SECTOR .....	113
8.2. SOLID WASTE DISPOSAL ON LAND (CRF 6.A.).....	114
8.2.1. Source category description .....	114
8.2.2. Methodological issues .....	114
8.2.3. Uncertainties and time-series consistency .....	116
8.2.4. Source-specific QA/QC and verification .....	117
8.2.5. Source specific recalculations .....	117
8.2.6. Source-specific planned improvements.....	117
8.3. WASTEWATER HANDLING (CRF 6.B.) .....	119
8.3.1. Source category description .....	119
8.3.2. Methodological issues .....	119
8.3.3. Uncertainties and time-series consistency .....	120
8.3.4. Source-specific QA/QC and verification .....	121
8.3.5. Source specific recalculations .....	121
8.3.6. Source-specific planned improvements.....	121
8.4. WASTE INCINERATION (CRF 6.C.) .....	122
8.4.1. Source category description .....	122
8.4.2. Methodological issues .....	122
8.4.3. Uncertainties and time-series consistency .....	122
8.4.4. Source-specific QA/QC and verification .....	122
8.4.5. Source specific recalculations .....	123
8.4.6. Source-specific planned improvements.....	123
8.5. EMISSION OVERVIEW .....	124
8.6. REFERENCES .....	125
<b>9. RECALCULATIONS AND IMPROVEMENTS.....</b>	<b>126</b>
9.1. EXPLANATIONS AND JUSTIFICATIONS FOR RECALCULATIONS. INCLUDING IN RESPONSE TO THE REVIEW PROCESS .....	126
9.1.1. Changes or refinements in methods.....	126
9.1.2. Correction of errors.....	129
9.2. THE IMPLICATION OF THE RECALCULATIONS ON THE LEVEL AND TREND. INCLUDING TIME SERIES CONSISTENCY.....	130
9.3. PLANNED IMPROVEMENTS TO THE INVENTORY .....	132
<b>ANNEX 1 .....</b>	<b>.....</b>
KEY CATEGORIES.....	.....
A1.1. DESCRIPTION OF METHODOLOGY USED FOR IDENTIFYING KEY CATEGORIES .....	ANNEX 1-1
A1.2. TABLES 7.A1-7.A3 OF THE IPCC GOOD PRACTICE GUIDANCE.....	ANNEX 1-3
<b>ANNEX 2.....</b>	<b>.....</b>
DETAILED DISCUSSION OF ACTIVITY DATA AND EMISSION FACTORS FOR ESTIMATING CO <sub>2</sub> EMISSIONS FROM FOSSIL FUEL COMBUSTION .....	.....
<b>ANNEX 3.....</b>	<b>.....</b>
CO <sub>2</sub> REFERENCE APPROACH AND COMPARISON WITH SECTORAL APPROACH, AND RELEVANT INFORMATION ON THE NATIONAL ENERGY BALANCE .....	.....

**ANNEX 4** .....

ASSESSMENT OF COMPLETENESS AND (POTENTIAL) SOURCES AND SINKS OF  
GREENHOUSE GAS EMISSIONS AND REMOVALS EXCLUDED.....

**ANNEX 5** .....

TABLE 6.1 OF THE IPCC GOOD PRACTICE GUIDANCE.....

**ANNEX 6** .....

QA/QC FRAMEWORK PLAN.....

## **LIST OF ABBREVIATIONS**

CDM	- Clean Development Mechanism (CDM)
CFC	- Chlorofluorocarbons
COPERT	- Computer Programme to Calculate Emissions from Road Transport
CORINAIR	- Core Inventory of Air Emissions in Europe
CPS Molve	- Central Gas Station Molve
CRF	- Common Reporting Format
EMEP	- Co-operative Programme for Monitoring and Evaluation of the Long Rang Transmission of Air Pollutants in Europe
ERT	- Expert Review Team
ET	- Emissions Trading
FAO	- Food and Agriculture Organization of the United Nations
GHG	- Greenhouse gas
GWP	- Global Warming Potential
HEP	- Croatian Electricity Utility Company
IEA	- International Energy Agency
IPCC	- Intergovernmental Panel on Climate Change
ISWA	- International Solid Waste Association
JI	- Joint Implementation
NGGIP	- National Greenhouse Gas Inventories Programme
NM VOC	- Non-methane Volatile organic Compounds
OECD	- Organisation for Economic Co-operation and Development
UNEP	- United Nations Environment Programme
UNFCCC	- United Nations Framework Convention on Climate Change
CBS	- Central Bureau of Statistics
EIHP	- Energy Institute "Hrvoje Požar"
CEE	- Cadastre of Emission in Environment
MZOPU	- Ministry of Environmental Protection and Physical Planning
INA	- Croatian Oil and Gas Company
ZGO	- Zagreb's Environmental Protection and Waste Management Company
APO	- Hazardous Waste Management Agency



## LIST OF TABLES AND FIGURES

### LIST OF TABLES

Table ES.2-1: Global warming potentials for certain gases .....	viii
Table ES.3-1: Emissions/removals of GHG by sectors for the period 1990-2005 (Gg CO <sub>2</sub> -eq) ....	ix
Table ES.3-2: Emissions/removals of GHG by gases for the period 1990-2005 (Gg CO <sub>2</sub> -eq) .....	ix
Table ES.3-3: CO <sub>2</sub> emission/removal by sectors from 1990-2005 (Gg CO <sub>2</sub> ) .....	xi
Table ES.3-4: CO <sub>2</sub> emission by sub-sectors from 1990-2005 (Gg CO <sub>2</sub> ) .....	xii
Table ES.3-5: CO <sub>2</sub> emission comparison due to fuel combustion (Gg).....	xii
Table ES.3-6: CO <sub>2</sub> emission from Industrial Processes for the period from 1990-2005 .....	xiii
Table ES.3-7: CH <sub>4</sub> emission in Croatia in the period from 1990-2005 (Gg CH <sub>4</sub> ).....	xv
Table ES.3-8: N <sub>2</sub> O emission in Croatia for the period from 1990-2005 (Gg N <sub>2</sub> O) .....	xvi
Table ES.4-1: Emissions of ozone precursors and SO <sub>2</sub> by different sectors (Gg).....	xviii
Table 1.4-1: Data sources for GHG inventory preparation .....	5
Table 2.2-1: Aggregated emissions and removals of GHG by gases (1990-2005) .....	11
Table 2.3-1: Aggregated emissions and removals of GHG by sectors (1990-2005) .....	13
Table 2.4-1: Emissions of ozone precursors and SO <sub>2</sub> by different sectors (Gg) .....	15
Table 3.1-1: The fuel consumption and CO <sub>2</sub> emissions from fuel combustion.....	18
Table 3.1-2: Fuel consumption and GHG emissions for inter. aviation and marine bunkers.....	19
Table 3.2-1: The CO <sub>2</sub> -eq emissions (Gg) from Energy Industries.....	21
Table 3.2-2: Generating capacities of HPPs, TPPs and NPP Krško.....	22
Table 3.2-3: Processing Capacities of Oil & Lube Refineries .....	23
Table 3.2-4: The CO <sub>2</sub> -eq emissions (Gg) from Manufacturing Industries and Construction.....	24
Table 3.2-5: The CO <sub>2</sub> -eq emissions (Gg) from Transport.....	25
Table 3.2-6: The CO <sub>2</sub> -eq emissions (Gg) from small stationary energy sources .....	27
Table 3.2-7: Emissions of ozone precursors and SO <sub>2</sub> from fuel combustion (Gg) .....	28
Table 3.3-1: The fugitive emissions of ozone precursors and SO <sub>2</sub> from oil refining .....	37
Table 3.3-2: The CO <sub>2</sub> emissions (Gg) from natural gas scrubbing in CGS Molve.....	37
Table 3.4-1: The CO <sub>2</sub> -eq emissions (Gg) from Energy sector .....	40
Table 4.2.1: Clinker production and emission factors (1990 - 2005).....	44
Table 4.2.2: Import/export quantities of clinker (1990 - 2005).....	45
Table 4.2-3: Cement production (1990-2005) .....	46
Table 4.2-4: Lime production (1990-2005) .....	49
Table 4.2-5: CO <sub>2</sub> emissions from quicklime and dolomitic lime production (1990-2005).....	49
Table 4.2-6: Limestone and dolomite use (1990-2005) .....	52
Table 4.2-7: Soda ash use (1990-2005).....	54
Table 4.3-1: Consumption and composition of gas in Ammonia Production (1990-2005).....	58
Table 4.3-2: Nitric acid production (1990-2005) .....	61

<i>Table 4.3-3: Production of other chemicals (1990-2005)</i> .....	63
<i>Table 4.3-4: Emissions of CH<sub>4</sub> from Production of Other Chemicals (1990-2005)</i> .....	64
<i>Table 4.4-1: Steel production (1990-2005)</i> .....	67
<i>Table 4.4-2: Production of ferroalloys (1990-2005)</i> .....	69
<i>Table 4.6-1: Emissions of HFCs (Gg CO<sub>2</sub>-eq) (1990 – 2005)</i> .....	75
<i>Table 4.8-1: Emissions of GHGs from Industrial Processes (1990-2005)</i> .....	78
<i>Table 4.8-2: Gases generated from different non-energy industrial process</i> .....	83
<i>Table 4.8-3: Emissions of indirect GHGs from Industrial Processes (1990-2005)</i> .....	83
<i>Table 5.1-1: NMVOC emissions from Solvent and Other Product Use (Gg) (1990-2005)</i> .....	86
<i>Table 6.1-1: Emission of greenhouse gases from agriculture (Gg)</i> .....	92
<i>Table 6.1-2: Emission of greenhouse gases from agriculture CO<sub>2</sub>-eq (Gg)</i> .....	92
<i>Table 7.2-1: Forest areas and annual increment of forests in Croatia according to Forest Management Plan (1986-1996; 1996-2006)</i> .....	108
<i>Table 7.2-2: Emission factors used in estimations</i> .....	110
<i>Table 7.2-3: Annual change in Carbon Stock in living biomass in Forest Land Remaining Forest Land (Gg CO<sub>2</sub>)</i> .....	110
<i>Table 8.2-1: Country-specific composition of waste</i> .....	114
<i>Table 8.2-2: Total annual MSW disposed to SWDSs and related MCF (1990-2005)</i> .....	115
<i>Table 8.3-1: Data for N<sub>2</sub>O emission calculation from Human Sewage (1990-2005)</i> .....	120
<i>Table 8.5-1: Emissions from Waste (1990-2005)</i> .....	124
<i>Table 9.2-1: Differences between NIR 2006 and NIR 2007 for 1990-2004 due to recalculation</i>	130
<i>Table 9.2-2: Differences between NIR 2006 and NIR 2007 for the emission trends 1990-2004</i>	131
<i>Table A1-1: Categories Assessed in Key Category Analysis</i> .....	A1-1
<i>Table A1-2: Key categories analysis – Level Assessment - Tier 1 (Including LULUCF)</i> .....	A1-3
<i>Table A1-3: Key categories analysis – Level Assessment - Tier 1 (Including LULUCF)</i> .....	A1-4
<i>Table A1-4: Key categories analysis – Trend Assessment - Tier 1 (Excluding LULUCF)</i> .....	A1-5
<i>Table A1-5: Key categories analysis – Trend Assessment - Tier 1 (Including LULUCF)</i> .....	A1-6
<i>Table A1-6: Key categories for Croatia – summary (Excluding LULUCF)</i> .....	A1-7
<i>Table A1-7: Key categories for Croatia – summary (Including LULUCF)</i> .....	A1-8
<i>Table A1-8: Changes in Key categories for Croatia based on the Level and Trend</i> .....	A1-9
<i>Table A2-1: The GHG emissions from Thermal Power Plants</i> .....	A2-1
<i>Table A2-2: The GHG emissions from Public Cogeneration Plants</i> .....	A2-1
<i>Table A2-3: The GHG emissions from Public Heating Plants</i> .....	A2-2
<i>Table A2-4: The GHG emissions from TPPs and PCPs (Tier 2), year 2005</i> .....	A2-3
<i>Table A2-5: The GHG emissions from Petroleum refining – own use of energy</i> .....	A2-4
<i>Table A2-6: The GHG emissions from Petroleum refining – heating/cogeneration plants*</i> .....	A2-4
<i>Table A2-7: The GHG emissions from manufact. of solid fuels and other energy industries</i> ....	A2-6
<i>Table A2-8: The GHG emissions from Manuf. Ind. and Constr. – liquid fuels and natural gas</i> .....	A2-6
<i>Table A2-9: The GHG emissions from Manuf. Industries and Construction – solid fuels</i> .....	A2-7
<i>Table A2-10: The number of road motor vehicles in Croatia</i> .....	A2-8

Table A2-11: GHG emissions from Road Transport.....	A2-9
Table A2-12: The GHG emissions from Domestic Air Transport .....	A2-9
Table A2-13: The GHG emissions from National Navigation.....	A2-9
Table A2-14: The GHG emissions from Railways .....	A2-10
Table A2-15: The GHG emissions from Commercial/Institutional.....	A2-11
Table A2-16: The GHG emissions from Residential sector .....	A2-12
Table A2-17: The GHG emissions from Agriculture/Forestry/Fishing .....	A2-13
Table A2-18: Methane emissions from Coal Mining and Handling from 1990 to 1999.....	A2-15
Table A2-19: Methane emissions from Oil and Gas Activities .....	A2-16
Table A3-1: Fuel combustion CO <sub>2</sub> emissions (Reference and Sectoral Approach).....	A3-1
Table A3-2: Net calorific values for different fossil fuels from 1990 to 2004.....	A3-2
Table A3-3: National energy balance for 2005 .....	A3-3
Table A4-1 GHGs and source/sink categories not considered in the Croatian inventory .....	A4-1
Table A5-1: Tier 1 Uncertainty Calculation and Reporting – excluding LULUCF .....	A5-1
Table A5-2: Tier 1 Uncertainty Calculation and Reporting – including LULUCF .....	A5-4
Table A6-1: The framework of the Croatian QA/QC Plan .....	A6-1
Table A6-2: An example of Inventory Data Record Sheet for 2005 in Waste .....	A6-2

## LIST OF FIGURES

Figure ES.3-1: Emissions and removals of the greenhouse gases in the Republic of Croatia by sectors in the period from 1990-2005 (Gg CO <sub>2</sub> -eq) .....	x
Figure ES.3-2: CO <sub>2</sub> emission removal in forestry sector from 1990-2005 (Gg CO <sub>2</sub> ) .....	xv
Figure ES.3-3: Halogenated carbons emission in the period from 1990-2005 (Gg CO <sub>2</sub> -eq) .....	xvii
Figure 2.2-1: Trend of GHG emissions, by gases .....	11
Figure 2.3-1: Trend of GHG emissions, by sectors .....	13
Figure 3.1-1: The contribution of different subsectors to GHG emission, year 2005.....	17
Figure 3.1-2: Structure of energy consumption .....	18
Figure 3.2-1: The CO <sub>2</sub> -eq emissions from energy industries .....	21
Figure 3.2-2: The CO <sub>2</sub> -eq emissions from Manufacturing Industries and Construction.....	25
Figure 3.2-3: The CO <sub>2</sub> -eq emissions from transport.....	26
Figure 3.2-4: The CO <sub>2</sub> -eq emissions from small stationary sources .....	27
Figure 3.3-1: The fugitive emissions of methane from coal mines.....	35
Figure 3.3-2: The fugitive emissions of methane from oil and gas activities .....	36
Figure 3.4-1: The CO <sub>2</sub> -eq emissions from Energy sector .....	40
Figure 4.1-1: Emissions of GHGs from Industrial Processes (1990-2005).....	43
Figure 4.2-1: Emissions of CO <sub>2</sub> from Cement Production (1990-2005) .....	46
Figure 4.2-2: Emissions of CO <sub>2</sub> from Lime Production (1990-2005) .....	50
Figure 4.2-3: Emissions of CO <sub>2</sub> from Limestone and Dolomite Use (1990-2005) .....	52

<i>Figure 4.2-4: Emissions of CO<sub>2</sub> from Soda Ash Use (1990-2005)</i> .....	54
<i>Figure 4.3-1: Emissions of CO<sub>2</sub> from Ammonia Production (1990-2005)</i> .....	59
<i>Figure 4.3-2: Emissions of N<sub>2</sub>O from Nitric Acid Production (1990-2005)</i> .....	62
<i>Figure 4.4-1: Emissions of CO<sub>2</sub> from Steel Production (1990-2005)</i> .....	67
<i>Figure 4.4-2: Emissions of CO<sub>2</sub> from Ferroalloys Production (1990-2005)</i> .....	69
<i>Figure 5.1-1: Emissions of NMVOC from Solvent and Other Product Use (1990-2005)</i> .....	85
<i>Figure 5.1-2: Emissions of CO<sub>2</sub> from Solvent and Other Product Use (1990-2005)</i> .....	88
<i>Figure 6.1-1: Agriculture GHG Sources (year 2005)</i> .....	91
<i>Figure 6.2-1: CH<sub>4</sub> emission from Enteric fermentation (Gg)</i> .....	93
<i>Figure 6.2-2: Number of dairy cattle, cattle, swine sheep, goats and horses in thousands</i> .....	94
<i>Figure 6.2-3: Number of poultry in thousands</i> .....	94
<i>Figure 6.3-1: CH<sub>4</sub> emission from Manure Management (Gg)</i> .....	95
<i>Figure 6.4-1: N<sub>2</sub>O Emissions from Manure Management (Gg)</i> .....	97
<i>Figure 6.5-1: Total N<sub>2</sub>O Emissions from Agricultural Soils (Gg)</i> .....	98
<i>Figure 6.5-2: Direct N<sub>2</sub>O Emissions from Agricultural Soils (Gg)</i> .....	99
<i>Figure 6.5-3: Direct N<sub>2</sub>O Emissions from Animals (Gg)</i> .....	101
<i>Figure 6.5-4: Indirect N<sub>2</sub>O Emissions from Agricultural (Gg)</i> .....	102
<i>Figure 8.1-1: Emissions of GHGs from Waste (1990-2005)</i> .....	113
<i>Figure 8.2-1: Emissions of CH<sub>4</sub> from Solid Waste Disposal on Land (1990-2005)</i> .....	116
<i>Figure 8.3-1: Emissions of N<sub>2</sub>O from Human Sewage (1990-2005)</i> .....	120

## EXECUTIVE SUMMARY

### ES.1. BACKGROUND INFORMATION ON GHG INVENTORIES AND CLIMATE CHANGE

Annual greenhouse gas inventory report of the Republic of Croatia is being prepared according to the UNFCCC reporting guidelines on annual inventories and IPCC methodology, continuously from 2001, when the first National Inventory Report was created in the framework of the preparation of the First National Communication. The preparation and submission of the National Inventory Report is supervised by the Ministry of Environmental Protection, Physical Planning and Construction. The preparation of the National Inventory Report (NIR) itself is outsourced to the national institutions which have the experience and capacities for data collection and emissions estimation. The quality assurance of the emission inventory is achieved through technical reviews organized by the UNFCCC Secretariat and carried out by nominated international expert reviewers. The main goal of the process of inventory preparation and inventory check is continuous improvement of its quality in the sense of accuracy, completeness, integrity, transparency and consistency. In this NIR, the inventory of the emissions and removals of the greenhouse gases is reported for the period from 1990 to 2005.

For the purposes of the greenhouse gas inventory preparation the methodology which is described in *Revised 1996 IPCC Guidelines for National GHG Inventories, (IPCC/UNEP/OECD/IEA) and Good Practice Guidance and Uncertainty Management in National GHG Inventories, 2000 (IPCC/NGGIP)* is being used. The important part of the inventory preparation is uncertainty assessment of the calculation and verification of the input data and results, all this with the aim to increase the quality and reliability of the emissions estimation.

The estimation includes the emissions which are the result of anthropogenic activities and these include the following greenhouse gases: carbon dioxide (CO<sub>2</sub>), methane (CH<sub>4</sub>), nitrous oxide (N<sub>2</sub>O), halogenated carbons (HFCs, PFCs), sulphur hexafluoride (SF<sub>6</sub>) and indirect greenhouse gases: carbon monoxide (CO), oxides of nitrogen (NO<sub>x</sub>), non-methane volatile organic compounds (NMVOCs) and sulphur dioxide (SO<sub>2</sub>). The greenhouse gases covered by the Montreal Protocol are reported in the framework of this international agreement and therefore are excluded from this Report.

Greenhouse gas emission sources and sinks are divided into six main sectors: Energy, Industrial Processes, Solvent and Other Product Use, Agriculture, Land Use, Land-Use Change and Forestry and Waste. Generally, the methodology for emission calculation could be described as a product of the particular economic activity (e.g. fuel consumption, cement production, number of animals, increase of wood stock etc.) with corresponding emission factor. The use of specific national emission factors is recommended wherever possible and justified, whereas on the contrary, the IPCC methodology gives default values of emission factors for all relevant activities of the particular sectors.

The greenhouse gas emission inventory will have important role in the first commitment period of the Kyoto Protocol, from 2008 to 2012, and particularly, in monitoring of the implementation of GHG mitigation policy and measures.

#### **ES.1.1. INSTITUTIONAL AND ORGANIZATIONAL STRUCTURE OF GREENHOUSE GAS EMISSIONS INVENTORY PREPARATION**

The important prerequisite for the efficient data management system and inventory preparation is clearly defined organization, authority and responsibility of the institutions participating in the process of inventory preparation, which includes number of steps in data collection and processing, calculating, control and verification of emission calculation, documentation, archiving and reporting to relevant international bodies. It could be stated that Croatia, respectively the Ministry of Environmental Protection, Physical Planning and Construction as a competent authority for inventory preparation uses decentralized model in which it has outsourced individual tasks in inventory preparation process to collaborative institutions which are partially public or governmental and partially in private ownership.

The main activity data sources for greenhouse gas emission calculation are Energy Institute "Hrvoje Požar" which prepares the national energy balance; Central Bureau of Statistics which collects data on raw materials and products for activities defined by National classification of economic activities on grounds of statistical research programmes; Croatian Center for Vehicles and Ministry of Internal Affairs which owns database on road and off-road vehicles; Ministry of Agriculture, Forestry and Water Management which owns data on forest-covered area. For the inventory preparation, the data gathered through questionnaires directly from individual emission sources and other scientific or professional institutions are also being used either for calculation or control of data obtained from the official publications. EKONERG - Energy Research and Environmental Protection Institute is national inventory focal point, responsible for emission calculation and preparation of annual greenhouse gas emission inventories according to contract with Ministry of Environmental Protection, Physical Planning and Construction. It should be noted that the financing of the inventory in the past period was mainly provided through the EC programme LIFE – Third Countries. This financial arrangement has certain advantages primarily in a sense of the efficient use of the existing resources, but on the other hand there are certain weaknesses of this system regarding middle-term and long-term planning and inventory improvement.

Considering the need for sustainable system of greenhouse gas emission monitoring, and commitment for establishment of the National System for the purposes of the Kyoto Protocol, the Ministry of Environmental Protection, Physical Planning and Construction under Article 46 of the Air Protection Act (Official Gazette No. 178/2004) has started the process of preparation of the Regulation on the Greenhouse Gases Emissions Monitoring in the Republic of Croatia in July 2006, which is confirmed in January 2007 (Official Gazette No. 1/2007). It should improve existing system of greenhouse gas emission monitoring and reporting in accordance with the requirements of the Kyoto protocol and relevant legislation of the EU. It is expected that the Regulation will enter into force in 2008.

## ES.2. SUMMARY OF NATIONAL EMISSION AND REMOVAL RELATED TRENDS

In this chapter the results of the greenhouse gas emission calculation in the Republic of Croatia are presented for the period from 1990 to 2005. The results are presented as total emissions of all greenhouse gases in CO<sub>2</sub> equivalents over sectors and then as emissions for the individual greenhouse gas by sectors. Since the certain greenhouse gases have different irradiation properties, and consequently different contribution to the greenhouse effect, it is necessary to multiply the emission of every pollutant with proper Global Warming Potential (GWP). The Global Warming Potential is a measure of the impact on greenhouse effect of the certain pollutant compared to CO<sub>2</sub> impact which is accordingly defined as a referent value. In that case the emission of greenhouse gases is presented as the equivalent emission of carbon dioxide (CO<sub>2</sub>-eq). If the removal of greenhouse gases occurs (e.g. the absorption of CO<sub>2</sub> at increase of wood stock in forests) than it refers to sinks of greenhouse gases and the amount is presented as a negative value. Table ES.2-1 shows the global warming potentials for particular gases.

*Table ES.2-1: Global warming potentials for certain gases (100- year time horizon)*

Gas	Global Warming Potential
Carbon dioxide (CO <sub>2</sub> )	1
Methane (CH <sub>4</sub> )	21
Nitrous oxide (N <sub>2</sub> O)	310
HFC-32	650
HFC-125	2800
HFC-134a	1300
HFC-143a	3800
CF <sub>4</sub>	6500
C <sub>2</sub> F <sub>6</sub>	9200
SF <sub>6</sub>	23900

### ES.3. OVERVIEW OF SOURCES AND SINK CATEGORY EMISSION ESTIMATES AND TRENDS

Total emission/removal of greenhouse gases for the period 1990-2005 and their trend in Sectors is given in table ES.3-1, while the contribution of the individual gases is given in table ES.3-2.

*Table ES.3-1: Emissions/removals of GHG by sectors for the period 1990-2005 (Gg CO<sub>2</sub> eq)*

Source	Emissions and removals of GHG (Gg CO <sub>2</sub> eq)							
	1990	1995	2000	2001	2002	2003	2004	2005
Energy	22504	16871	19287	20329	21531	22986	22442	22806
Industrial Processes	4206	2144	2874	2830	2746	2886	3252	3442
Solvent and Other Product Use	80	80	69	75	99	108	135	155
Agriculture	4464	3101	3130	3236	3276	3320	3561	3495
Waste	298	380	475	504	533	555	642	583
<b>Total emission (excluding net CO<sub>2</sub> from LULUCF)</b>	<b>31552</b>	<b>22576</b>	<b>25835</b>	<b>26974</b>	<b>28185</b>	<b>29853</b>	<b>30031</b>	<b>30481</b>
Removals (LULUCF)	-6281	-10629	-8012	-8880	-8537	-7878	-7948	-7779
<b>Total emission (including LULUCF)</b>	<b>25271</b>	<b>11947</b>	<b>17823</b>	<b>18094</b>	<b>19648</b>	<b>21976</b>	<b>22083</b>	<b>22702</b>

*Table ES.3-2: Emissions/removals of GHG by gases for the period 1990-2005 (Gg CO<sub>2</sub> eq)*

Source	Emissions and removals of GHG (Gg CO <sub>2</sub> eq)							
	1990	1995	2000	2001	2002	2003	2004	2005
Carbon dioxide (CO <sub>2</sub> )	23307	16877	19947	20945	22032	23501	23148	23567
Methane (CH <sub>4</sub> )	3247	2540	2548	2691	2746	2925	3015	2956
Nitrous oxide (N <sub>2</sub> O)	4017	3116	3317	3289	3358	3263	3680	3608
HFCs, PFCs and SF <sub>6</sub>	980	43	23	49	49	164	189	349
<b>Total emission (excluding net CO<sub>2</sub> from LULUCF)</b>	<b>31552</b>	<b>22576</b>	<b>25835</b>	<b>26974</b>	<b>28185</b>	<b>29853</b>	<b>30031</b>	<b>30481</b>
Removals (LULUCF)	-6281	-10629	-8012	-8880	-8537	-7878	-7948	-7779
<b>Total emission (including LULUCF)</b>	<b>25271</b>	<b>11947</b>	<b>17823</b>	<b>18094</b>	<b>19648</b>	<b>21976</b>	<b>22083</b>	<b>22702</b>

Figure ES.3-1 graphically represents the contribution of the individual sectors to total emissions and removals of the greenhouse gases. The largest contribution to the greenhouse gas emission in 2005 has the Energy Sector with 74.8 percent, followed by Agriculture with 11.5 percent, Industrial Processes with 11.3 percent, Waste with 1.9 percent and Solvent and Other product Use with 0.5 percent. This structure is with minor changes consistent through all the observed period from 1990 to 2005. In the year 2005 the amount of removed emissions of the greenhouse gases by CO<sub>2</sub> from the forestry sector was 25.5 percent.






Figure ES.3-1: Emissions and removals of the greenhouse gases in the Republic of Croatia by sectors in the period from 1990-2005 (Gg CO<sub>2</sub>-eq)

Energy sector, is the largest contributor to greenhouse gas emissions. In this sector, in the year 2005 the total energy consumption was 1.6 percent higher than in the former year 2004, whereat the total largest increase was in coal consumption (4.0 percent), which is also the most intensive energy-generating product from the point of view of CO<sub>2</sub> emission (92.7 t/TJ). The CO<sub>2</sub> emission from electric and heat power production in thermal power plants, public heating plants and public cogeneration plants was 4.6 millions of tons in 2005, representing 15.1 percents in total greenhouse emission in the Republic of Croatia.

Emission of CH<sub>4</sub> and N<sub>2</sub>O in the Agricultural sector is conditioned by different agricultural activities. For the emission of CH<sub>4</sub> the most important source is livestock farming (Enteric Fermentation). The number of cattle showed continuous decrease in the period from 1990 to 2000. As a consequence, this led to CH<sub>4</sub> emission reduction. In the year 2000, the number of cattle has started increasing and this trend was retained until 2005. The emission of N<sub>2</sub>O is considered as a direct emission from cultivation of agricultural soils, emission from the animal manure (Manure Management) and indirect emission. Likewise as the CH<sub>4</sub> emission, the increasing trend of N<sub>2</sub>O emission from 2000 is recorded due to increased use of mineral nitrogen fertilizers.

In Industrial Processes sector the key emission sources are Cement Production, Ammonia Production, Nitric Acid Production and Consumption of HFCs in Refrigeration and Air Conditioning Equipment, which all together contribute with 93.1 percent in total sectoral emission in 2005. The iron production in blast furnaces and aluminium production were ended in 1992, and ferroalloys production ended in 2002. The cement production in the period from 1997-2005 was constantly increasing. The aim of the producer is maximum use of the existing capacities which amounts about 3.1 millions of tons of clinker in total per year, whereas in the

year 2005, 2.9 millions of tons of clinker was produced. The ammonia production in 2005 was 1 percent lower in comparison to the previous year. Also, the nitric acid production in 2005 was 2.4 percent lower in comparison to 2004. The level of emissions from these sub-sectors strongly depends on consumer's demand for particular type of mineral fertilizer at the market.

Waste sector includes waste disposal, waste water management and waste incineration, whereas the waste disposal represents dominant CH<sub>4</sub> emission source from that sector in the Republic of Croatia. The emission depends on the amount and composition of municipal solid waste, management practices on-site including implementation of measures for collection and utilization of landfill gas. It is necessary to emphasize that in 2006, according to recommendations of the ERT, the First Order Decay (FOD) model was used for the first time, which replaced the former Tier 1 model. Application of more complex model led to significantly lower CH<sub>4</sub> emissions from the Waste sector during the all observed period, as it was expected. The results of the model show that the emissions from this sector continuously increases mainly as a consequence of the larger amounts of waste disposed at the landfills and absence of secondary measures for emission reduction at the landfills (flaring, electricity production from the landfill gas). It should be emphasized that the Waste sector contributes to total greenhouse gas emissions with 1.9 percent.

### ES.3.1. CARBON DIOXIDE EMISSION (CO<sub>2</sub>)

Carbon dioxide is the most significant anthropogenic greenhouse gas. As in the majority of countries, the most significant anthropogenic sources of CO<sub>2</sub> emissions in Croatia are the processes of fossil fuel combustion for electricity or/and heat production, transport and industrial processes (cement and ammonia production). The results of the CO<sub>2</sub> emission calculation in Croatia are presented in table ES.3-3.

*Table ES.3-3: CO<sub>2</sub> emission/removal by sectors from 1990-2005 (Gg CO<sub>2</sub>)*

<b>Sector</b>	<b>1990</b>	<b>1995</b>	<b>2000</b>	<b>2001</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>
Energy	20945	15540	17888	18814	19939	2135	20758	21107
Industrial processes	2283	1257	1930	2056	1994	2059	2255	2305
Solvent and Other Product Use	80	80	69	75	99	108	135	155
LULUCF	-6281	-10629	-8012	-8880	-8537	-7878	-7948	-7779
<b>Total CO<sub>2</sub> emission</b>	<b>23308</b>	<b>16877</b>	<b>19947</b>	<b>20945</b>	<b>2032</b>	<b>23501</b>	<b>23148</b>	<b>23567</b>
<b>Net CO<sub>2</sub> emission</b>	<b>17026</b>	<b>6249</b>	<b>11935</b>	<b>12065</b>	<b>13495</b>	<b>15624</b>	<b>15199</b>	<b>15788</b>

#### ES.3.1.1. Energy sector

This sector covers all the activities which include fossil fuel consumption and fugitive emission from fuels. Fugitive emission arises from production, transport, processing, storage and distribution of fossil fuels. The Energy sector is the main source of the anthropogenic greenhouse gas emission with share of 75 percent in total greenhouse gas emission. CO<sub>2</sub> emission from fuel combustion makes the largest part of it (more than 90 percent of emission in the Energy sector). Emission by sub-sectors is presented in table ES.3-4.

Table ES.3-4: CO<sub>2</sub> emission by sub-sectors from 1990-2005 (Gg CO<sub>2</sub>)

Source	1990	1995	2000	2001	2002	2003	2004	2005
Energy Industries	6823	5176	5882	6294	7213	7877	6771	6798
Manufacturing Industries & Constr.	6015	3489	3501	3646	3495	3608	4041	4134
Transport (Road & Off-Road)	4070	3392	4515	4613	4912	5287	5435	5655
Comm./Inst., Resid., Agr /For./Fish.)	3620	2785	3357	3574	3653	3880	3801	3830
Natural gas scrubbing (CGS Molve)	416	697	633	688	665	684	710	691
<b>Total CO<sub>2</sub> emission</b>	<b>20945</b>	<b>15540</b>	<b>17888</b>	<b>18814</b>	<b>19939</b>	<b>21335</b>	<b>20758</b>	<b>21107</b>

Emission calculation is based on fuel consumption data recorded in annual national energy balance, where the fuel consumption and supply is presented at the sufficient level of detail which enables more detailed calculation by sub-sectors in the framework of the formal IPCC methodology (i.e. Sectoral approach). Furthermore, the simplest method of the calculation was carried out (i.e. Reference approach) which takes into account only the total balance of fuel, without sub-sector analysis. The relative deviation of CO<sub>2</sub> emissions between sectoral and reference approach for Croatia is around 2.5 percent which is within the acceptable values (table ES.3-5).

Table ES.3-5: CO<sub>2</sub> emission comparison due to fuel combustion (Gg)

	1990	1995	2000	2001	2002	2003	2004	2005
Reference approach (Gg CO <sub>2</sub> )	20842	15140	17739	18543	19805	21171	20557	20921
Sectoral approach (Gg CO <sub>2</sub> )	20529	14843	17255	18126	19273	20651	20048	20416
<b>Relative Difference (%)</b>	<b>1.5</b>	<b>2.0</b>	<b>2.8</b>	<b>2.3</b>	<b>2.8</b>	<b>2.5</b>	<b>2.5</b>	<b>2.5</b>

Two energy most intensive sub-sectors are energy transformation (thermal power plants, heating plants, refineries and oil and gas field combustion) and manufacturing industry and construction. In the framework of the sub-sector Manufacturing Industry and Construction, the largest CO<sub>2</sub> emissions are the result of fuel combustion in construction material industry and than in iron and steel industry, non-metal industry, chemical industry, industry of pulp, paper and print, food and drink production, tobacco production etc. Furthermore, this sub-sector includes electricity and heat production in manufacturing industry for manufacturing processes.

Transport is also one of more important CO<sub>2</sub> emission sources. The largest part of the emission arises from Road transportation (86 – 94 percent depending on the year) followed by railways and domestic civil aviation and navigation. Emission from fuel sold for the international aviation and marine transportation is reported separately and it's not included in total national emission balance. In the year 2005, emission from Transport sector contributed with 18.4 percent to total greenhouse gas emission.

Biomass combustion (fuel wood and waste wood, biodiesel, biogas) also results in greenhouse gas emissions. CO<sub>2</sub> emission from biomass is not included in balance according the guidelines, due to assumption that life-cycle CO<sub>2</sub> emitted is formerly absorbed for the growth of biomass. Sinks or CO<sub>2</sub> emissions resulted in change of forest biomass is calculated in sector Land Use, Land-Use Change and Forestry.

Fugitive greenhouse gas emission from coal, liquid fuels and natural gas, resulted from exploration of minerals, production, processing, transport, distribution and activities during mineral use is also included in this sector. Although this emission is not characteristic for CO<sub>2</sub>, yet for CH<sub>4</sub>, there is a CO<sub>2</sub> emission present during the process of scrubbing of natural gas in Central Gas Station Molve. The natural gas exploited on Croatian fields is rich in carbon dioxide (more than 15 percent) and before the natural gas is distributed in commercial gas pipeline it is necessary to remove the CO<sub>2</sub> (scrubbing) so that the maximum volume share of CO<sub>2</sub> in natural gas is 3 percent. Emission assessment during the removal is based on material balance method and amounts up to 4 percent of the total CO<sub>2</sub> emission in Energy sector.

### ES.3.1.2. Industrial processes

The greenhouse gas emission is a by-product in various industrial processes where the raw material is chemically transformed in final product. Industrial processes where the contribution to CO<sub>2</sub> emission is identified as relevant are: cement production, lime, ammonia, ferroalloys production as well as limestone use and soda ash use in various industrial activities.

General methodology used for emission calculation from industrial processes, recommended by the Convention, includes the product of annual produced or consumed amount of a product or material with appropriate emission factor per unit of this production or consumption. Annual production or consumption data for particular industrial processes are extracted, in most cases, from monthly industrial reports published by Central Bureau of Statistics. Certain activity data was collected from voluntary survey of manufacturers. The results of the CO<sub>2</sub> emission in industrial processes are shown in table ES.3-6.

*Table ES.3-6: CO<sub>2</sub> emission from Industrial Processes for the period from 1990-2005 (Gg CO<sub>2</sub>)*

<b>Source</b>	<b>1990</b>	<b>1995</b>	<b>2000</b>	<b>2001</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>
Cement production	1047.5	592.4	1233.2	1395.0	1361.1	1371.5	1458.7	1495.0
Lime production	159.8	62.3	124.3	143.5	164.0	161.0	174.3	196.4
Limestone and dol. use	43.2	11.2	8.4	9.2	9.6	11.8	11.5	12.1
Soda ash prod. and use	25.7	14.4	11.0	12.4	12.2	14.7	16.5	17.2
Ammonia production	491.6	462.9	525.3	425.9	383.7	431.8	522.6	511.2
Ferroalloys production	194.5	33.9	20.5	0.5	0.0	0.0	0.0	0.0
Aluminum production	111.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0
<b>Total CO<sub>2</sub> emission</b>	<b>2073.7</b>	<b>1177.0</b>	<b>1922.7</b>	<b>1986.5</b>	<b>1930.7</b>	<b>1990.8</b>	<b>2183.6</b>	<b>2231.9</b>

The most significant CO<sub>2</sub> industrial processes emission sources are cement production and ammonia production. The CO<sub>2</sub> emission from cement production contributes, depending on the year, with 22 to 51 percent of total CO<sub>2</sub> emission from industrial sector, and the CO<sub>2</sub> emission from ammonia production contributes with 12 to 24 percent of the total sectoral emission. Generally, CO<sub>2</sub> emissions from industrial processes declined from 1990 to 1995, due to the decline in industrial activities. However in the next period from 1996-2005 the emission was increasing to the level reported in 1990.

The quantity of the CO<sub>2</sub> emitted during cement production is directly proportional to the lime content of the clinker. Therefore, the CO<sub>2</sub> emissions are calculated using an emission factor, in tones of CO<sub>2</sub> released per tone of clinker produced, to the annual clinker output corrected with the fraction of clinker that is lost from the kiln in the form of Cement Kiln Dust (CKD). The emission factor and correction factor for CKD is determined according to *Revised 1996 IPCC Guidelines* and *Good Practice Guidance*. Country specific emission factor and CKD was estimated using data from individual plants. The activity data for clinker production were collected from survey of cement manufacturers and cross-checked with cement production data from monthly industrial reports published by Central Bureau of Statistics.

Emission of CO<sub>2</sub> from ammonia production is stehiometrically determined based on carbon content in natural gas. One part of the CO<sub>2</sub> produced in ammonia production is further used as feedstock in urea production, i.e. mineral fertilizer. Emission of intermediately bound CO<sub>2</sub> occurs during the use of urea as a fertilizer in agriculture. However, according to IPCC methodology this approach is not distinguished. Therefore, the total CO<sub>2</sub> emission from natural gas used as a feedstock for ammonia production is reported in this sub-sector.

#### **ES.3.1.3. CO<sub>2</sub> removals**

According to Forest Management Area Plan of the Republic of Croatia (1996-2005), the forests and the forest land cover 43.5 percent of the total surface area. By its origin, approximately 95 percent of the forests in Croatia were formed by natural regeneration and the 5 percent of the forests are grown artificially. Out of the total surface area occupied by forests and the forest land, 2089607 ha (84 percent) is the forest-covered area, 327630 ha (13 percent) is non forest land, and 74063 ha (3 percent) is bare unproductive and unfertile forestland.

The Republic of Croatia reports only CO<sub>2</sub> emissions related with data for changes in the forest and other woody biomass stocks. For other segments in the sector Land Use, Land-Use Change and Forestry like forest land and grassland converted to cropland or other land, and carbon change in soil, there were no reliable input data available.

Annual carbon increment in Croatian forests is 9643000 m<sup>3</sup>. Increment is an increase in forest wood stock over a certain time period. It is calculated as annual, periodical and average increment. Different methods have been developed in forest management to identify the forest increment. The methods mostly used in Croatia are a check method and a method of bore-spills. Different methods of forest cultivation can make the increment larger both in terms of their quantity and quality. A described cutting is a part of the forest wood stock planned for commercial cutting over a time period (1 year, 10 years, 20 years) expressed in wood stock (m<sup>3</sup>, m<sup>3</sup>/ha) or in an area (ha). In order to satisfy the basic principal of forest management and principles of sustainability the described cut should not be larger than the increment value. The problem of deforestation in Croatia doesn't exist. According to present data the total forest area has not been reduced in the last 100 years. Clear-cuttings, as a measure of forest recovery are prohibited, according to "Forestry Act" of the Republic of Croatia from 2005, and natural regeneration, is the main way of forest renewal.

The methodology used for CO<sub>2</sub> removal calculation is taken from the IPCC and it is based on data on annual increment and cutting. The figure ES.3-2 shows the CO<sub>2</sub> emission removal trend in the forestry sector.




Figure ES.3-2: CO<sub>2</sub> emission removal in forestry sector from 1990-2005 (Gg CO<sub>2</sub>)

### ES.3.2. METHANE EMISSION (CH<sub>4</sub>)

The major sources of methane (CH<sub>4</sub>) emission are fugitive emission from production, processing, transportation and activities related with fuel use in Energy sector, Agriculture and Waste Disposal on Land. In table ES.3-7, sectoral and total CH<sub>4</sub> emissions are reported.

Table ES.3-7: CH<sub>4</sub> emission in Croatia in the period from 1990-2005 (Gg CH<sub>4</sub>)

Source	1990	1995	2000	2001	2002	2003	2004	2005
Energy	68.5	58.8	59.4	64.5	67.0	68.5	69.7	70.4
Industrial Processes	0.8	0.4	0.3	0.3	0.3	0.3	0.3	0.3
Agriculture	74.8	48.0	43.0	43.6	42.7	48.6	47.4	46.
Waste	10.5	13.7	18.6	19.7	20.8	22.0	26.2	23.4
<b>Total CH<sub>4</sub> emission</b>	<b>154.6</b>	<b>120.9</b>	<b>121.4</b>	<b>128.1</b>	<b>130.8</b>	<b>139.3</b>	<b>143.6</b>	<b>140.8</b>

Fugitive methane emission is mainly the result of exploration, production, processing, transportation and distribution of natural gas (about 97 percent). The fugitive emission from oil accounts with about 1 percent; venting and flaring of gas/oil production accounts with approximately 2 percent. In 1999 by closing of the coal mines in Istra, large amount of fugitive emissions arising from the exploration, processing and transportation of coal, were avoided.

In the Agricultural sector there are two significant methane emission sources present: enteric fermentation in the process of digestion of ruminants (dairy cows represent the major source) and different activities related with storage and use of organic fertilizers (manure management). The total methane emission for domestic animals is being calculated as a sum of emission from enteric fermentation and emission related to manure management.

Methane emission from solid waste disposal sites (SWDSs) is a result of anaerobic decomposition of organic waste by methanogenic bacteria. The amount of methane emitted during the process of decomposition is directly proportional to the fraction of degradable organic carbon (DOC) which is defined as carbon content in different types of organic biodegradable wastes. In Croatia, more than million tons of municipal solid waste is produced annually and the average composition of its biodegradable part is: paper and textile (21-22 percent), garden and park waste (18-19 percent), food waste (23-24 percent), wood waste and straw (3 percent). As for the wastewater handling in Croatia, aerobic biological process is used mostly in wastewater treatment. Anaerobic process is applied in some industrial wastewater treatment. Total amount of gas is flared in these treatments, and therefore all methane from gas is oxidized to carbon dioxide and water vapour. Disposal of domestic and commercial wastewater, particularly in rural areas where systems such as septic tanks are used, are partly anaerobic without flaring, which results with CH<sub>4</sub> emissions.

### ES.3.3. NITROUS OXIDE EMISSION (N<sub>2</sub>O)

The most important sources of N<sub>2</sub>O emissions in Croatia are agricultural activities, nitric acid production, but as well, the N<sub>2</sub>O emissions occur in energy sector and waste management. In table ES.3-8 the N<sub>2</sub>O emission is reported according to sectors.

*Table ES.3-8: N<sub>2</sub>O emission in Croatia for the period from 1990-2005 (Gg N<sub>2</sub>O)*

<b>Source</b>	<b>1990</b>	<b>1995</b>	<b>2000</b>	<b>2001</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>
Energy	0.4	0.3	0.5	0.5	0.6	0.7	0.7	0.7
Industrial Processes	3.0	2.7	2.8	2.3	2.3	2.1	2.6	2.5
Agriculture	9.3	6.8	7.2	7.5	7.7	7.4	8.3	8.1
Waste	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
<b>Total N<sub>2</sub>O emission</b>	<b>13.0</b>	<b>10.1</b>	<b>10.700</b>	<b>10.6</b>	<b>10.8</b>	<b>10.5</b>	<b>11.9</b>	<b>11.6</b>

In the framework of Agricultural sector, three N<sub>2</sub>O emission sources are determined: direct N<sub>2</sub>O emission from agricultural soils, direct N<sub>2</sub>O emission from livestock farming and indirect N<sub>2</sub>O emission induced by agricultural activities. The largest emission is a result of direct emission from agricultural soils and it includes total amount of carbon which occurs in systems of plant cultivation. According to IPCC methodology, the mineral nitrogen, nitrogen from organic fertilizers, amount of nitrogen in fixing crops, amount of nitrogen emitted due to decomposition of crop waste and amount of nitrogen which is released from crop residue mineralization and soil nitrogen mineralization due to cultivation of histosols, are separately analyzed.

In the sector Industrial Processes the N<sub>2</sub>O emission occurs in nitric acid production, which is used as a raw material in nitrogen mineral fertilizers. In the framework of the N<sub>2</sub>O reduction measure analysis, the possibility for application of non-selective catalytic reduction device was considered, whereby the nitric acid production influence on N<sub>2</sub>O emissions would be practically eliminated.

In Energy sector the emission was calculated on the basis of fuel consumption and adequate emission factors (IPCC). The N<sub>2</sub>O emission increase in Energy sector is the consequence of

greater use of three-way catalytic converters in road transport motor vehicles, which have about 30 times greater  $\text{N}_2\text{O}$  emission comparing to vehicles without a catalytic converter.

$\text{N}_2\text{O}$  emission from the Waste sector indirectly occurs from the human sewage. It is calculated on the basis of the total number of inhabitants and annual protein consumption per inhabitant. Data on the annual per capita Protein intake value were obtained by the FAOSTAT Statistical Database. Adjustment method (extrapolation) has been used for calculation of insufficient data.

#### ES.3.4. HALOGENATED CARBONS (HFCs, PFCs) AND $\text{SF}_6$ EMISSIONS

Synthetic greenhouse gases include halogenated carbons (HFCs and PFCs) and sulphur hexafluoride ( $\text{SF}_6$ ). Although on an absolute scale their emissions are not great, due to their high global warming potential (GWP) their contribution to global warming is considerable. According to survey carried out among major agents, users and consumers of these gases, information related to import and export of HFCs (provided by the Ministry of Environmental Protection, Physical Planning and Construction) was used for emission calculation which is presented in Gg of  $\text{CO}_2$  eq and showed on the figure ES.3-3.




Figure ES.3-3: Halogenated carbons emission in the period from 1990-2005 (Gg  $\text{CO}_2$ -eq)

#### ES.4. EMISSION OF INDIRECT GREENHOUSE GASES

The photochemically active gases, carbon monoxide ( $\text{CO}$ ), oxides of nitrogen ( $\text{NO}_x$ ) and non-methane volatile organic compounds (NMVOCs) indirectly contribute to the greenhouse gas effect. These are generally called indirect greenhouse gases or ozone precursors, because they are involved in creation and degradation of ozone which is also one of the greenhouse gases. Sulphur dioxide ( $\text{SO}_2$ ), as a precursor of sulphate and aerosols, is believed to contribute negatively to the greenhouse effect. The calculation of aggregated results for the emissions of indirect gases in the period 1990-2005 are given in table ES.4-1.



Table ES.4-1: Emissions of ozone precursors and SO<sub>2</sub> by different sectors (Gg)

Gas	Emissions (Gg)							
	1990	1995	2000	2001	2002	2003	2004	2005
<b>NO<sub>x</sub> Emission</b>	<b>83.9</b>	<b>57.7</b>	<b>68.4</b>	<b>69.4</b>	<b>68.1</b>	<b>68.7</b>	<b>67.4</b>	<b>73.3</b>
Energy Industries	13.6	10.3	12.0	11.6	13.2	13.8	11.2	12.0
Manufacturing Ind. & Construction	17.5	8.9	9.7	10.6	10.2	9.7	11.8	16.3
Transport	37.0	29.8	33.3	33.0	31.0	31.1	30.6	31.3
Other Energy (fuel combustion)	15.0	8.1	12.8	13.6	13.1	13.6	13.2	13.1
Fugitive Emission from Fuels	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.3
Industrial Processes	0.4	0.3	0.3	0.3	0.3	0.2	0.3	0.3
<b>CO Emission</b>	<b>502.1</b>	<b>339.4</b>	<b>378.0</b>	<b>322.9</b>	<b>305.8</b>	<b>321.5</b>	<b>308.5</b>	<b>329.6</b>
Energy Industries	1.5	1.0	1.2	1.0	1.0	1.4	1.2	0.9
Manufacturing Ind. & Construction.	40.4	41.3	37.8	38.6	37.9	38.0	41.6	32.7
Transport	252.5	176.6	188.7	165.0	151.0	137.5	125.4	159.2
Other Energy (fuel combustion)	204.0	116.7	146.5	115.2	113.0	141.4	136.4	133.0
Fugitive Emission from Fuels	0.6	0.5	0.5	0.4	0.4	0.4	0.5	0.5
Industrial Processes	3.1	3.3	3.3	2.7	2.5	2.8	3.4	3.3
<b>NM VOC Emission</b>	<b>144.8</b>	<b>141.8</b>	<b>218.0</b>	<b>177.7</b>	<b>290.9</b>	<b>416.3</b>	<b>481</b>	<b>437.6</b>
Energy Industries	0.3	0.2	0.3	0.3	0.3	0.3	0.3	0.3
Manufacturing Ind. & Construction	1.7	1.4	1.4	1.5	1.4	1.7	1.9	3.3
Transport	40.9	30.0	32.9	27.4	25.6	22.1	19.4	18.5
Other Energy (fuel combustion)	12.1	6.9	9.0	7.5	7.3	8.8	8.5	8.3
Fugitive Emission from Fuels	8.2	7.8	9.7	10.4	10.8	10.5	9.7	9.1
Industrial Processes	81.6	95.5	164.7	130.6	245.5	372.9	441.2	398.1
Solvent Use	27.4	27.4	23.4	25.5	33.8	36.7	46.1	52.9
<b>SO<sub>2</sub> Emission</b>	<b>175.2</b>	<b>81.5</b>	<b>69.9</b>	<b>69.9</b>	<b>75.6</b>	<b>74.2</b>	<b>61.6</b>	<b>68.0</b>
Energy Industries	78.5	39.0	25.4	26.0	23.3	35.7	25.7	32.8
Manufacturing Ind. & Construction	55.8	24.7	22.6	24.9	29.9	15.6	11.8	10.3
Transport	5.3	3.5	6.0	4.9	6.3	7.4	7.9	8.5
Other Energy (fuel combustion)	23.9	4.7	6.5	6.1	8.0	7.5	6.9	6.6
Fugitive Emission from Fuels	6.4	5.0	4.8	4.5	4.5	4.5	4.7	4.6
Industrial Processes	5.3	4.6	4.6	3.5	3.6	3.5	4.6	5.2

## 1. INTRODUCTION

### 1.1. BACKGROUND INFORMATION ON GHG INVENTORIES AND CLIMATE CHANGE

The Republic of Croatia became a party to the United Nations Framework Convention on Climate Change (UNFCCC) in 1996, by the Act on Ratification (Official Gazette No. 2/96). Pursuant to that Decree, the Republic of Croatia has under Article 22 of the UNFCCC undertaken the commitments outlined in Annex I as a country with economy in transition. Croatia has thus committed itself to maintain greenhouse gas emissions at their 1990 level. The Republic of Croatia is also a signatory of the Kyoto Protocol. Upon its ratification (Official Gazette – International agreement No. 05/07) and entering into force, Croatia has committed to reduce its greenhouse gas emissions by 5 percent in relation to the base year, over the commitment period from 2008 to 2012.

One of the commitments outlined in Article 4.1 of the UNFCCC is that Parties are required to *develop, periodically update, publish and make available to the Conference of the Parties, in accordance with Article 12, national inventories of anthropogenic emissions by sources and removals by sinks of all greenhouse gases not controlled by the Montreal Protocol, using comparable methodologies to be agreed upon by the Conference of the Parties.*

Furthermore, Article 5.1 of the Kyoto Protocol requires that *each Party included in Annex I shall have in place, no later than one year prior to the start of the first commitment period, a national system for the estimation of anthropogenic emissions by sources and removals by sinks of all greenhouse gases not controlled by the Montreal Protocol.* A national system includes all institutional, legal and procedural arrangements made within a Party included in Annex I for estimating anthropogenic emissions by sources and removals by sinks of all greenhouse gases not controlled by the Montreal Protocol, and for reporting and archiving inventory information.

The Republic of Croatia is also a country which is currently in the process of accession to the EU. Accession is conditioned by the harmonization, adoption and implementation of the entire *acquis communautaire*, i.e. the body of legislation and rules already implemented in the EU. This process is very complex and requires changes that are systemic in its nature particularly in institutional and legislative sphere. As a future EU member state, Croatia will have to implement legislation concerning a mechanism for monitoring Community greenhouse gas emissions and for implementing the Kyoto Protocol, which also stipulates establishment of mechanism for monitoring emissions by sources and removals by sinks of greenhouse gases, evaluating progress towards meeting commitments in respect of these emissions and for implementing the UNFCCC and the Kyoto Protocol, as regards national programmes, inventories, national system and registries.

In compliance with the UNFCCC reporting requirements, Croatia has submitted national emission inventory reports on an annual basis since 2001. In this NIR, the inventory of the emissions and removals of the greenhouse gases is reported for the period from 1990 to 2005. The NIR is prepared in accordance with the UNFCCC reporting guidelines on annual Inventories as adopted by the COP by its Decision 18/CP.8. The methodologies used in the

calculation of emissions are based on the *Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories (IPCC Guidelines)* and the *IPCC Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories (IPCC Good Practice Guidance)* prepared by the Intergovernmental Panel on Climate Change (IPCC). As recommended by the IPCC Guidelines country specific methods have been used where appropriate and where they provide more accurate emission data. The important part of the inventory preparation is uncertainty assessment of the calculation and verification of the input data and results, all this with the aim to increase the quality and reliability of the calculation.

Furthermore, since the introduction of annual technical reviews of the national inventories by experts review teams (ERT), Croatia has undergone three reviews so far, in-country review in 2004 and centralized reviews in 2005 and 2006. Issues recommended by the ERT have been included in this report as far as possible.

The calculation includes the emissions which are the result of anthropogenic activities and these include the following greenhouse gases: carbon dioxide (CO<sub>2</sub>), methane (CH<sub>4</sub>), nitrous oxide (N<sub>2</sub>O), halogenated carbons (HFCs, PFCs) and sulphur hexafluoride (SF<sub>6</sub>) and indirect greenhouse gases: carbon monoxide (CO), oxides of nitrogen (NO<sub>x</sub>), non-methane volatile organic compounds (NMVOCs) and sulphur dioxide (SO<sub>2</sub>). The greenhouse gases covered by Montreal Protocol on the pollutants related to ozone depletion (freons) are reported in the framework of this protocol and therefore are excluded from this Report.

Greenhouse gas emission sources and sinks are divided into six main sectors: Energy, Industrial Processes, Solvent and Other Product Use, Agriculture, Land Use, Land-Use Change and Forestry and Waste. Generally, the methodology for emission calculation could be described as a product of the particular economic activity (e.g. fuel consumption, cement production, number of animals, increase of wood stock etc.) with corresponding emission factors. The use of specific national emission factors is recommended wherever possible and justified, whereas on the contrary, the methodology gives typical values of emission factors for all relevant activities of the particular sectors.

## **1.2 BRIEF DESCRIPTION OF THE INSTITUTIONAL ARRANGEMENT FOR INVENTORY PREPARATION**

Ministry of Environmental Protection, Physical Planning and Construction (MEPPPC) is a national focal point for the UNFCCC, responsible for overall inventory management activities including organization of the institutions responsible for and involved in preparing the national inventory, coordination between these institutions and communication to the UNFCCC secretariat. It is expected that MEPPPC will delegate full scope or part of these tasks to Croatian Environment Agency (CEA) in the near future.

EKONERG - Energy Research and Environmental Protection Institute is an inventory focal point authorized by the MEPPPC and contractually obliged for the preparation of annual emission inventories, which include compilation of national inventory report, archiving of relevant data, documentation of activity data, emission factors and used methods, validation of data, and verification of inventory estimates.

There are few key collaborating institutions in the inventory preparation process, mainly activity data providers, including Energy Institute "Hrvoje Požar" (EIHP) which prepares national energy balance on annual basis, Croatian Bureau of Statistics (CBS) and Croatian Center for Vehicles (CCV). These institutions are not formally/contractually bound to provide data to national focal point or national executive institutions but rather they provide these data on voluntary basis or through regular statistical publications.

One of the goals of the National GHG inventory improvement strategy is to prepare secondary legislation which will stipulate responsibilities and mandates for national institutions involved in inventory preparation process.

### **1.3. BRIEF DESCRIPTION OF THE PROCESS OF INVENTORY PREPARATION**

The important prerequisite for the efficient data management system and inventory preparation is clearly defined organization, authority and responsibility of the abovementioned institutions participating in the process of inventory preparation, which consists of number of steps in data collection and processing, calculating, control and verification of emission calculation, documentation, archiving and reporting to competent international institutions. It can be stated that Croatia, respectively the Ministry of Environmental Protection, Physical Planning and Construction, as a competent authority for inventory preparation uses, in organizational sense, decentralized model in which it transfers the authorization for preparation of individual tasks in process of inventory preparation on collaborative institutions which are partially public or governmental and partially in private ownership.

The main data sources for greenhouse gas emission calculation are Energy Institute "Hrvoje Požar" which prepares the national energy balance, Central Bureau of Statistics which collects data on raw materials and products for activities defined at National classification of economic activities on grounds of programme of statistical research, Croatian Center for Vehicles and Ministry of Internal Affairs which possesses data bases on road and off-road vehicles and Ministry of Agriculture, Forestry and Water Management which possesses data on forest-covered area. For the inventory preparation, the data gathered through questionnaires directly from individual emission sources and other scientific or professional institutions are also being used either for calculation or control of data obtained from the official publications. EKONERG - Energy Research and Environmental Protection Institute is executive institution, responsible for data collection, emission calculation and preparation of annual greenhouse gas emission inventories according to contract with Ministry of Environmental Protection, Physical Planning and Construction. It should be noted that the financing of the inventory in the past period was mainly provided through programme LIFE – Third Countries and GEF of the European Commission. The system organized in that way has certain advantages primarily in a sense of the efficient use of the existing resources, but on the other hand there are certain weaknesses of this system regarding middle-term and long-term planning and inventory improvement.

Considering the routine and need for sustainable system of greenhouse gas emission monitoring, and commitment for establishment of the National System for the purposes of the Kyoto Protocol, the Ministry of Environmental Protection, Physical Planning and Construction under Article 46 of the Air Protection Act (Official Gazette No. 178/2004) has started in July

2006 the process of preparation of the Regulation on the Greenhouse Gases Emissions Monitoring in the Republic of Croatia, which is confirmed in January 2007 (Official Gazette No. 1/2007). It should improve existing system of greenhouse gas emission monitoring and reporting in accordance with the requirements of the Kyoto protocol and relevant legislation of the EU. It is expected that the Regulation will enter into force in 2008.

#### 1.4. BRIEF DESCRIPTION OF METHODOLOGIES AND DATA SOURCES USED

The IPCC methodology from *Revised 1996 IPCC Guidelines for National GHG Inventories* and *Good Practice Guidance and Uncertainty Management in National GHG Inventories*, recommended by the UNFCCC, were used to calculate greenhouse gas emissions. This methodology covers following gases, which are result of anthropogenic activities: CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O, HFCs, PFCs, SF<sub>6</sub>, CO, NO<sub>x</sub>, NMVOCs, and SO<sub>2</sub>. Carbon dioxide (CO<sub>2</sub>), methane (CH<sub>4</sub>), and nitrous oxide (N<sub>2</sub>O) are principal greenhouse gases and though they occur naturally in the atmosphere, their recent atmospheric build-up appears to be largely the result of human activities. Synthetic gases such as halogenated hydrocarbons (PFCs, HFCs) and sulphur hexafluoride (SF<sub>6</sub>) are also considered as greenhouse gases and they are solely the result of human activities. The methodology does not include the CFCs which are the subject of the Montreal Protocol. In addition, there are other photochemically active gases such as carbon monoxide (CO), oxides of nitrogen (NO<sub>x</sub>) and non-methane volatile organic compounds (NMVOCs) that, although not considered as greenhouse gases, contribute indirectly to the greenhouse effect in the atmosphere. These are generally referred to as ozone precursors, because they participate in the creation and destruction of tropospheric and stratospheric ozone (which is also GHG). Sulphur dioxide (SO<sub>2</sub>), as a precursor of sulfate and aerosols, is believed to exacerbate the greenhouse effect because the creation of aerosols removes heat from the environment.

The emission estimates are divided into following IPCC sectors: Energy, Industrial Processes, Solvent and Other Product Use, Agriculture, Land Use, Land-Use Change and Forestry and Waste. Generally, methodology applied to estimate emissions involves the product of activity data (e.g. fuel consumption, cement production, wood stock increment and so forth) and an associated emission factor. The use of country-specific emission factors, if available, is recommended but these cases should be based on well-documented research. Otherwise, the *Revised 1996 IPCC Guidelines* provides a default values for emission factors.

Data sources for GHG inventory preparation are presented in the Table 1.4-1, but more detailed information is given in sectoral chapters.

Table 1.4-1: Data sources for GHG inventory preparation

CRF Sector/Sub-sector	Type of data	Source of data
<b>Energy</b>	Energy balance data Fuel consumption and fuel characteristic data for every boiler or gas turbine Database of motor vehicles Aggregated data about number of motor vehicles Fuel characteristic data Natural gas processed (scrubbed), CO <sub>2</sub> content before scrubbing and CO <sub>2</sub> emission	<ul style="list-style-type: none"> <li>▪ Energy Institute "Hrvoje Požar"</li> <li>▪ HEP - Croatian Electric Utility Company</li> <li>▪ Croatian Centre for Vehicles</li> <li>▪ Central Bureau of Statistics</li> <li>▪ INA - Oil and Gas Company</li> <li>▪ Central Gas Station MOLVE (INA)</li> </ul>
<b>Industrial Processes</b>	Activity data on production/consumption of material for particular industrial process  Data on consumption and composition of natural gas in ammonia production	<ul style="list-style-type: none"> <li>▪ Central Bureau of Statistics, Department of Manufacturing and Mining</li> <li>▪ Voluntary survey of manufacturers</li> <li>▪ Ministry of Environmental Protection, Physical Planning and Construction</li> <li>▪ Voluntary survey of ammonia manufacturer (Petrokemija Kutina)</li> <li>▪ Central Bureau of Statistics, Department of Manufacturing and Mining</li> </ul>
<b>Solvent and Other Product Use</b>	Activity data on production for particular source category and number of inhabitants	<ul style="list-style-type: none"> <li>• Central Bureau of Statistics, Department of Manufacturing and Mining</li> </ul>
<b>Agriculture</b>	Livestock number  Production of N-fixing crops and non N-fixing crops Area of histosols Synthetic fertiliser	<ul style="list-style-type: none"> <li>▪ Report on agricultural production and Inventory of agriculture (Central Bureau of Statistics)</li> <li>▪ Report on agricultural production (Central Bureau of Statistics)</li> <li>▪ Expert judgment</li> <li>▪ Expert judgment, Fertiliser company (Petrokemija Kutina)</li> </ul>
<b>LULUCF</b>	Forest area and Commercial harvest	<ul style="list-style-type: none"> <li>▪ Ministry of Agriculture, Forestry and Water Management</li> <li>▪ Public enterprise "Hrvatske šume"</li> </ul>
<b>Waste</b>	The total annual municipal solid waste disposed to different types of SWDSs  Data on waste composition	<ul style="list-style-type: none"> <li>▪ Report of Estimation of the Quantities of Municipal Solid Waste in Croatia (1990-1998 &amp; 1998-2010)</li> <li>▪ Report of Environment Condition, Ministry of Environmental Protection, Physical Planning and Construction</li> <li>▪ Interpolation/extrapolation methods</li> <li>▪ Waste Management Strategy</li> <li>▪ Waste Management Plan</li> <li>▪ Report: The basis for methane emissions estimation in Croatia (1990-1998), part B: Data on Municipal Solid Waste in Croatia 1990-1998</li> </ul>

## 1.5. BRIEF DESCRIPTION OF KEY CATEGORIES

According to the *Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories* key categories are those which represent 95% of the total annual emissions in the last reported year or belonging to the total trend, when ranked from contributing the largest to smallest share in annual total and in the trend.

The analysis is based on the contribution of CO<sub>2</sub> equivalents from different sources and sinks on the sectoral level. The recommended IPCC categories as well as the categories recommended in *Good Practice Guidance for Land Use, Land-Use Change and Forestry* to be assessed in the key category analysis are presented in Table A1-1 of the Annex 1. Furthermore, Croatian experts determined certain sub-categories which are particularly significant, such as CO<sub>2</sub> Emission from Natural Gas Scrubbing (also shown in Table A1-1 of the Annex 1). Additionally, CO<sub>2</sub> emissions from Solvent and Other Product Use and Other not-specified Non Energy Use (NEU) categories were for the first time included in the key categories analysis.

The results of the Level Assessment including/excluding LULUCF are shown in Table A1-2 and Table A1-3 respectively, with the key categories shaded. The key categories are sorted in descending order of magnitude and the cumulative total is included in the final column of the table.

The results of the Trend Assessment including/excluding LULUCF are shown in Table A1-4 and Table A1-5 respectively, with the key categories shaded. The key categories are sorted in descending order of magnitude, and the cumulative total is shown in the final column of the table.

Finally, the results of the Key Category analysis including/excluding LULUCF are summarized by sector and gas in Table A1-6 and A1-7 respectively. The tables indicate whether the key category arises from the level assessment or the trend assessment or both level and trend assessment.

Some changes in the Key Categories occurred in this NIR in relation to NIR 2006, e.g. the CO<sub>2</sub> Emissions from Solvent and Other Product Use and CO<sub>2</sub> Emissions from Non-energy Use were included in calculation. Furthermore, some key categories were excluded in 2006 submission due to an error in determining the key categories from the correctly calculated sheets. This refers to the ERT recommendation to include all the sources that add up to over 95 per cent of total. These changes are shown in Table A1-8.

## 1.6. INFORMATION ON THE QA/QC PLAN INCLUDING VERIFICATION AND TREATMENT OF CONFIDENTIALITY ISSUES

### 1.6.1. QA/QC PLAN AND PROCEDURES

According to *Good Practice Guidance and Uncertainty Management in National GHG Inventories*, QA/QC plan is an internal document to organize, plan, and implement QA/QC activities. Croatia has drafted QA/QC framework plan following the recommendations from document *Quality Assurance and Quality Control Plan, Samples and Manual for Development* which was prepared under regional UNDP/GEF project Capacity building for improving the quality of GHG inventories (RER/01/G31), Table A6-1, Annex 6.

Generally, QA/QC framework plan follows the proposed cycle of activities including:

- Development and approval of QA/QC plan (QA/QC manager and Inventory team leader);
- Data checking and inventory reviewing activities (QA/QC manager and sectorial experts);
- Compilation of findings (QA/QC manager);
- Recommendations for corrective actions (QA/QC manager);
- Implementing and reporting corrective actions (sectorial experts);
- Reporting (QA/QC manager).

Quality control activities are focused on following elements of inventory preparation process:

- Activity data gathering and handling activities;
- Activity data documentation;
- Choice of emission factors and emissions estimation.

Quality assurance activities should include one or more of the following activities: peer review by third party, public review and technical review by the UNFCCC expert review team.

QA/QC framework plan is presented in Annex 6.

Although inventory team has not prepared written general and source-specific QC procedures for each QC activity outlined in *Good Practice Guidance and Uncertainty Management in National GHG Inventories* for each IPCC sector, a *Sectorial methodological guidelines* were prepared in order to support inventory team with comprehensive guidelines for choice of methodology, emission factors and activity data, uncertainty estimates, QA/QC activities, reporting and documentation and inventory improvement plan. These guidelines are in accordance with *IPCC Guidelines* and *Good Practice Guidance* but also contain detail information on national circumstances particularly related to status of activity data, data gaps and short- and medium-term actions for improvement of the inventory.

For the purposes of transparency of the emission calculation, inventory team has continued with preparation of *Inventory Data Record Sheets* which were introduced in 2001 submission and which contain details of the person and/or organization responsible for an emission estimate, the primary or secondary sources of activity data and emission factors used, the methodology



applied, data gaps, ways to cross-check, suggestion for future improvement in the estimates and relevant bibliographic references. The information provided in *Inventory Data Record Sheets* is available for each source category and for the entire time-series. An example of Inventory Data Record Sheet for 2005 in Waste sector is presented in Annex 6, Table A6-2.

During the preparation of the NIR a number of checks were carried out by sector experts related to completeness, consistency, comparability, recalculation and uncertainty of activity data, emission factors and emission estimates. The details on these issues are elaborated in the NIR and the CRF.

Finally, before submitting this NIR an audit has been carried out by designated QA/QC manager. The audit covered all IPCC sectors in the NIR with purpose to check which quality control elements, both general and specific, as defined in the *IPCC Good Practice Guidance*, are already implemented by sector experts and which improvements and corrective actions should be carried out in the future submissions.

### 1.6.2. VERIFICATION AND CONFIDENTIALITY ISSUES

The verification process of calculation is aimed at the improvement of the input quality and identification of the calculation reliability. The *IPCC Guidelines* recommend that inventories should be verified through the use of a set of simple checks for completeness and accuracy, such as checks for arithmetic errors, checks of country estimates against independently published estimates, checks of national activity data against international statistics and checks of CO<sub>2</sub> emissions from fuel combustion calculated using sectoral methods with the IPCC Reference Approach. Further verification checks may be done through comparison with other national inventory calculation data.

In the development of the Croatian inventory, certain steps and some of these checks were performed:

- Comparison with the national inventory data of other countries was conducted by comparing CRF tables or through a direct communication;
- Activity data were compared using different sources such as Croatian Bureau of Statistics and individual emission sources.
- The CO<sub>2</sub> emissions from fossil fuel combustion, within the framework of IPCC methodology, are estimated using two approaches: (1) Reference Approach and (2) Sectoral Approach (Tier 1).

National inventory report 2006 was technically reviewed by the UNFCCC expert review team in January 2007 (centralized review). The main findings of the ERT are that the inventory is well documented and the NIR and CRF are in conformity with the *Guidelines for the preparation of national communications by Parties included in Annex I to the Convention, Part I: UNFCCC reporting guidelines on annual inventories* and the *IPCC good practice guidance*. The ERT recognized the improvements made since the previous submission. Beside areas for further improvement identified by the Party and ERT it could be concluded that the NIR and the CRF have improved compared to the 2005 submission.

## 1.7. GENERAL UNCERTAINTY EVALUATION

The uncertainties associated with both annual estimates of emissions, and emission trends over time are reported according to the *Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories*. The Croatian inventory team estimates uncertainties using Tier 1 method described by the IPCC, which provides estimates of uncertainties by pollutant. The uncertainties are estimated for both excluding LULUCF and including LULUCF due to the *Good Practice Guidance for Land Use, Land-Use Change and Forestry*.

According to the uncertainty analysis the total uncertainty excluding LULUCF is 13.7 percent, while the total uncertainty including LULUCF is 15.5 percent. The uncertainty introduced into the trend in total national emissions excluding LULUCF is estimated to be 3.3 percent and including LULUCF 9.7 percent. The combined uncertainty as a share of total emissions is dominated by CH<sub>4</sub> emissions from fugitive emissions from oil and gas operations (uncertainty of the 13.1 percent excluding LULUCF and 10.5 percent including LULUCF). Furthermore, LULUCF sources/sink shows quite large uncertainty of 11 percent.

The results of the Tier 1 approach are shown in Table A5-1 and A5-2 (Annex 5), where the shaded rows represent key categories. In 2006 submission the error occurred in formula for combined uncertainty for limestone and dolomite use which was corrected according to the recommendation of the ERT. Therefore, the total uncertainty and uncertainty introduced into the trend including LULUCF are considerably smaller than in previous submission.

## 1.8. GENERAL ASSESSMENT OF THE COMPLETENESS

Croatian inventory consists of the emission estimates for the period from 1990-2005.

The completeness is evaluated following the IPCC methodology and appropriate use of the following notation keys: *NO* (not occurred); *NE* (not estimated); *NA* (not applicable); *IE* (included elsewhere); *C* (confidential). Detailed description by activities and gases of the status of the emission calculation is given in corresponding CRF tables.

Generally, the objective of the completeness is achieved in compliance with the capabilities of the Republic of Croatia in collecting adequate and acceptable activity data. The problems related with lack of activity data are described in sectoral chapters where necessary. The aim of the Croatian inventory is to include all antropogenic sources of GHGs in the future.

The summary of the “not estimated” sources/sinks is given in Annex 4 – Assessment of completeness and (potential) sources and sinks of greenhouse gas emissions and removals excluded, Table A4-1.

## **2. TRENDS IN GREENHOUSE GAS EMISSIONS**

### **2.1. BRIEF DESCRIPTION AND INTERPRETATION OF EMISSION TRENDS FOR AGGREGATED GREENHOUSE GAS EMISSIONS**

The total GHG emissions in 2005, excluding removals by sinks, amounted to 30.5 mill. t CO<sub>2</sub>-eq (equivalent CO<sub>2</sub> emissions), which represents 3.4 percent emissions reduction compared to GHG emission in the year 1990.

Overall decline of economic activities and energy consumption in the period 1991-1994, which was mainly the consequence of the war in Croatia, had directly caused the decline in total emissions of greenhouse gases in that period. With the entire national economy in transition process, some energy intensive industries reduced their activities or phased out certain productions (e.g. blast furnaces in Sisak, primary aluminium production in Šibenik, coke plant in Bakar), which was considerably reflected in GHG emissions reduction. Emissions have started to increase in the period 1995-1999 at an average rate of 3.6 percent per year, because of economy revitalisation. Last five years average increase was even 2.6 percent per year. The main reason of GHG emission increase was Energy sector, because of emission growth in sub-sectors Energy Industries and Transport.

## 2.2. BRIEF DESCRIPTION AND INTERPRETATION OF EMISSION TRENDS BY GAS

The shares of GHG emission have not significantly changed during the entire period. The CO<sub>2</sub> is the largest anthropogenic contributor to total national GHG emissions. In 2005 the shares of GHG emissions were as follows: 77.3 percent CO<sub>2</sub>, 9.7 percent CH<sub>4</sub>, 11.8 percent N<sub>2</sub>O and 0.1 percent HFCs. The trend of aggregated emissions/removals, divided by gasses, is shown in the Table 2.2-1 and the Figure 2.2-1.

Table 2.2-1: Aggregated emissions and removals of GHG by gases (1990-2005)

Gas	Emissions and removals of GHG (Gg CO <sub>2</sub> -eq)							
	1990	1995	2000	2001	2002	2003	2004	2005
CO <sub>2</sub>	23307	16877	19947	20945	22032	23501	23148	23567
CH <sub>4</sub> as CO <sub>2</sub> -eq	3247	2540	2548	2691	2746	2925	3015	2956
N <sub>2</sub> O as CO <sub>2</sub> -eq	4017	3116	3317	3289	3358	3263	3680	3608
HFCs as CO <sub>2</sub> -eq	43	43	23	49	49	164	189	349
PFCs as CO <sub>2</sub> -eq	937	0	0	0	0	0	0	0
SF <sub>6</sub> as CO <sub>2</sub> -eq	NE	NE	NE	NE	NE	NE	NE	NE
<b>Total GHG emission</b>	<b>31552</b>	<b>22576</b>	<b>25835</b>	<b>26974</b>	<b>28185</b>	<b>29853</b>	<b>30031</b>	<b>30481</b>
Removals (CO <sub>2</sub> )	<b>-6281</b>	<b>-10629</b>	<b>-8012</b>	<b>-8880</b>	<b>-8537</b>	<b>-7878</b>	<b>-7948</b>	<b>-7779</b>
<b>Total emission (including LULUCF)</b>	<b>25271</b>	<b>11947</b>	<b>17823</b>	<b>18094</b>	<b>19648</b>	<b>21975</b>	<b>22083</b>	<b>22702</b>




Figure 2.2-1: Trend of GHG emissions, by gases

### 2.2.1. CARBON DIOXIDE – CO<sub>2</sub>

The most significant anthropogenic greenhouse gas is carbon dioxide (CO<sub>2</sub>). In 2005, CO<sub>2</sub> emission were 1 percent larger than in 1990. CO<sub>2</sub> removals by sinks were almost 24 percent larger than removals in 1990. The largest CO<sub>2</sub> emission growth was in Energy sector (Road Transport and Public Electricity & Heat Production) and Industrial processes. There was a permanent increase in mobility (number of road vehicles) and therefore increase in motor fuel consumption in last ten years. There was also a significant increase in electricity demand and

supply. Consequently, two new thermal power plants were installed in last few years (coal burned thermal power plant - 210 MW and combined cycled gas turbine – 200 MW). The largest CO<sub>2</sub> emission growth in Industrial Processes is in Mineral production.

#### **2.2.2. METHANE – CH<sub>4</sub>**

The CH<sub>4</sub> emission in 2005 was 9 percent below the emission in 1990, largely due to decrease in emission in Agriculture sector (Enteric Fermentation and Manure Management), as a consequence of lower number of domestic animals.

#### **2.2.3. NITROUS OXIDE – N<sub>2</sub>O**

The N<sub>2</sub>O emission in 2005 was 10 percent lower than emission in 1990. Decrease of emission was in Energy Sector (Manufacturing Industries and Construction and Other Sectors), Industrial Processes (Nitric Acid Production) and Agriculture (Direct emission from agriculture soils, Direct N<sub>2</sub>O emissions from animals, Indirect N<sub>2</sub>O emission from nitrogen used in agriculture).

#### **2.2.4. FLUOROCARBONS – HFCS AND PFCs**

PFCs emissions were generated in the production of primary aluminium. The Croatian aluminium industry was still operational in 1990/1991, but production was stopped in 1992. HFCs were used as substitutes for cooling gases in refrigerating and air-conditioning systems that deplete the ozone layer. According to provided calculations, the contribution of F-gases in total national GHG emission in 2005 was 1 percent.

#### **2.2.5. SULPHUR HEXAFLUORIDE SF<sub>6</sub>**

The SF<sub>6</sub> emission estimation is still not included in the inventory, because the input data is not reliable.

## 2.3. BRIEF DESCRIPTION AND INTERPRETATION OF EMISSION TRENDS BY CATEGORY

According to the UNFCCC reporting guidelines and IPCC methodological guidelines, total national emission are divided into six sectors: Energy, Industrial Processes, Solvent and Other Product Use, Agriculture, Land Use, Land-Use Change and Forestry and Waste. The total national GHG emissions and removals, divided by sectors, are presented in the Table 2.3-1 and the Figure 2.3-1.

Table 2.3-1: Aggregated emissions and removals of GHG by sectors (1990-2005)

Source	Emissions and removals of GHG (Gg CO <sub>2</sub> -eq)							
	1990	1995	2000	2001	2002	2003	2004	2005
Energy	22504	16871	19287	20329	21531	22986	22442	22806
Industrial Processes	4206	2144	2874	2830	2746	2886	3252	3442
Solvent and Oth.Prod.Use	80	80	69	75	99	108	135	155
Agriculture	4464	3101	3130	3236	3276	3320	3561	3495
Waste	298	380	475	504	533	555	642	583
<b>Total GHG emission</b>	<b>31552</b>	<b>22576</b>	<b>25835</b>	<b>26974</b>	<b>28185</b>	<b>29853</b>	<b>30031</b>	<b>30481</b>
Removals (LULUCF)	<b>-6281</b>	<b>-10629</b>	<b>-8012</b>	<b>-8880</b>	<b>-8537</b>	<b>-7878</b>	<b>-7948</b>	<b>-7779</b>
<b>Total emission (including LULUCF)</b>	<b>25271</b>	<b>11947</b>	<b>17823</b>	<b>18094</b>	<b>19648</b>	<b>21976</b>	<b>22083</b>	<b>22702</b>




Figure 2.3-1: Trend of GHG emissions, by sectors

### 2.3.1. ENERGY

The most important IPCC sector in Croatia is Energy. The Energy sector accounted for some 75 percent of the total national GHG emissions (presented as equivalent emission of CO<sub>2</sub>). In 2005 the GHG emission from Energy was 1 percent larger than emission in 1990.

### **2.3.2. INDUSTRIAL PROCESSES**

Industrial Processes contributes to total GHG emission with approximately 10 percent, depending on the year. There was a significant decrease of GHG emission from industrial processes. The GHG emission in 2005 was 18 percent lower than emission in 1990.

### **2.3.3. SOLVENT AND OTHER PRODUCT USE**

Solvent and Other Product Use contributes to total GHG emission with some 0.25 - 0.5 percent of the total national GHG emissions (presented as equivalent emission of CO<sub>2</sub>). The GHG emission in 2005 was still 93 percent larger than emission in 1990.

### **2.3.4. AGRICULTURE**

The GHG emissions from Agriculture have also a decreasing trend. The GHG emission in 2005 was 22 percent lower in comparison with 1990 emission. According to estimation of Croatian experts for agriculture, approximately 12 percent of total GHG emissions belong to Agriculture.

### **2.3.5. WASTE**

Emissions from Waste sector have been constantly increasing in the period 1990-2005. Increasing emissions are a consequence of greater quantities of waste. The GHG emission in 2005 was 95 percent larger in comparison with 1990 emission. Contribution of waste sector to total GHG emission is approximately 2 percent.

## 2.4. BRIEF DESCRIPTION AND INTERPRETATION OF EMISSION TRENDS FOR INDIRECT GREENHOUSE GASSES AND SO<sub>2</sub>

Although they are not considered as greenhouse gases, photochemical active gases such as carbon monoxide (CO), oxides of nitrogen (NO<sub>x</sub>) and non-methane volatile organic compounds (NMVOCs) indirectly contribute to the greenhouse effect. These are generally referred to as indirect greenhouse gases or ozone precursors, because they take effect in the creation and degradation of O<sub>3</sub> as one of the GHGs. Sulphur dioxide (SO<sub>2</sub>), as a precursor of sulphate and aerosols, is believed to contribute negatively to the greenhouse effect. The emissions of ozone precursors and SO<sub>2</sub> are shown in the Table 2.4-1.

Table 2.4-1: Emissions of ozone precursors and SO<sub>2</sub> by different sectors (Gg)

Gas	Emissions (Gg)							
	1990	1995	2000	2001	2002	2003	2004	2005
<b>NO<sub>x</sub> Emission</b>	<b>83.9</b>	<b>57.7</b>	<b>68.4</b>	<b>69.4</b>	<b>68.1</b>	<b>68.7</b>	<b>67.4</b>	<b>73.3</b>
Energy Industries	13.6	10.3	12.0	11.6	13.2	13.8	11.2	12.0
Manufacturing Ind. & Construction	17.5	8.9	9.7	10.6	10.2	9.7	11.8	16.3
Transport	37.0	29.8	33.3	33.0	31.0	31.1	30.6	31.3
Other Energy (fuel combustion)	15.0	8.1	12.8	13.6	13.1	13.6	13.2	13.1
Fugitive Emission from Fuels	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.3
Industrial Processes	0.4	0.3	0.3	0.3	0.3	0.2	0.3	0.3
<b>CO Emission</b>	<b>502.1</b>	<b>339.4</b>	<b>378</b>	<b>322.9</b>	<b>305.8</b>	<b>321.5</b>	<b>308.5</b>	<b>329.6</b>
Energy Industries	1.5	1.0	1.2	1.0	1.0	1.4	1.2	0.9
Manufacturing Ind. & Construction.	40.4	41.3	37.8	38.6	37.9	38.0	41.6	32.7
Transport	252.5	176.6	188.7	165.0	151.0	137.5	125.4	159.2
Other Energy (fuel combustion)	204.0	116.7	146.5	115.2	113.0	141.4	136.4	133.0
Fugitive Emission from Fuels	0.6	0.5	0.5	0.4	0.4	0.4	0.5	0.5
Industrial Processes	3.1	3.3	3.3	2.7	2.5	2.8	3.4	3.3
<b>NMVOC Emission</b>	<b>144.8</b>	<b>141.8</b>	<b>218</b>	<b>177.7</b>	<b>290.9</b>	<b>416.3</b>	<b>481</b>	<b>437.6</b>
Energy Industries	0.3	0.2	0.3	0.3	0.3	0.3	0.3	0.3
Manufacturing Ind. & Construction	1.7	1.4	1.4	1.5	1.4	1.7	1.9	3.3
Transport	40.9	30.0	32.9	27.4	25.6	22.1	19.4	18.5
Other Energy (fuel combustion)	12.1	6.9	9.0	7.5	7.3	8.8	8.5	8.3
Fugitive Emission from Fuels	8.2	7.8	9.7	10.4	10.8	10.5	9.7	9.1
Industrial Processes	81.6	95.5	164.7	130.6	245.5	372.9	441.2	398.1
Solvent Use	27.4	27.4	23.4	25.5	33.8	36.7	46.1	52.9
<b>SO<sub>2</sub> Emission</b>	<b>175.2</b>	<b>81.5</b>	<b>69.9</b>	<b>69.9</b>	<b>75.6</b>	<b>74.2</b>	<b>61.6</b>	<b>68</b>
Energy Industries	78.5	39.0	25.4	26.0	23.3	35.7	25.7	32.8
Manufacturing Ind. & Construction	55.8	24.7	22.6	24.9	29.9	15.6	11.8	10.3
Transport	5.3	3.5	6.0	4.9	6.3	7.4	7.9	8.5
Other Energy (fuel combustion)	23.9	4.7	6.5	6.1	8.0	7.5	6.9	6.6
Fugitive Emission from Fuels	6.4	5.0	4.8	4.5	4.5	4.5	4.7	4.6
Industrial Processes	5.3	4.6	4.6	3.5	3.6	3.5	4.6	5.2



### 3. ENERGY (CRF sector 1)

#### 3.1. OVERVIEW OF SECTOR

##### 3.1.1. INTRODUCTION

This sector covers all activities that involve fuel combustion from stationary and mobile sources, and fugitive emission from fuels.

The Energy sector is the main cause for anthropogenic emission of greenhouse gases. It accounts for approximately 75 percent of the total emission of all greenhouse gases presented as equivalent emission of CO<sub>2</sub>. Looking at its contribution to total emission of carbon dioxide (CO<sub>2</sub>), the energy sector accounts for about 90 percent. The contribution of energy in methane (CH<sub>4</sub>) emission is substantially smaller (49 percent) while the contribution of nitrous oxide (N<sub>2</sub>O) is quite small (about 6 percent).

During full combustion, the carbon contained in fuel oxidizes and transforms into CO<sub>2</sub>, while through the incomplete combustion the small amounts of CH<sub>4</sub>, CO and NMVOC emissions also appear. The CO<sub>2</sub> is the most important greenhouse gas from fuel combustion. This was the reason for making a detailed estimate by IPCC methodology. The emission of CO<sub>2</sub> depends on the quantity and type of the fuel used. The specific emission is the largest during combustion of coal, then oil and natural gas. A rough ratio of specific emission during combustion of the stated fossil fuels is 1 : 0.75 : 0.55 (coal : oil : gas).

There are some other gases generated from fuel combustion such as methane (CH<sub>4</sub>) and nitrous oxide (N<sub>2</sub>O), and indirect greenhouse gases such as nitrogen oxides (NO<sub>x</sub>), carbon monoxide (CO) and non-methane volatile organic compounds (NMVOC). The indirect greenhouse gases participate in the process of ozone creating and destroying, which is one of the GHGs. In the framework of the IPCC methodology, the calculation of sulphur dioxide (SO<sub>2</sub>) emission is also recommended. The sulphur dioxide, as a precursor of sulphate and aerosols, is believed to have a negative impact on the greenhouse effect because the creation of aerosols removes heat from the environment.

The fuel fugitive emission is also estimated, which is generated during production, transport, processing, storing, and distribution of fossil fuels. These activities produce mainly the emission of CH<sub>4</sub>, and smaller quantities of NMVOC, CO and NO<sub>x</sub>.

Emissions from fossil fuel combustion comprise the majority (more than 90 percent) of energy-related emissions. Contribution of individual subsectors to emission of greenhouse gases, for the last estimated year (2005), is presented in the Figure 3.1-1.

Greenhouse gases are also generated during combustion of biomass and biomass-based fuels. The CO<sub>2</sub> emission from biomass, in line with IPCC recommendations, is not included into the national emission totals because emitted CO<sub>2</sub> had been previously absorbed from the atmosphere for growth and development of biomass. Removal or emission of CO<sub>2</sub> due to the changes in the forest biomass is estimated in the sector of Land Use, Land-use Change and Forestry.

The emission from fuel combustion in international air and waterborne transport is reported separately and it has not been included in the national emission totals.




Figure 3.1-1: The contribution of different subsectors to GHG emission, year 2005

### 3.1.2. ENERGY STRUCTURE

The basis for an estimate of the GHGs emission from Energy sector is the national energy balance, production, imports, exports, stock change and consumption of fuels are shown in the national energy balance report in natural units (kg or m<sup>3</sup>) or energy units (J). National energy balance for 2005 is presented in Annex 3.

For easier data comparison in energy balance the natural units are transformed to energy units using proper national net calorific values for different fuels. The structure of energy consumption of fossil fuels from 1990 to 2005 is shown in Figure 3.1-2.




Figure 3.1-2: Structure of energy consumption

Liquid fossil fuels are mainly used with share between 60 to 70 percent, and natural gas with approximately 30 percent, while share of solid fossil fuels is 3-11 percent. Fuel woods and biomass-based fuels are neutral with regard to CO<sub>2</sub> emission. Therefore, they are not shown in the Figure 3.1-2.

### 3.1.3. COMPARISON OF THE SECTORAL APPROACH WITH THE REFERENCE APPROACH

The methodology used for estimating CO<sub>2</sub> emissions follows the *Revised 1996 IPCC Guidelines*. The emission of CO<sub>2</sub> is calculated using 2 different approaches: Reference approach and Sectoral approach. Sectoral emission estimates are based on fuel consumption data given in National Energy Balance, where energy demand and supply is given at sufficiently detailed level, what allows emissions estimation by sectors and subsectors. In Reference approach the input data are production, import, export, international bunkers and stock change for primary and secondary fuel. Comparison between these approaches was made and presented in Annex 3. The total differences in fuel consumption and CO<sub>2</sub> emissions for chosen years are given in Table 3.1-1.

Table 3.1-1: The fuel consumption and CO<sub>2</sub> emissions from fuel combustion (Reference & Sectoral approach)

	1990	1995	2000	2001	2002	2003	2004	2005
Fuel consumption (PJ)								
Reference appr.	316.9	232.1	268.0	278.4	296.1	316.0	310.1	311.8
Sectoral appr.	287.2	216.2	251.0	261.8	277.0	294.7	287.6	290.4
Apparent energy consumption (PJ)	289.0	217.7	250.6	263.3	281.3	297.2	287.6	291.3
<b>Relative Differen. (%)</b>	0.6	0.7	-0.2	0.6	1.6	0.8	0.0	0.3
CO <sub>2</sub> Emission (Gg)								
Reference appr.	20842	15140	17739	18543	19805	21171	20557	20921
Sectoral appr.	20529	14843	17255	18126	19273	20651	20048	20416
<b>Relative Differen. (%)</b>	1.5	2.0	2.8	2.3	2.8	2.5	2.5	2.5

The CO<sub>2</sub> emission calculated by Reference approach is higher compare to Sectoral approach. The reason is CO<sub>2</sub> emission from non-energy fuel consumption which is calculated under Reference approach while under Sectoral approach is not.

### 3.1.4. INTERNATIONAL BUNKER FUELS

The CO<sub>2</sub> emissions from the consumption of fossil fuels for aviation and marine international transport activities, as required by the IPCC methodology, are reported separately, and not included in national emission totals. The fuel consumption (PJ) for International Aviation and Marine Bunkers and GHG emissions for observed period are shown in the Table 3.1-2.

International marine bunkers are included in national energy balance for the period from 1994 to 2005, as separate data. Until the year 1994, international marine bunkers are based on expert estimation. According to suggestion of review team the disaggregation of fuel between international and domestic aviation was recalculated based on International Energy Agency (IEA) data. International aviation bunker was included in national energy balance data first time for the year 2004.

*Table 3.1-2: Fuel consumption and GHG emissions for International aviation and marine bunkers, from 1990 to 2005*

	1990	1991	1992	1993	1994	1995	1996	1997
<b>Fuel combustion (PJ)</b>								
Aviation bunkers	2.86	0.00	0.00	1.41	2.81	2.51	2.37	2.51
Marine bunkers	1.44	0.95	1.07	1.52	1.83	1.36	1.52	0.97
<b>Total bunkers</b>	<b>4.30</b>	<b>0.95</b>	<b>1.07</b>	<b>2.92</b>	<b>4.64</b>	<b>3.86</b>	<b>3.90</b>	<b>3.48</b>
<b>CO<sub>2</sub>-eq emission (Gg)</b>								
Aviation bunkers	202.26	0.00	0.00	99.57	199.15	177.37	168.03	177.37
Marine bunkers	108.54	71.34	80.62	114.54	138.33	102.01	114.91	73.63
<b>Total bunkers</b>	<b>310.80</b>	<b>71.34</b>	<b>80.62</b>	<b>214.11</b>	<b>337.48</b>	<b>279.38</b>	<b>282.94</b>	<b>251.00</b>

*Table 3.1-2: Fuel consumption and GHG emissions for International aviation and marine bunkers, from 1990 to 2005 (cont.)*

	1998	1999	2000	2001	2002	2003	2004	2005
<b>Fuel combustion (PJ)</b>								
Aviation bunkers	2.68	1.58	1.41	0.88	0.84	1.01	1.23	1.70
Marine bunkers	1.08	0.88	0.76	1.19	0.98	0.91	0.97	1.05
<b>Total bunkers</b>	<b>3.76</b>	<b>2.46</b>	<b>2.16</b>	<b>2.07</b>	<b>1.81</b>	<b>1.92</b>	<b>2.20</b>	<b>2.74</b>
<b>CO<sub>2</sub>-eq emission (Gg)</b>								
Aviation bunkers	189.81	112.02	99.57	62.23	59.12	71.57	87.13	120.11
Marine bunkers	81.00	65.68	57.02	89.37	73.24	68.67	73.06	78.98
<b>Total bunkers</b>	<b>270.81</b>	<b>177.70</b>	<b>156.59</b>	<b>151.60</b>	<b>132.36</b>	<b>140.24</b>	<b>160.19</b>	<b>199.09</b>

### 3.1.5. FEEDSTOCKS AND NON-ENERGY USE OF FUELS

Non-energy fuel consumptions (fuels used as feedstock) and appropriate emissions, where one part or even the whole carbon is stored in product for a longer time and the other part oxidizes and goes to atmosphere, are described here. The feedstock use of energy carriers occurs in

chemical industry (natural gas consumption for ammonia production, production of naphtha, ethane, paraffin, and wax), construction industry (bitumen production), and other products such as motor oil, industrial oil, grease etc. As a result of non-energy use of bitumen in construction industry there is no CO<sub>2</sub> emission because all carbon is bound to the product. In order to avoid double counting, CO<sub>2</sub> emission in non-energy consumption of natural gas in ammonia production, as well as non-energy consumption of naphtha, lubricants, ethane and other, was estimated in sector Industrial Processes.

### **3.1.6. CO<sub>2</sub> CAPTURE FROM FLUE GASES AND SUBSEQUENT CO<sub>2</sub> STORAGE**

There are no plants for recovery and storage of CO<sub>2</sub> in Croatia. Natural gas produced in Croatian gas fields contains a large amount of CO<sub>2</sub>, more than 15 percent, and before coming to commercial pipeline has to be cleaned (scrubbed), but CO<sub>2</sub> was emitted without capture and storage. The CO<sub>2</sub> emission from gas scrubbing in Central Gas Station Molve, estimated by material balance method, is described in the Chapter 3.3.1.2.

### **3.1.7. COUNTRY-SPECIFIC ISSUES**

There are also a few technical country-specific issues, which are connected to GHG emission calculation in Energy sector:

- The methodology for estimating CO<sub>2</sub> emission from natural gas scrubbing is not given in IPCC Guidelines. The CO<sub>2</sub> emission is determined on the base of differences in CO<sub>2</sub> content before and after scrubbing units and quantity of scrubbed natural gas (material balance method).
- Country-specific net calorific values obtained from national energy balance are used in GHG emission calculation (Annex 2).

## 3.2. FUEL COMBUSTION ACTIVITIES (CRF 1.A.)

### 3.2.1. SOURCE CATEGORY DESCRIPTION

#### 3.2.1.1. Energy Industries (CRF 1.A.1.)

This subsector comprises emission from fuel combustion in public electricity and heat production plants, petroleum refining plants, solid transformation plants, oil and gas extraction and coal mining. The total GHG emission from Energy Industries is given in the Table 3.2-1 and Figure 3.2-1.

Table 3.2-1: The CO<sub>2</sub>-eq emissions (Gg) from Energy Industries

CO <sub>2</sub> -eq emission (Gg)	1990	1995	2000	2001	2002	2003	2004	2005
Public Electric. and Heat Product.	3724	2973	3817	4475	5167	5756	4598	4660
Petroleum Refining	2575	1892	1792	1603	1799	1881	1916	1811
Other Energy Industries	545	327	291	234	269	264	279	351
<b>Total Energy Industries</b>	<b>6844</b>	<b>5192</b>	<b>5901</b>	<b>6312</b>	<b>7235</b>	<b>7901</b>	<b>6793</b>	<b>6821</b>




Figure 3.2-1: The CO<sub>2</sub>-eq emissions from energy industries

It should be stressed out that a large part of the electrical energy is generated without GHG emission; therefore the emission from this sector is relatively small, 28-35 percent of emission from total Energy sector. The largest part (53-73 percent) of the emission is a consequence of fuel combustion in thermal power plants, then the combustion in oil refineries 24-40 percent. The remaining combustion in oil and gas fields, coal mines and the coke plant accounts for some 3-9 percent.

#### **Public Electricity and Heat Production (CRF 1.A.1.a)**

The installed electricity generating capacities in the Republic of Croatia include power plants owned by the HEP Group (Croatian Power Company), a certain number of industrial power plants and a few privately owned power plants (wind power plants, small hydro power plants).

Total capacities serving the needs of the Croatian electric power system amount to 4049 MW (with 50% of nuclear unit Krško in Slovenia). Out of this amount, 1632 MW is placed in thermal power plant, 2079 MW in hydro power plant and 338 MW in the nuclear unit Krško (50% of total

available capacity). Generating capacities of HPPs, TPPs and NPP Krško are presented in the Table 3.2-2.

*Table 3.2-2: Generating capacities of HPPs, TPPs and NPP Krško*

	Available Power (MW)		Fuel
	Generator	Net Output	
HPPs		2079	water
NPP Krško*	353.5	338	UO2
TPP Plomin 1	105	98	coal
TPP Plomin 2**	210	192	coal
TPP Rijeka	320	303	fuel oil
TPP Sisak	2x210	396	fuel oil / n. gas
CHP Zagreb (east)	25 + 120 + 210	337	f.oil / n.gas / ELO
CHP Zagreb (west)	12.5 + 32 + 52	92	f.oil / n.gas / ELO
CPP Osijek	45 + 2x25	89	f.oil / n.gas / ELO
CCGT Jertovec	2x42.5	83	n.gas / ELO
Emergency diesel (4)	29	29	D2
Emergency diesel (1)	13	13	2GT
<b>Total (HPPs+NPP+TPPs)</b>		<b>4049</b>	

UO2 - uranium oxide

ELO - extra light oil

D2/2GT - special fuel oil for operation of emergency TPPs

\* - 50% of NPP Krško is owned by HEP

\*\* - TPP Plomin 2 Ltd. (HEP and RWE Power Co-ownership – share 50% : 50%)

During the observed period between 1990 and 2005 in Croatia only 18 to 38 percent of Croatian electricity demands were covered by thermal power plants. The largest contribution to electricity production in Croatia had hydro power plants 48 to 74 percent. Nuclear power plant Krško delivered 50 percent of it's electricity to Croatian power system until 1998 (the delivery was stopped). The delivery of electricity from NPP Krško started again in 2003. The past few years the electricity demand was compensated with import. Therefore, in 2000 the electricity import was larger than production in all Croatian thermal power plants (TPPs). In 2005, the import of electricity was about 37 percent of total electricity consumption in Croatia.

In this subsector, there are few types of plants:

- Thermal Power Plants (TPPs), which produce only electricity
- Public Cogeneration Plants (PCPs), which produce combined heat and electricity
- Public Heating Plants (PHPs), which produce only heat

Electricity and heat production, fuel consumption and GHG emissions for the years 1990, 1995, 2000, 2001, 2002, 2003, 2004 and 2005 are presented in tables A2-1 to A2-2 of the Annex 2.

### **Petroleum Refining (CRF 1.A.1.B)**

Croatia has two oil refineries in Rijeka and Sisak, while lubricants are produced in Rijeka and Zagreb. Processing capacities of the Croatian refineries, which belong to INA – oil and gas company, are shown in the Table 3.2-3.

*Table 3.2-3: Processing Capacities of Oil & Lube Refineries*

<b>PROCESSING CAPACITIES</b>	<b>INSTALLED (1000 t/year)</b>
<b>Oil Refinery Rijeka (Urinj)</b>	
atmospheric distillation	5000
reforming	730
fluidized-bed catalytic cracking (FCC)	1000
visbreaking	600
isomerisation	250
hydrodesulphurisation (HDS)	1040
mild hydrocracking (MHC)	560
<b>Lube Refinery Rijeka (Mlaka)</b>	
vacuum distillation	630
deasphalting	110
furfural extraction	220
deparaffination	140
ferofining	230
deoiling	30
bitumen	350
<b>Oil Refinery Sisak</b>	
atmospheric distillation	4000
reforming	720
fluidized-bed catalytic cracking (FCC)	500
coking	240
vacuum distillation	800
bitumen	350
<b>Lube Refinery Zagreb</b>	
atmospheric distillation	-
lubricants	60

In the refineries, there are two types of fuel combustion – for heating and/or cogeneration and for own use of energy for production processes. National energy balance gives separate numbers for own use of energy in refineries, while fuel combustion for heating/cogeneration plants is presented together with similar industrial plants. Because of that, cogeneration and heating plants in refineries were calculated (previous submission) in the sub-sector Manufacturing Industries and Construction instead of Energy Industries (Petroleum Refining).

Fuel consumption and GHG emissions from petroleum refining are presented in tables A2-5 and A2-6 of the Annex 2.



### **Manufacturing of Solid Fuels and Other Energy Industries (CRF 1.A.1.C)**

In Croatia the coal production was rather low. In 1999, last coal mines in Istria were closed. Coke-oven plant in Bakar, nearby Rijeka, was also closed in 1994.

Crude oil is produced from 35 oil fields and gas condensation products from 10 gas-condensations fields, which covers about 35 percent of the total domestic demand.

Natural gas is produced from 20 gas fields, which covers about 60 percent of the total demand. The largest quantities come from the Molve, Kalinovac and Stari Gradec, where the Central Gas Stations (CGS Molve) for gas processing and transport preparation were built – Molve I, II and III. Their capacities are:

- 1 mill. m<sup>3</sup>/day for Molve I
- 3 mill. m<sup>3</sup>/day for Molve II
- 5 mill. m<sup>3</sup>/day for Molve III

The underground gas storage Okoli was designed with the nominal capacity of 550 mill. m<sup>3</sup>. Maximal injection capacity is 3.8 mill. m<sup>3</sup>/day and maximal drawdown capacity is 5 mill. m<sup>3</sup>/day.

Fuel consumption and GHG emissions from manufacturing of solid fuels and other energy industries are presented in table A2-7.

Fuel consumption and GHG emissions from Manufacturing Industries and Construction are presented in tables A2-8 and A2-9.

#### **3.2.1.2. Manufacturing Industries and Construction (CRF 1.A.2.)**

Manufacturing Industries and Construction include the emissions from fuel combustion in different industries, such as iron and steel industries, industries of non-ferrous metals, chemicals, pulp and paper, food processing, beverages and tobacco, construction and building material industries. This sector also includes the emissions from fuel used for the generation of electricity and heat in industry (industrial cogeneration plants and industrial heating plants). The total GHG emission from Manufacturing Industries and Construction is given in the Table 3.2-4 and Figure 3.2-2.

*Table 3.2-4: The CO<sub>2</sub>-eq emissions (Gg) from Manufacturing Industries and Construction*

	1990	1995	2000	2001	2002	2003	2004	2005
Iron & Steel Industry				93	70	81	65	89
Non-Ferrous Metals				16	20	16	27	21
Chemical				540	420	481	655	473
Pulp, Paper & Print				123	109	114	125	130
Food Processing, Bev. & Tab.				451	515	469	488	542
Other (Constr. Material...)				2.437	2.376	2.463	2.700	2.897
<b>Total Manufact. Ind. &amp; Constr.</b>	<b>6.045</b>	<b>3.505</b>	<b>3.516</b>	<b>3.661</b>	<b>3.510</b>	<b>3.624</b>	<b>4.060</b>	<b>4.151</b>




Figure 3.2-2: The CO<sub>2</sub>-eq emissions from Manufacturing Industries and Construction

The emission from this sector contributes 14-25 percent of the total emission from Energy sector. In national energy balance the fuel combustion in industrial cogeneration and heating plants is not divided on appropriate industrial branches, for which electricity and/or thermal energy is produced. The fuel consumed in industrial cogeneration and heating plants is divided by industrial subsectors only for the years 2001, 2002, 2003, 2004 and 2005. The largest contributions to emissions have the fuel combustion in industry of contraction materials (subsector: Other in Figure 3.2-2), then follows chemical industry, food processing industry, iron and steel industry, industry of glass and non-metal, non-ferrous metal and paper industry.

The GHG emissions from Manufacturing Industries and Construction by fuels is shown in Table A2-9 and Table A2-10 of the Annex 2.

### 3.2.1.3. Transport (CRF 1.A.3.)

The emission from combustion and evaporation of fuel for all transport activities is included in this sector. In addition to road transport, this sector includes the emission from air, rail and marine transport as well. The total GHG emission from Transport sector is given in the Table 3.2-5 and Figure 3.2-3.

Table 3.2-5: The CO<sub>2</sub>-eq emissions (Gg) from Transport

	1990	1995	2000	2001	2002	2003	2004	2005
Road Transport	3583	3179	4353	4415	4723	5131	5285	5485
Domestic Aviation	298	88	126	163	157	142	160	174
Railways	138	107	86	88	87	88	92	96
National Navigation	134	99	86	92	111	112	91	100
<b>Total Transport</b>	<b>4153</b>	<b>3472</b>	<b>4651</b>	<b>4758</b>	<b>5078</b>	<b>5473</b>	<b>5630</b>	<b>5855</b>




Figure 3.2-3: The CO<sub>2</sub>-eq emissions from transport

The emissions from fuel consumption in aircraft or marine vessel engaged in international transport are excluded from the national total. These emissions are reported separately.

The contribution from Transport to total emissions from Energy sector was 18-25 percent. The most of the emission comes from road transport (86-94 percent), than from domestic air, rail and marine transport (Figure 3.2-3). The increase of emissions from this sector is a consequence of growth of mobility and number of road motor vehicles.

### Road Transport

The COPERT III package (Tier 2/3 method) was used for CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O emissions (and other pollutants) calculation from road transport in the period from 1990 to 2005. For calculating emissions, COPERT III emission factors per fuel types were used.

The aggregate number of road motor vehicles is presented in the Table A2-10 of the Annex 2. Fuel consumption and GHG emissions from Road Transport are presented in tables A2-11 of the Annex 2.

Key assumption – motor fuel tanked (filled in vehicle reservoir) abroad and consumed in Croatia is equal with fuel tanked in Croatia and consumed abroad. Fuel consumption calculated by COPERT multiplying number of vehicles and annual average vehicle mileage should be equal with appropriate data from national energy balance (difference is less than 1%). It is necessary to avoid inconsistency in trend emissions.

### Off-road Transport

The GHG emission calculation from off-road transport was calculated using Tier 1 approach, based on fuel consumption data (national energy balance) and default IPCC emission factors. The fuel consumption and appropriate GHG emissions for domestic air transport, national navigation and railway transport are shown in the tables A2-12, A2-13 and A2-14 of the Annex 2. According to suggestion of review team the disaggregation of fuel between international and domestic aviation was recalculated based on International Energy Agency (IEA) data.

### 3.2.1.4. Small Stationary Energy Sources (CRF 1.A.4.)

This sector includes emission from fuel combustion in commercial and institutional buildings, emission from fuel combustion in residential sector and the emission from fuel combustion in agriculture, forestry and fishing. The total GHG emissions from abovementioned small stationary energy sources are shown in the Table 3.2-6 and Figure 3.2-4.

Table 3.2-6: The CO<sub>2</sub>-eq emissions (Gg) from small stationary energy sources

	1990	1995	2000	2001	2002	2003	2004	2005
Commercial/Institutional	790	612	606	711	751	777	773	752
Residential	2178	1689	2009	2156	2257	2469	2443	2489
Agriculture/Forestry/Fishing	843	583	861	801	741	755	702	712
<b>Total</b>	<b>3810</b>	<b>2883</b>	<b>3476</b>	<b>3669</b>	<b>3750</b>	<b>4001</b>	<b>3918</b>	<b>3953</b>




Figure 3.2-4: The CO<sub>2</sub>-eq emissions from small stationary sources

The CO<sub>2</sub>-eq emissions from these subsectors were about 17-20 percent of the total emissions from energy sector. The most of the emission comes from small household furnaces and boiler rooms (55-62 percent), then from service sector (15-21 percent), while the combustion of fuel in agriculture, forestry and fishing accounts for 18 to 25 percent (Figure 2.3-4).

The GHG emission calculation from these subsectors was calculated using Tier 1 approach, based on fuel consumption data (national energy balance) and default IPCC emission factors. The fuel consumption and GHG emissions for Commercial/Institutional, Residential and Agriculture/Forestry/Fishing are presented in the tables A2-15, A2-16 and A2-17, Annex 2.

### 3.2.1.5. Ozone Precursors and SO<sub>2</sub> Emissions

The emission of indirect greenhouse gases (NO<sub>x</sub>, CO and NMVOC) and SO<sub>2</sub> is described in this chapter. Ozone precursors are cause of greenhouse gas - tropospheric ozone, whereas SO<sub>2</sub> was added to a list of pollutants first time in Revised 1996 IPCC Guidelines for National GHG Inventories due to the importance of this gas from the position of acidification and eutrophication.

The emission of NO<sub>x</sub> is the largest from road transport (about 50 percent), then from energy industries and manufacturing industries and construction. Emissions of CO and NMVOC are mainly from road transport and small household furnaces using firewood or coal. The emission of SO<sub>2</sub> mainly originates from stationary energy sources, such as thermal power plants and refineries, and depends on the quantity of fuel used and the sulphur content of fuel.

The emissions of ozone precursors and SO<sub>2</sub> are shown in the Table 3.2-7.

*Table 3.2-7: Emissions of ozone precursors and SO<sub>2</sub> from fuel combustion (Gg)*

<b>Emission (Gg)</b>	<b>1990</b>	<b>1995</b>	<b>2000</b>	<b>2001</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>
<b>NO<sub>x</sub> Emission</b>	<b>83.1</b>	<b>57.2</b>	<b>67.9</b>	<b>68.8</b>	<b>67.6</b>	<b>68.2</b>	<b>66.8</b>	<b>72.9</b>
Energy Industries	13.6	10.3	12.0	11.6	13.2	13.8	11.2	12.0
Manuf. Ind. & Construction	17.5	8.9	9.7	10.6	10.2	9.7	11.8	16.3
Transport	37.0	29.8	33.3	33.0	31.0	31.1	30.6	31.3
Other Energy	15.0	8.1	12.8	13.6	13.1	13.6	13.2	13.1
<b>CO Emission</b>	<b>498.4</b>	<b>335.6</b>	<b>374.2</b>	<b>319.8</b>	<b>302.9</b>	<b>318.3</b>	<b>304.7</b>	<b>325.8</b>
Energy Industries	1.5	1.0	1.2	1.0	1.0	1.4	1.2	0.9
Manuf. Ind. & Construction	40.4	41.3	37.8	38.6	37.9	38.0	41.6	32.7
Transport	252.5	176.6	188.7	165.0	151.0	137.5	125.4	159.2
Other Energy	204.0	116.7	146.5	115.2	113.0	141.4	136.4	133.0
<b>NMVOC Emission</b>	<b>55.1</b>	<b>38.5</b>	<b>43.6</b>	<b>36.6</b>	<b>34.6</b>	<b>32.9</b>	<b>30.2</b>	<b>30.3</b>
Energy Industries	0.3	0.2	0.3	0.3	0.3	0.3	0.3	0.3
Manuf. Ind. & Construction	1.7	1.4	1.4	1.5	1.4	1.7	1.9	3.3
Transport	40.9	30.0	32.9	27.4	25.6	22.1	19.4	18.5
Other Energy	12.1	6.9	9.0	7.5	7.3	8.8	8.5	8.3
<b>SO<sub>2</sub> Emission</b>	<b>163.6</b>	<b>71.8</b>	<b>60.5</b>	<b>61.9</b>	<b>67.5</b>	<b>66.2</b>	<b>52.3</b>	<b>58.2</b>
Energy Industries	78.5	39.0	25.4	26.0	23.3	35.7	25.7	32.8
Manuf. Ind. & Construction	55.8	24.7	22.6	24.9	29.9	15.6	11.8	10.3
Transport	5.3	3.5	6.0	4.9	6.3	7.4	7.9	8.5
Other Energy	23.9	4.7	6.5	6.1	8.0	7.5	6.9	6.6

### 3.2.2. METHODOLOGICAL ISSUES

The GHG emission calculation is mainly provided using Tier 1 approach. There are two exceptions, as follows:

- Thermal power plants and public cogeneration plants (Energy Industries, CRF 1.A.1.a)
- Road transport (Transport, CRF 1.A.3.b)

#### 3.2.2.1. Tier 1 Approach

##### CO<sub>2</sub> emissions

The majority of energy-related GHG emissions belong to CO<sub>2</sub>. That is the reason why it is analysed in greater detail by IPCC methodology given in the *Revised 1996 IPCC Guidelines for National GHG Inventories*.

The CO<sub>2</sub> emission is estimated by two approaches: (1) Reference approach and (2) Sectoral approach. Inputs in the Reference approach are production, import, export, international bunkers, and stock change for primary and secondary fuel. The Sectoral approach is used to

identify the emission by means of fuel consumption for each group of sources (sectors). The energy data from the national energy balance are recalculated from natural units into energy units by means of own net calorific values for each fuel. Calorific values are also taken from the energy balance. The emission factors used for calculation are taken from *IPCC Guidelines (Revised 1996 IPCC Guidelines for National GHG Inventories, Workbook, Page 1.6)*.

Since the combustion processes are not 100 percent efficient, the part of carbon stored is not emitted to the atmosphere so it occurs as soot, ash and other by-products of inefficient combustion. Therefore, it is necessary to know the fraction of carbon which oxidizes. This value was taken from *IPCC Guidelines* as recommended (Workbook, Page 1.8).

Non-energy uses of fossil fuels can result in storage (in products) of some or all of the carbon contained in the fuel for a certain period of time, depending on the end-use. The fraction of carbon stored in products is suggested in *IPCC Guidelines* (Workbook, auxiliary worksheet 1-1, page 1.37).

According to the *IPCC guidelines* the emission from international transport activities was not included in national totals.

### **Emissions of CH<sub>4</sub>, N<sub>2</sub>O and indirect greenhouse gases**

Emissions of CH<sub>4</sub>, N<sub>2</sub>O and indirect greenhouse gases (NO<sub>x</sub>, CO and NMVOC) have been identified by Tier 1 method in such a way that the fuel used in each sector is multiplied by the emission factor suggested in *Revised 1996 IPCC Guidelines for National GHG Inventories* (Reference Manual, page 1.33-1.42). The basis for the estimate is the fuel used in different energy sectors. The used fuel is grouped into basic fossil fuels categories according to its aggregate condition: coal, natural gas and oil, and biomass-based fuel. Data about quantities of the fuel used are taken from the national energy balance.

In order to identify the SO<sub>2</sub> emission, besides the data on the type and the quantity of fuel consumed it is necessary to know the sulphur content in fuel. The available data on the sulphur content were collected from INA - Oil and Gas Company (for petroleum derivatives: gasoline, residual oil, diesel oil, jet fuel) and from HEP – Croatian Electric Utility Company (for fossil fuels consumed in thermal power plants).

#### **3.2.2.2. Tier 2/3 Approach**

##### **Thermal power plants and public cogeneration plants (CRF 1.A.1.a)**

The GHG emissions from thermal power plants and public cogeneration plants, for last four years (2001, 2002, 2003, 2004 and 2005), were calculated using more detailed Tier 2 approach. Tier 2 approach is based on bottom-up fuel consumption data from every boiler or gas turbine in plant. There were available data about monthly fuel consumption and detailed fuel characteristics data (net calorific value, sulphur and ash content...). For estimation of CO<sub>2</sub> emissions, default IPCC emission factors were used, while implied emission factors for CH<sub>4</sub> and N<sub>2</sub>O are based on technology type and configuration (Tier 2).

### **Road transport (CRF 1.A.3.b)**

The COPERT III package (Tier 2/3 method) was used for air emission calculation from road transport emission in the period from 1990 to 2005 for CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O emissions.

Very detailed set of input data is necessary for COPERT implementation. In Croatian case, main data provider is Croatian Centre for Vehicles, which is responsible for compilation of detailed motor vehicle database. The database assures the following information about:

- type of vehicles (passenger cars, light duty vehicles, heavy duty vehicles, buses, mopeds, motorcycles)
- type of motor (gasoline four-stroke, gasoline two-stroke, diesel, rotation motor and electromotor)
- cylinder capacity (<1.4 lit, 1.4-2.0 lit, >2.0 lit)
- weight class (<3.5 t, 3.5-7.5 t, 7.5-16 t, 16-32 t, >32 t)
- age of vehicles (distribution of vehicles per ECE categories according to EC directives)
- registration area (country distribution)
- annual average vehicle mileage

Fuel consumption data (from Energy Institute "Hrvoje Požar") and fuel characteristics data (from INA - Oil and Gas Company) are also necessary for calculation of emissions from road transport using COPERT software.

Additional data, like highway, rural and urban transport mileage, average speed of different kind of vehicles and different road types, average daily trip distance, beta value (the fraction of the monthly mileage driven before the engine and any exhaust components have reached their nominal operation temperature) and temperature per month are estimated (based on data from statistics) or COPERT default data are used.

COPERT calculates emission factors according to driving conditions data (the average speed per vehicle type and per road), fuel variables and climate conditions (average monthly temperatures data).

### **3.2.3. UNCERTAINTIES AND TIME-SERIES CONSISTENCY**

#### **3.2.3.1. Uncertainty of CO<sub>2</sub> emissions**

The CO<sub>2</sub> emission, from the fossil fuel combustion, depends of the amount of fuel consumed (from energy balance), net calorific values (from energy balance), carbon emission factors (IPCC recommendation), the fraction of carbon stored (IPCC recommendation) and the fraction of carbon oxidised (IPCC recommendation).

The national energy balance is based on data from all available sources. The data from Central Bureau of Statistics about production, usage of raw material and consumption of fuels in all industrial facilities in Croatia are used. The data from questionnaires about monthly use of natural gas in certain sectors from all distributive companies in Croatia, about annual consumption of coal in certain sectors and the data from Customs Administration about export

and import of fossil fuels are also used. The data from these sources and other necessary data are organised in related database. The estimated uncertainty of data from energy balance is below 5 percent.

The accuracy of data on net calorific values, which are also taken from national energy balance, is high.

There are more uncertainties in data on international marine and aviation bunkers. Nevertheless, possible errors in estimated values do not significantly affect on the accuracy of data of national emission, as marine and aviation transport have relatively small influence. The estimated CO<sub>2</sub> emissions for International Marine and Aviation Transport are not included in national totals.

The other data needed for calculation, such as, carbon emission factors, the fraction of carbon stored for non-energy uses of fuel and the fraction of carbon oxidized, are taken from *Revised 1996 IPCC Guidelines for National GHG Inventories*. Experts believe that CO<sub>2</sub> emission factors for fuels are generally well determined within 5 percent, as they are primarily dependent on the carbon content of the fuel.

For example, for the same primary fuel type (e.g. coal), the amount of carbon contained in the fuel per unit of useful energy can vary. Non-energy uses of the fuel can also create situations where the carbon is not emitted to the atmosphere (e.g. plastics, asphalt, etc.) or is emitted at a much-delayed rate. Additionally, inefficiencies in the combustion process, which can result in ash or soot remaining unoxidized for long periods, were also assumed. These factors all contribute to the uncertainty in the CO<sub>2</sub> estimates. However, these uncertainties are believed to be relatively small. Overall uncertainty for CO<sub>2</sub> emission estimates from the fossil fuel combustion are considered accurate within 7 percent.

### **3.2.3.2. Uncertainty of CH<sub>4</sub>, N<sub>2</sub>O and indirect greenhouse gases emissions**

Estimates of CH<sub>4</sub>, N<sub>2</sub>O and ozone precursor emissions are based on fuel (coal, natural gas, oil and bio-fuels) and aggregate emission factors for different sectors. Uncertainties in estimates are due to the fact that emissions are estimated on the base of emission factors representing only a limited subset of combustion conditions.

Using the aggregate emission factors for each sector, the differences between various types of coal and especially liquid fuel are not included, nor are the differences in the technology and the contribution of equipment for emission reduction. Therefore, the uncertainties associated with emission estimates of these gases are greater than estimates of CO<sub>2</sub> emissions from the fossil fuel combustion.

The uncertainty of CH<sub>4</sub> emission is estimated to  $\pm 40$  percent, while the uncertainty of N<sub>2</sub>O emission is estimated to factor 2 (the emission could be twice larger or smaller than the estimated one). The largest part of uncertainty refers to the emission factor applied while the fuel consumption data (national energy balance) are rather good. Implementation of Tier 2/3 approach for estimation of CH<sub>4</sub> and N<sub>2</sub>O emissions from thermal power plants and public



cogeneration plants (CRF 1.A.1.a) and road transport (CRF 1.A.3.b) lead to certain uncertainty reduction (Annex 4).

### 3.2.3.3. Time-series consistency

Activity data, emission factors and methodology implied for GHG emission calculation from fuel combustion activities is very consistent for entire period. Negligible inconsistency is a consequence of implementation of more detailed approach (Tier 2/3) for last four years in Energy Industries.

### 3.2.4. SOURCE-SPECIFIC QA/QC

Quality control activities were divided in two phases, first phase included activities during the inventory preparation performed by sector expert, and the second phase included audit conducted by the expert designated for QA/QC after the preparation of final draft of the NIR.

During the preparation of the inventory submission activities related to quality control were mainly focused on completeness and consistency of emission estimates and on proper use of notation keys in the CRF tables. Also, several checks have been carried out in order to ensure correct aggregation from lower to higher reporting level and correct use of conversion factors.

The basis for emission estimates in Energy sector is Energy balance prepared by Energy Institute "Hrvoje Požar" and mainly default emission factors provided by the IPCC. Background information and assumptions for entire time-series are transparently recorded in *Inventory Data Record Sheets* which allow third party to evaluate quality of estimates in this sector.

After preparation of final draft of this chapter an audit was carried out to check selected activities from Tier 1 General inventory level QC procedures and Tier 2 source-specific QC procedures. The audit revealed that most of the Tier 1 QC activities were correctly carried out during preparation of the inventory despite the fact that formal QC procedures were not prepared.

Regarding to Tier 2 activities, emission factors and activity data were checked for key source categories. In Energy industries, Public Electricity and Heat Production a more detailed Tier 2 methodology was applied for the period 2001-2005, due to availability of detail information on fuel consumption in the facilities. These data are still not available for other sub-categories therefore Tier 1 methodology was applied. Also, inventory team used country-specific fuel net calorific values for emission estimates. In Mobile combustion – Road, a COPERT III model was used for whole period (1990-2005). This model requires a very detailed set of input data and could be considered as a Tier 3 methodology. In Mobile combustion – Domestic and International Aviation, a data from International Energy Agency statistics was used in order to reduce trend inconsistency, but it was point out that uncertainty of international bunkers is relatively higher comparing to other data.

### 3.2.5. SOURCE-SPECIFIC RECALCULATIONS

All recalculations were made in response to the review process, according to suggestions given by expert review team.

#### Road Transportation (1.A.3.b.)

According to suggestion of review team, Tier 3 method for CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O emissions calculation is used, because of trend consistency reasons. Consequently, CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O emissions from Road transport (1990-2005) were recalculated, as follows:

Years:	1990-2005
Gases:	CO <sub>2</sub> , CH <sub>4</sub> and N <sub>2</sub> O
Method:	Tier 3 methodology for calculation of CO <sub>2</sub> , CH <sub>4</sub> and N <sub>2</sub> O emission from Road transport is implied, because of trend consistency (in previous submission COPERT III was used for calculation of all GHG emissions, for period 2001-2004).
EF:	Emission factors based on COPERT III software for CO <sub>2</sub> , CH <sub>4</sub> and N <sub>2</sub> O is implied

#### Manufacturing industry and construction (1.A.2)

Energy consumption of natural gas in petrochemical industry is calculated under 1.A.2.f Manufacturing industries and construction – Other – Petrochemical Industry.

Years:	1990-2005
Gases:	CO <sub>2</sub> , CH <sub>4</sub> and N <sub>2</sub> O
Method:	Tier 1 methodology for calculation of CO <sub>2</sub> , CH <sub>4</sub> and N <sub>2</sub> O emissions from Manufacturing industries and construction – Other – Petrochemical Industry was used for calculation of all GHG emissions (in previous submission energy use of fuels in Petrochemical industry wasn't reported)
EF:	Default emission factors for CO <sub>2</sub> , CH <sub>4</sub> and N <sub>2</sub> O is implied

#### Other Stationary - Non energy fuel consumption (1.A.5)

According to suggestion of review team, reconciliation between the reference and sectoral approaches was made. Natural gas and liquid fuels from non energy fuel consumption is reported in 1.A.d. instead of 1.A.5.a.

Years:	1990-2005
Gases:	CO <sub>2</sub> , CH <sub>4</sub> and N <sub>2</sub> O
Method:	Tier 1 methodology for calculation of CO <sub>2</sub> , CH <sub>4</sub> and N <sub>2</sub> O emissions from Non energy fuel consumption was used (in previous submission non energy fuel consumption was reported in the 1.A.5.a instead of 1.A.d)
EF:	Default emission factors for CO <sub>2</sub> , CH <sub>4</sub> and N <sub>2</sub> O is implied

### **Indirect GHG (NO<sub>x</sub>, CO, NMVOC, SO<sub>2</sub>)**

Emissions of indirect GHGs for whole time period (1990-2005) was set up according to the SNAP 97 nomenclature of EMEP/CORINAIR methodology. Emissions were taken from the emission inventory report which is Croatia's obligation in the framework of the Long Range Transboundary Air Pollution (LRTAP) Convention according to the Act on Air Protection (Official Gazette No. 178/04).

#### **3.2.6. SOURCE-SPECIFIC PLANNED IMPROVEMENTS**

For the purpose of GHG inventory improvement, missing data should be collected and also quality of existing data, emission factors and methods should be improved. Implementation of well-documented country specific emission factors and appropriate detailed methods are recommended. Consequently, the main objectives of the GHG inventory improvement plan are:

- data gaps reduction,
- data collection improvement,
- activity data and emission factors uncertainties reduction,
- activities on improvement methodologies and emission factors, documentation and description of inventory system.

As a result of comprehensive analysis of GHG inventory quality, based on information prepared in the framework of Centralized Review Report, short-term and long-term goals for GHG inventory improvement are obtained.

##### Short-term goals (< 1 years)

Generally, the changes from Tier 1 to Tier 2/3 estimation methodologies for Energy key sources, as much as possible, are recommended. The priority should be the key sources with high uncertainties of emission estimation. But, significant constraints are availability of activity data, especially for the beginning years of concerned period. Consequently, implementation of more detailed methodology approach (Tier 2/3) for key sources, for entire period, will be very difficult.

The extensive use of detailed methodology (Tier 2/3) for Energy Industries is also one of the short-term goals. For achievement of abovementioned goal is necessary to ensure delivery of detailed activity data for Energy Industries. The good example is the usage of technology/plant-specific data for sub-sector Thermal power plants and public cogeneration plants (Tier 2) for the last five inventory years.

##### Long-term goals (> 1 years)

The extensive use of plant-specific data collected in the framework of Cadastre of Emissions to Environment (CEE) is recommended ("bottom up" approach). In addition, usage of more source-specific QA/QC procedures will improve the quality of GHG inventory in Energy sector.

### 3.3. FUGITIVE EMISSIONS FROM FUELS (CRF 1.B.)

#### 3.3.1. SOURCE CATEGORY DESCRIPTION

This section describes fugitive emission of greenhouse gases from coal, oil and natural gas activities. This category includes all emissions from mining, production, processing, transportation, and use of fossil fuels. During all stages from the extraction of fossil fuels to their final use, the escape or release of gaseous fuels or volatile components may occur.

##### 3.3.1.1. Solid fuels (CRF 1.B.1.)

All underground and opencast coal mines release methane during their regular operation. The amount of methane generated during mining is primarily a function of the coal rank and mining depth, as well as other factors such as moisture. After coal has been mined, small amounts of methane retained in coal are released during post-mining activities, such as coal processing, transportation and utilization.

In Croatia the coal production was rather low. Until 1999 only underground coal mines in Istria were in operation (Tupljak, Ripenda and Koromačno) and they produced some 0.015 to 0.174 mill. tons of coal. Global Average Method (Tier 1) was used for the methane emission estimation and the estimated emission was 0.2 to 2.3 Gg. The emissions of methane from mining and post-mining activities are showed in the Figure 3.3-1 and Table A2-18, Annex 2.




Figure 3.3-1: The fugitive emissions of methane from coal mines

##### 3.3.1.2. Oil and natural gas (CRF 1.B.2.)

The fugitive emission of methane is inevitable during all the activities involving oil and natural gas. This category includes the fugitive emission from production, refining, transportation, processing, and distribution of crude oil or oil products and gas. The fugitive emission also includes the emission of methane, which is the result of incomplete combustion of gas during flaring, and the emission from venting during oil and gas production.


The most significant fugitive emissions after methane among the activities relating to oil and gas are the emissions of non-methane volatile organic compounds (NMVOCs). They are produced

by evaporation when fuel oil gets in contact with air during refining, transportation, and distribution of oil products. In addition to NMVOCs there are fugitive emissions of NO<sub>x</sub>, CO and SO<sub>2</sub> during various processes in oil refineries.

### **Fugitive emission of methane**

For estimating the fugitive emission of methane the simplest procedure has been used (Tier 1), which is based on production, unloading, processing, and consumption of oil and gas.

According to IPCC, all countries are divided into regions with relatively homogenous characteristics of oil and gas systems. Croatia is included in the region that covers the countries of Central & East Europe and former Soviet Union. For this region higher emission factors are provided, especially for the gas system. In the absence of better data, average emission factors provided for the region are used for estimating the fugitive emission of methane. Estimated results are given in Figure 3.3-2 and Table A2-19, Annex 2.



*Figure 3.3-2: The fugitive emissions of methane from oil and gas activities*

The fugitive emission of methane is mainly (about 97 percent) consequence of production, transmission, and distribution of natural gas. The fugitive emission from oil accounts for about 1 percent and venting and flaring of gas/oil production accounts for approximately 2 percent.

### **Fugitive emission of ozone precursors and SO<sub>2</sub>**

A simplified Tier 1 procedure was used to make a fugitive emission estimate of ozone precursors and SO<sub>2</sub> from oil refineries for the entire period from 1990 to 2005. The simplified procedure is based on the quantity of crude oil processed in oil refineries. Default emission factors were used for the estimation. A summary of estimated results of the fugitive emissions of CO, NO<sub>x</sub> and NMVOC and SO<sub>2</sub> are illustrated in the table 3.3-1.

Table 3.3-1: The fugitive emissions of ozone precursors and SO<sub>2</sub> from oil refining

Emissions (Gg)	1990	1995	2000	2001	2002	2003	2004	2005
CO emission	0.62	0.48	0.46	0.43	0.43	0.44	0.46	0.45
NO <sub>x</sub> emission	0.41	0.32	0.31	0.29	0.29	0.29	0.30	0.30
NM VOC emission	8.23	7.77	9.73	10.41	10.81	10.51	9.70	9.05
SO <sub>2</sub> emission	6.38	4.96	4.80	4.49	4.49	4.52	4.72	4.60

### **CO<sub>2</sub> emission from natural gas scrubbing**

Fugitive emission of greenhouse gases from coal, oil and natural gas, due to mining, production, processing, transportation and use of fossil fuels is also part of Energy sector. Although these emission sources are not characteristic in respect of CO<sub>2</sub> emission, specifically in Croatia emission of CO<sub>2</sub> from natural gas scrubbing in Central Gas Station Molve is assigned here. IPCC doesn't offer methodology for estimating CO<sub>2</sub> emission scrubbed from natural gas and subsequently emitted into atmosphere.

Natural gas produced in Croatian gas fields (Molve, Kalinovac and Stari Gradac) contains a large amount of CO<sub>2</sub>, more than 15 percent, and before coming to commercial pipeline has to be cleaned (scrubbed). Since the maximum volume content of CO<sub>2</sub> in commercial natural gas is 3 percent, it is necessary to clean the natural gas before transporting through pipeline to end-users. Because of that, the Scrubbing Units exist at largest Croatian gas field. The estimated CO<sub>2</sub> emissions, by the material balance method, are presented in Table 3.3-2.

Table 3.3-2: The CO<sub>2</sub> emissions (Gg) from natural gas scrubbing in CGS Molve

CO <sub>2</sub> emission (Gg)	1990	1995	2000	2001	2002	2003	2004	2005
Central Gas Station MOLVE	416	697	633	688	665	684	710	691

### **3.3.2. METHODOLOGICAL ISSUES**

The fugitive emission of methane from coal, oil, and gas has been identified by Tier 1 method with average emission factors given in *Revised 1996 IPCC Guidelines for National GHG Inventories* (Workbook, page 1.26 and 1.30). Data about quantities of the mined coal and production, unloading, transportation, processing, storing and consumption of oil and gas are taken from the national balance energy supply and demand.

Inputs on processed crude oil in refineries are taken from national energy balance while emission factors are taken from *IPCC Guidelines* (Reference Manual, page 1.133 and 1.134).

The methodology for estimating CO<sub>2</sub> emission from natural gas scrubbing is not given in IPCC Guidelines. The CO<sub>2</sub> emission is determined on the base of differences in CO<sub>2</sub> content before and after scrubbing units and quantity of scrubbed natural gas.

### 3.3.3. UNCERTAINTIES AND TIME SERIES CONSISTENCY

#### 3.3.3.1. Uncertainty

The fugitive emission of methane from coal mining and handling is determined by use of Global Average Method (Tier 1), which is based on multiplication of coal produced and emission factor. The amount of coal produced is taken from energy balance and that value is very accurate. The main uncertainty of calculation depends on accuracy of used emission factor. The arithmetic average value of emission factor has been chosen from IPCC for the region to which Croatia belongs. The estimated uncertainty of methane emissions, for underground mining may be as high as a factor of 2 and for post-mining activities a factor of 3.

The Production-Based Average Emission Factors Approach is used to determine fugitive emission from oil and natural gas activities. This approach is based on activity data (production, transport, refining and storage of fossil fuels) and average emission factors. Due to the complexity of the oil and gas industry, it is difficult to quantify the uncertainties. The uncertainty of calculation is linked mostly to the emission factor, just like the determination of fugitive emission of methane from coal mining and handling. The expert estimated that accuracy of calculation of fugitive emission from oil is better than from fugitive emission from gas, but the uncertainty of both estimations is pretty high. Similarly, the uncertainty of calculation of emission of ozone precursors and SO<sub>2</sub> is also very high.

The CO<sub>2</sub> emission from scrubbing of natural gas is also shown here. The calculation is based on material balance which gives much better accuracy ( $\pm 10$  percent).

#### 3.3.3.2. Time-series consistency

Activity data, emission factors and methodology implied for fugitive emission from fuels is consistent for entire period.

#### 3.3.4. SOURCE-SPECIFIC QA/QC

Quality control activities were divided in two phases, first phase included activities during the inventory preparation performed by sector expert, and the second phase included audit conducted by the expert designated for QA/QC after the preparation of final draft of the NIR.

During the preparation of the inventory submission activities related to quality control were mainly focused on completeness and consistency of emission estimates and on proper use of notation keys in the CRF tables. Also, several checks have been carried out in order to ensure correct aggregation from lower to higher reporting level and correct use of conversion factors.

The basis for emission estimates in Energy sector is Energy balance prepared by Energy Institute "Hrvoje Požar" and mainly default emission factors provided by the IPCC. Background information and assumptions for entire time-series are transparently recorded in *Inventory Data Record Sheets* which allow third party to evaluate quality of estimates in this sector.

After preparation of final draft of this chapter an audit was carried out to check selected activities from Tier 1 General inventory level QC procedures and Tier 2 source-specific QC

procedures. The audit revealed that most of the Tier 1 QC activities were correctly carried out during preparation of the inventory despite the fact that formal QC procedures were not prepared.

For Fugitive emissions from oil and gas operations a Tier 1 method was applied and emission factor is a mean value of the range proposed in the IPCC Manual. The CO<sub>2</sub> emission from natural gas scrubbing in CPS Molve was estimated using country specific methodology since IPCC Guidelines does not provide methodology for this source category.

### 3.3.5. SOURCE-SPECIFIC RECALCULATIONS

#### Production and processing 1.B.2.b (ii)

According to suggestion of review team, CO<sub>2</sub> emission from scrubbing is reported in subcategory 1.B.2.b(ii) Production and processing instead of 1.B.2.d Other, for whole time series, although activity data doesn't exist (country specific issue).

Years: 1990-2005

Gas: CO<sub>2</sub>

Method: The CO<sub>2</sub> emission is determined on the base of differences in CO<sub>2</sub> content before and after scrubbing units and quantity of scrubbed natural gas

AD: Doesn't exist

EF: Doesn't exist

### 3.3.6. SOURCE-SPECIFIC PLANNED IMPROVEMENTS

For estimation of fugitive emissions from oil and natural gas operations, a Tier 1 method was applied. Used emission factors are an average value of the range proposed in the IPCC Manual. However, fugitive emission from natural gas is key source and implementation of rigorous source-specific evaluations approach (Tier 3) is necessary. The Tier 3 approach will generally involve compiling the following types of information:

- detailed inventories of the amount and types of process infrastructure (e.g. wells, field installations and production/processing facilities),
- production disposition analyses oil and gas production, vented, flared and reinjected volumes of gas, and fuel gas consumption,
- accidental releases (i.e. well blow-outs and pipeline ruptures),
- typical design and operating practices and their impact on the overall level of emission control.

For implementation of rigorous source-specific evaluations approach (Tier 3) is necessary additional technical and financial resources.



### 3.4. OVERVIEW OF GHG EMISSIONS FROM ENERGY SECTOR

This chapter gives overview of the GHG emissions. The contribution of individual energy subsectors to the total emissions of greenhouse gases for the observed period is given in the Table 3.3-3 and Figure 3.3-3.

Table 3.3-3: The CO<sub>2</sub>-eq emissions (Gg) from Energy sector

	1990	1995	2000	2001	2002	2003	2004	2005
Energy Industries	6844	5192	5901	6312	7235	7901	6793	6821
Manufac. Ind. and Const.	6045	3505	3516	3661	3510	3624	4060	4151
Transport	4153	3472	4651	4758	5078	5473	5630	5855
Other Energy	3810	2883	3476	3669	3750	4001	3918	3953
Fugitive Emissions	1651	1818	1744	1929	1958	1987	2041	2025
<b>Total emission (Gg CO<sub>2</sub>-eq)</b>	<b>22504</b>	<b>16871</b>	<b>19287</b>	<b>20329</b>	<b>21531</b>	<b>22986</b>	<b>22442</b>	<b>22806</b>




Figure 3.3-3: The CO<sub>2</sub>-eq emissions from Energy sector

The Energy sector was the main cause for anthropogenic emission of greenhouse gases. It accounted for approximately 75 percent of the total emission of all greenhouse gases presented as equivalent emission of CO<sub>2</sub>. Looking at its contribution to total emission of carbon dioxide (CO<sub>2</sub>), the energy sector accounts for approximately 90 percent. The contribution of energy in methane (CH<sub>4</sub>) emission is substantially smaller (48 percent) while the contribution of nitrous oxide (N<sub>2</sub>O) is quite small (6 percent).

The largest part (28 to 35 percent) of the emissions are a consequence of fuel combustion in Energy Industries, then the combustion in Transport with increasing trend (18 percent in 1990; 26 percent in 2005) and the combustion in Manufacturing Industries and Construction with decreasing trend (25 percent in 1990; 18 percent in 2005). Small stationary energy sources, such as Commercial/Institutional, Residential and Agriculture/Forestry/Fishing, contribute to total emission from Energy sector with 17 to 20 percent, while fugitive emissions from fuels contribute with about 10 percent. The majority of energy-related GHG emissions belong to CO<sub>2</sub> (91 to 93 percent), then follows CH<sub>4</sub> (6 to 8 percent) and N<sub>2</sub>O (less than 1 percent).

### 3.5. REFERENCES

- Central Bureau of Statistics (2006) *Statistical Yearbook – for period from 1990 till 2005*, Zagreb
- Croatian Centre for Vehicles (2006) *Motor Vehicle Databases for 2001-2005*, data on CD
- FCCC/ARR/2005/HRV (2007) *Report of the individual review of the greenhouse gas inventory of Croatia submitted in 2006*, Centralized Review Report
- EEA (2000) *COPERT III Computer Programme to Calculate Emissions from Road Transport*, Denmark
- HEP (2005) *Annual Report of Air Emissions from HEP's Thermal Power Plants (draft)*, prepared by EKONERG, Zagreb
- INA (2006) *Data about natural gas scrubbing in Central Gas Station MOLVE and sulphur content in liquid fossil fuels*, INA – Oil and Gas Company, data sent by fax or by mail
- IPCC (2000) *Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories*, Japan
- IPCC/UNEP/OECD/IEA (1997) *Greenhouse Gas Inventory – Workbook*, Revised 1996 IPCC Guidelines for National Greenhouse Inventories, Volume 2, United Kingdom
- IPCC/UNEP/OECD/IEA (1997) *Greenhouse Gas Inventory – Reference Manual*, Revised 1996 IPCC Guidelines for National Greenhouse Inventories, Volume 3, United Kingdom
- Ministry of Economy, Labour and Entrepreneurship (2007) *Energy in Croatia 2005*, Annual Energy Report, Energy Institute "Hrvoje Požar", Zagreb
- Ministry of Environmental Protection, Physical Planning and Construction (2006) *National Inventory Report 2006, Croatian greenhouse gas inventory for the period 1990 – 2004*, EKONERG, Zagreb
- Ministry of Environmental Protection, Physical Planning and Construction (2006) *Second, Third and Fourth National Communication of the Republic of Croatia under the United Nations Framework Convention on Climate Change (UNFCCC)*, Zagreb
- Vuk B. *National Energy Balances – for period from 1990 till 2005*, Energy Institute Hrvoje Požar, Zagreb

## 4. INDUSTRIAL PROCESSES (CRF sector 2)

### 4.1. OVERVIEW OF SECTOR

Greenhouse gas emissions are produced as by-products of non-energy industrial processes in which raw materials are chemically transformed to final products. During these processes different greenhouse gases (GHGs) such as carbon dioxide (CO<sub>2</sub>), methane (CH<sub>4</sub>) or nitrous oxide (N<sub>2</sub>O) are released in the atmosphere.

Industrial processes whose contribution to CO<sub>2</sub> emissions was identified as significant are production of cement, lime, ammonia, ferroalloy, as well as use of limestone and soda ash in different industrial activities. Nitric acid production is source of N<sub>2</sub>O emissions. Emissions of CH<sub>4</sub> are appeared in production of other chemicals, as well as carbon black, ethylene and dichloroethylene.

Consumption of halocarbons (HFCs), which are used as substitution gases in refrigeration and air conditioning systems, is source of emissions of fluorinated compounds.

Some industrial process, particularly petrochemical, generate emissions of short-lived ozone and aerosol precursor gases such as carbon monoxide (CO), nitrogen oxides (NO<sub>x</sub>), non-methane volatile organic compounds (NMVOC) and sulphur dioxide (SO<sub>2</sub>). These gases indirect contribute to greenhouse effect.

The general methodology applied to estimate emissions associated with each industrial process, as recommended by Revised 1996 IPCC Guidelines and Good Practice Guidance and Uncertainty Management in National GHG Inventories involves the product of amount of material produced or consumed, and an associated emission factor per unit of production/consumption.

The activity data on production/consumption for particular industrial process are, in most cases, extracted from Monthly Industrial Reports, published by Central Bureau of Statistics, Department of Manufacturing and Mining. These reports cover industrial activities according to prescribed national classification of activities and comprise data on production and consumption of raw materials on monthly basis. In cases when such data were insufficient or some production-specific data were required to calculate emissions, individual manufacturers were contacted and voluntary surveys were carried out.

Regulation on the Greenhouse Gases Emissions Monitoring in the Republic of Croatia (Official Gazette No. 1/07), which will go into force in 2008, prescribes obligation and procedure for emissions monitoring, which comprise estimation and/or reporting of all anthropogenic emissions and removals. According to requirement, sources of abovementioned greenhouse gases should report required activity data for more accurate emissions estimation.

Emission factors used for calculation of emissions are, in most cases, default emission factors according to *Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories*, and *Good Practice Guidance and Uncertainty Management in National GHG Inventories*, mainly due

to a lack of plant-specific emission factors. Country-specific emission factors for cement production as well as ammonia production were estimated by collecting the actual data from individual plants.

Uncertainty estimates associated with emission factors for some industrial processes are well reported in *Good Practice Guidance*, while those associated with activity data are based on expert judgements since statistics and manufacturers have not particularly assessed the uncertainties.

Generally, CO<sub>2</sub> emissions from industrial processes declined from 1990 to 1995, due to the decline in industrial activities caused by the war in Croatia, while in the period 1996-2005 emissions slightly increased. Production of iron and aluminium were stopped in 1992.

The total annual emissions of GHGs, expressed in Gg CO<sub>2</sub>-eq, from Industrial Processes in the period 1990-2005 are presented in the Figure 4.1-1.





Figure 4.1-1: Emissions of GHGs from Industrial Processes (1990-2005)

## 4.2. MINERAL PRODUCTS (CRF 2.A.)

### 4.2.1. CEMENT PRODUCTION

#### 4.2.1.1. Source category description

During cement production, calcium carbonate (CaCO<sub>3</sub>) is heated in a cement kiln at high temperatures to form lime (i.e. calcium oxide, CaO) and CO<sub>2</sub> in a process known as calcination or calcining:



Lime is combined with silica-containing materials (clays or shales) to form dicalcium and tricalcium silicates which are the main constituents of cement clinker, with the earlier CO<sub>2</sub> being released in the atmosphere as a by-product. The clinker is then removed from the cement kiln, cooled, pulverized and mixed with small amount of gypsum to form final product called Portland cement.

There are four manufacturers of cement in Croatia, producing mostly Portland cement. There is production of Aluminate cement in the minor quantities. CO<sub>2</sub> emitted during the cement production process represents the most important source of non-energy industrial process of total CO<sub>2</sub> emissions. Different raw materials are used for Portland cement and Aluminate cement production. The quantity of the CO<sub>2</sub> emitted during Portland cement production is directly proportional to the lime content of the clinker. Emissions of SO<sub>2</sub> (non-combustion emissions) in the cement production originate from sulphur in the clay raw material.

#### 4.2.1.2. Methodological issues

Estimation of CO<sub>2</sub> emissions is accomplished by applying an emission factor, in tonnes of CO<sub>2</sub> released per tonne of clinker produced, to the annual clinker output corrected with the fraction of clinker that is lost from the kiln in the form of Cement Kiln Dust (CKD), (Tier 2 method, *Good Practice Guidance*).

Country-specific emission factor for Portland cement was estimated by using data on CaO and MgO content of clinker produced from individual plants. Corrections for imports of CaO and MgO via raw materials were made. CO<sub>2</sub> from Cement Kiln Dust (CKD) leaving the kiln system was calculated based on the volumes of dust and an emission factor. Because CKD is not fully calcined, the correction factor for CKD (CF<sub>ckd</sub>) was determined based on the emission factor for clinker and the calcinations rate of the CKD, which is based on plant-specific data.

The emission factor for Aluminate cement, which was obtained by cement manufacturer, equals 0.325 tonnes of CO<sub>2</sub> per tonne of clinker produced.

The activity data for clinker production, data on the CaO and MgO content of the clinker and information on the CKD collection and recycling practices and likewise on the calcination fraction of the CKD were collected by EKONERG from survey of cement manufacturers. The data were cross-checked with cement production data from Monthly Industrial Reports published by Central Bureau of Statistics, Department of Manufacturing and Mining. The data on clinker production and emission factors are presented in Table 4.2-1. The quantity of clinker imported has not been considered in the emission estimations.

Table 4.2.1: Clinker production and emission factors (1990 - 2005)

Year	Clinker production Portland cement (tonnes) <sup>1</sup>	Clinker production Aluminate cement (tonnes) <sup>1</sup>	Actual clinker production (tonnes) <sup>2</sup>	Emission factor Portland cement (t CO <sub>2</sub> /t clinker)	Emission factor Aluminate cement (t CO <sub>2</sub> /t clinker)
1990	1986783	0	2006651	0.522	-
1991	1264391	0	1277035	0.513	-
1992	1512065	0	1527186	0.518	-
1993	1256472	0	1269037	0.518	-
1994	1542560	0	1557986	0.521	-
1995	1133436	0	1144770	0.518	-
1996	1225925	0	1238184	0.504	-
1997	1442183	0	1456605	0.516	-
1998	1571767	0	1587485	0.515	-

Table 4.2.1: Clinker production and emission factors (1990 - 2005), cont.

Year	Clinker production Portland cement (tonnes) <sup>1</sup>	Clinker production Aluminate cement (tonnes) <sup>1</sup>	Actual clinker production (tonnes) <sup>2</sup>	Emission factor Portland cement (t CO <sub>2</sub> /t clinker)	Emission factor Aluminate cement (t CO <sub>2</sub> /t clinker)
1999	2063838	0	2084476	0.507	-
2000	2308148	73999	2406708	0.519	0.325
2001	2645180	94065	2767578	0.511	0.325
2002	2627934	70664	2726291	0.504	0.325
2003	2609349	82741	2719838	0.508	0.325
2004	2764331	87911	2881644	0.510	0.325
2005	2827258	99320	2956837	0.510	0.325

<sup>1</sup> Clinker production according to survey of cement manufacturers

<sup>2</sup> Actual clinker productions calculated as a product of clinker production and CF<sub>ckd</sub>, which is, for Portland cement; in the range 1.009 – 1.014; for Aluminate cement CF<sub>ckd</sub> equals 1.02

Import/export quantities of clinker are presented in Table 4.2-2.

Table 4.2.2: Import/export quantities of clinker (1990 - 2005)

Year	Clinker import (Portland cement)/ tonnes	Clinker export (Portland cement)/ tonnes
1990	0	0
1991	0	0
1992	0	4376
1993	0	0
1994	0	0
1995	52500	0
1996	0	32715
1997	57973	63529
1998	116397	82451
1999	0	114868
2000	0	111226
2001	0	131565
2002	0	5029
2003	102501	0
2004	51791	53387
2005	0	195888

The resulting emissions of CO<sub>2</sub> from Cement Production in the period 1990-2005 are presented in the Figure 4.2-1.




Figure 4.2-1: Emissions of CO<sub>2</sub> from Cement Production (1990-2005)

The activity data for cement production (see Table 4.2-3) were collected by EKONERG from survey of cement manufacturers and cross-checked with cement production data from Monthly Industrial Reports published by Central Bureau of Statistics, Department of Manufacturing and Mining.

Table 4.2-3: Cement production (1990-2005)

Year	Cement production (tonnes)
1990	2564359
1991	1667161
1992	1782124
1993	1589106
1994	2044234
1995	1556711
1996	1620633
1997	1873472
1998	2161827
1999	2549726
2000	2992854
2001	3237249
2002	3491746
2003	3689697
2004	3643549
2005	3629053

SO<sub>2</sub> emissions originate from sulphur in the fuel and in the clay raw material. The fuel emissions are counted as energy emissions (these emissions are presented in the chapter on emissions from energy sources). SO<sub>2</sub> emissions from the clay are counted as process emissions and calculated on the basis of produced quantities of cement. About 70-95 percent of the SO<sub>2</sub> generated in the process is absorbed in the produced alkaline clinker. SO<sub>2</sub> emissions have been calculated by applying emission factor of 0.3 kg SO<sub>2</sub>/tonne cement according to *Revised 1996 IPCC Guidelines*.

The resulting emissions of SO<sub>2</sub> from Cement Production in the period 1990-2005 are presented in the review on indirect GHG emissions from non-energy industrial processes.

#### **4.2.1.3. Uncertainties and time-series consistency**

Uncertainties contained in CO<sub>2</sub> emissions estimates are primarily related to uncertainties in the fraction of lime in domestic cement clinker and the actual fraction of CKD. According to *Revised 1996 IPCC Guidelines* most of the cement currently produced in the world is of Portland cement type<sup>1</sup>, which contains 60-67 percent lime by weight.

Uncertainty estimate associated with default emission factors amounts 3 percent, accordingly to values (1 to 5 percent) reported in *Good Practice Guidance*. Uncertainty estimate associated with activity data amounts 3 percent (1 to 5 percent), based on expert judgements since statistics and manufacturers have not been particularly assessed the uncertainties.

Emissions from Cement Production have been calculated using the same method and data sets for every year in the time series.

#### **4.2.1.4. Source-specific QA/QC and verification**

During the preparation of the inventory submission activities related to quality control were mainly focused on completeness and consistency of emission estimates and on proper use of notation keys in the CRF tables.

After preparation of final draft of this chapter an audit was carried out to check selected activities from Tier 1 General inventory level QC procedures and Tier 2 source-specific QC procedures. Cement Production is one of the key source categories in Industrial Processes. Regarding to Tier 2 activities, emission factors and activity data were checked for key source categories. CO<sub>2</sub> emissions from cement production were estimated using Tier 2 method which is a *good practice*.

#### **4.2.1.5. Source-specific recalculations**

In the previous report, two different data source were used across the reporting years. For 1990-2001 years data were obtained from document "Data for drafting baseline scenarios and analysis of GHG emission reductions in the Republic of Croatia" (J. Garilovic), which were cross-checked with clinker and cement production data from Monthly Industrial Reports published by Central Bureau of Statistics, Department of Manufacturing and Mining. For 2002-2004 years data obtained from survey of cement manufacturers, which were cross-checked with data from Monthly Industrial Reports.

In this report, clinker and cement production data for the period 1990-2005 were collected from survey of cement manufacturers due to improving comparability and time series consistency. Also, country specific EFs and its CKD correction factor were developed. Thereupon, CO<sub>2</sub> emissions are recalculated for the period 1990-2004.

---

<sup>1</sup> In the period 1990-2005 average 98 percent of cement produced in Croatia were of Portland cement type.




#### 4.2.1.6. Source-specific planned improvements

Since Cement Production is a key source category, towards increasing of CO<sub>2</sub> estimation accuracy, more detailed plant-level data on the carbonate raw materials are planned to investigate.

### 4.2.2. LIME PRODUCTION

#### 4.2.2.1. Source category description

The production of lime involves a series of steps which include quarrying the raw material, crushing and sizing, calcination and hydration. CO<sub>2</sub> is generated during the calcination stage, when limestone (CaCO<sub>3</sub>) or dolomite (CaCO<sub>3</sub>\*MgCO<sub>3</sub>) are burned at high temperature (900-1200 °C) in a rotary kiln to produce quicklime (CaO) or dolomitic lime (CaO\*MgO) and CO<sub>2</sub> which is released in the atmosphere:



#### 4.2.2.2. Methodological issues

Calculation of CO<sub>2</sub> emissions from lime production is accomplished by applying an emission factor in tonnes of CO<sub>2</sub> released per tonne of quicklime or dolomitic lime produced, to the annual lime output. The emission factors were derived on the basis of calcination reaction depending on the type of raw material used in the process and assuming 100 percent pure products.

According to aforementioned, emission factors for production of quicklime and dolomitic lime equals 0.79 tonnes CO<sub>2</sub>/tonnes quicklime produced and 0.91 tonnes CO<sub>2</sub>/tonnes dolomitic lime produced, respectively (*Revised 1996 IPCC Guidelines*).

The activity data for total lime production (see Table 4.2-4) were extracted from Monthly Industrial Reports published by Central Bureau of Statistics, Department of Manufacturing and Mining, and also were collected by EKONERG from survey of lime manufacturer since national classification of activities does not distinguish quicklime and dolomitic lime production. Also, certain amounts of quicklime were produced in the blast furnace processes, during 1990 and 1991.

Table 4.2-4: Lime production (1990-2005)

Year	Quicklime production (tonnes)	Dolomitic lime production (tonnes) <sup>1</sup>
1990	202253	0
1991	121710	0
1992	68976	0
1993	76269	0
1994	75511	0
1995	78820	0
1996	57522	37042
1997	65231	55047
1998	72419	53367
1999	68684	53088
2000	77804	68999
2001	102802	68427
2002	98325	94831
2003	92263	96820
2004	153056	58711
2005	157097	79470

<sup>1</sup> According to survey of dolomitic lime manufacturer there was no dolomitic lime production in the period 1990-1995 (production of dolomitic lime started in 1996)

CO<sub>2</sub> emissions from quicklime and dolomitic lime production are presented in the Table 4.2.5.

Table 4.2-5: CO<sub>2</sub> emissions from quicklime and dolomitic lime production (1990-2005)

Year	Quicklime production (Gg CO <sub>2</sub> )	Dolomitic lime production (Gg CO <sub>2</sub> )
1990	159.78	0.00
1991	96.15	0.00
1992	54.49	0.00
1993	60.25	0.00
1994	59.65	0.00
1995	62.27	0.00
1996	45.44	33.71
1997	51.53	50.09
1998	57.21	48.56
1999	54.26	48.31
2000	61.47	62.79
2001	81.21	62.27
2002	77.68	86.30
2003	72.89	88.11
2004	120.91	53.43
2005	124.11	72.32

The resulting emissions of CO<sub>2</sub> from Lime Production in the period 1990-2005 are presented in the Figure 4.2-2.




Figure 4.2-2: Emissions of CO<sub>2</sub> from Lime Production (1990-2005)

The methodology for calculating SO<sub>2</sub> emissions from Lime Production is not available in *Revised 1996 IPCC Guidelines*. Process (non-combustion) SO<sub>2</sub> emissions depend on the sulphur content and mineralogical form of the stone feed, the quality of the lime produced and the type of kiln. Until more information becomes available, it is recommended that only emissions from fuel combustion (which are presented in the chapter on emissions from energy sources) are considered.

#### 4.2.2.3. Uncertainties and time-series consistency

Uncertainties contained in CO<sub>2</sub> estimates are due to provided default emission factors which assume 100 percent of CaO in lime (in some cases purity may range from 85 to 95 percent depending on lime type). Emissions estimation using default emission factors lead to overestimation of CO<sub>2</sub> emission, but at the moment there are no enough information concerning to purity of lime. It is only known that additive quantity may amounts on an average 1 to 3 percent. For industrial purpose, limestone usually consists of 97 to 99 percent of CaCO<sub>3</sub>, whereas dolomite mainly contains 40 to 43 percent of MgCO<sub>3</sub> and 57 to 60 percent of CaCO<sub>3</sub>.

Uncertainty estimate associated with default emission factors amounts 15 percent, accordingly to value recommended in *Good Practice Guidance*. Uncertainty estimate associated with activity data amounts 7.5 percent (5 to 10 percent), based on expert judgements since statistics and manufacturers have not been particularly assessed the uncertainties.

Emissions from Lime Production have been calculated using the same method and data sets for every year in the time series.

#### 4.2.2.4. Source specific QA/QC and verification

During the preparation of the inventory submission activities related to quality control were mainly focused on completeness and consistency of emission estimates and on proper use of notation keys in the CRF tables. After preparation of final draft of this chapter an audit was carried out to check selected activities from Tier 1 General inventory level QC procedures.

#### 4.2.2.5. Source specific recalculations

There are no source-specific recalculations in sub-sector Lime Production.

#### 4.2.2.6. Source-specific planned improvements

For the purpose of accurate calculation of national emission factors for quicklime and dolomitic lime production, knowledge of CaO and MgO content in raw materials which are used for lime production is favourably and plans to investigate, as well as CaO and MgO content in lime.

### 4.2.3. LIMESTONE AND DOLOMITE USE

#### 4.2.3.1. Source category description

Limestone ( $\text{CaCO}_3$ ) and dolomite ( $\text{CaCO}_3 \cdot \text{MgCO}_3$ ) are basic raw materials having commercial applications in a number of industries including metal production, glass and ceramic manufacture, refractory materials manufacture, chemical, agriculture, construction and environmental pollution control. For some of these applications carbonates are sufficiently heated to high temperature as part of the process to generate  $\text{CO}_2$  as a by-product. The major utilization of dolomite in Croatia is in glass, ceramic and refractory materials manufacture as well as the limestone use in the pig iron production (during 1990 and 1991).

#### 4.2.3.2. Methodological issues

Emissions of  $\text{CO}_2$  from use of limestone and dolomite have been calculated by multiplying annual consumption of raw material in processes (limestone/dolomite) by emission factors, which are based on a ratio between  $\text{CO}_2$  and limestone/dolomite used in a particular process. Emissions of  $\text{CO}_2$  from the use of limestone have been estimated by using emission factor which equals 440 kg  $\text{CO}_2$ /tonne limestone. Emissions of  $\text{CO}_2$  from the use of dolomite have been estimated by using emission factor which equals 477 kg  $\text{CO}_2$ /tonne dolomite, assuming 100 percent purity of raw material (*Revised 1996 IPCC Guidelines*).


The activity data for limestone use in the pig iron production for the 1990 and 1991 were collected by EKONERG from survey of iron manufacturer.

The activity data for dolomite use in glass, ceramic and refractory materials manufacture in the period 1990-1995 were extracted from Monthly Industrial Reports published by Central Bureau of Statistics, Department of Manufacturing and Mining. National classification of activities distinguished dolomite use in glass, ceramic and refractory materials manufacture in that period. After this period national classification of activities did not distinguish dolomite use in abovementioned activities and because of that, AD was collected by survey of manufacturer. The activity data for dolomite use in glass manufacture in the period 1996-2005 were collected by EKONERG from survey of glass manufacturer. Some of these activities (from the period 1990-1995) were halted in the meantime. According to statistical data and data from survey there was no limestone use in abovementioned processes (see Table 4.2-6).

Table 4.2-6: Limestone and dolomite use (1990-2005)

Year	Limestone use (tonnes)	Dolomite use (tonnes)
1990	41816	52031
1991	12037	40452
1992	0	22091
1993	0	20134
1994	0	32504
1995	0	23461
1996	0	17827
1997	0	15191
1998	0	18028
1999	0	16666
2000	0	17634
2001	0	19364
2002	0	20167
2003	0	24687
2004	0	24141
2005	0	25269

The resulting emissions of CO<sub>2</sub> from Limestone and Dolomite Use in the period 1990-2005 are presented in the Figure 4.2-3.

Figure 4.2-3: Emissions of CO<sub>2</sub> from Limestone and Dolomite Use (1990-2005)

#### 4.2.3.3. Uncertainties and time-series consistency

Uncertainties in CO<sub>2</sub> estimates are related to possible variations in the chemical composition of limestone and dolomite (carbonates may contain smaller amounts of impurities i.e. magnesia, silica, and sulphur). Uncertainties contained in these estimates are due to provided default emission factor which assume 100 percent purity of raw material.

Uncertainty estimate associated with default emission factors amounts 30 percent, based on expert judgements. Uncertainty estimate associated with activity data amounts 7.5 percent (5 to 10 percent), based on expert judgements since statistics and manufacturers have not been particularly assessed the uncertainties.

Emissions from Limestone and Dolomite Use have been calculated using the same method for every year in the time series. Data sets are different for the period 1990-1995 in relation to the period 1996-2005. As abovementioned, in the period 1990-1995 national classification of activities distinguished dolomite use in glass, ceramic and refractory materials manufacture. After this period national classification of activities did not distinguish dolomite use in abovementioned activities and because of that, AD was collected by survey of manufacturer. Some of these activities (from the period 1990-1995) were halted in the meantime, and there is no possibility to collect AD by the same data sets, for entire period.

#### **4.2.3.4. Source specific QA/QC and verification**

During the preparation of the inventory submission activities related to quality control were mainly focused on completeness and consistency of emission estimates and on proper use of notation keys in the CRF tables. After preparation of final draft of this chapter an audit was carried out to check selected activities from Tier 1 General inventory level QC procedures.

#### **4.2.3.5. Source specific recalculations**

There are no source-specific recalculations in sub-sector Limestone and Dolomite Use.

#### **4.2.3.6. Source-specific planned improvements**

For the purpose of accurate calculation of national emission factors for limestone and dolomite use, knowledge of chemical composition of dolomite which is used as raw materials in abovementioned commercial applications (glass, ceramic and refractory materials manufacture) is favourably and plans to investigate.

### **4.2.4. SODA ASH PRODUCTION AND USE**

#### **4.2.4.1. Source category description**

Soda ash (sodium carbonate,  $\text{Na}_2\text{CO}_3$ ) is a white crystalline solid that is commercially used as a raw material in a large number of industrial processes including glass and ceramic manufacture, soap and detergents, pulp and paper production and water treatment.

According to Department of Manufacturing and Mining (Central Bureau of Statistics) there was not any significant production, both natural and synthetic, of soda ash in Croatia in the period 1990-2005. Therefore, only  $\text{CO}_2$  emissions arising in soda ash consumption in glass and ceramic manufacture, and in the production of soap and detergents, have been estimated.

#### **4.2.4.2. Methodological issues**


Emissions of  $\text{CO}_2$  from the soda ash use have been calculated by multiplying annual consumption of soda ash by emission factor, which is based on a ratio between  $\text{CO}_2$  and soda ash used. Default emission factor equals 415 kg  $\text{CO}_2$ /tonne soda ash has been used (*Revised 1996 IPCC Guidelines*).

The activity data for soda ash use in glass and ceramic manufacture, and in the production of soap and detergents in the period 1990-1995, were extracted from Monthly Industrial Reports published by Central Bureau of Statistics, Department of Manufacturing and Mining. National classification of activities distinguished soda ash use in glass and ceramic manufacture and in the production of soap and detergents in that period. After this period national classification of activities did not distinguish soda ash use in abovementioned activities and because of that, AD was collected by survey of manufacturer. The activity data for soda ash use in glass manufacture in the period 1996-2005 were collected by EKONERG from survey of glass manufacturer (see Table 4.2-7).

*Table 4.2-7: Soda ash use (1990-2005)*

Year	Soda ash use (tonnes)
1990	62024
1991	52415
1992	35376
1993	30202
1994	36659
1995	34668
1996	27493
1997	23320
1998	27694
1999	25538
2000	26536
2001	29818
2002	29446
2003	35335
2004	39821
2005	41498

The resulting emissions of CO<sub>2</sub> from Soda Ash Use in the period 1990-2005 are presented in the Figure 4.2-4.



*Figure 4.2-4: Emissions of CO<sub>2</sub> from Soda Ash Use (1990-2005)*

#### **4.2.4.3. Uncertainties and time-series consistency**

Emissions of CO<sub>2</sub> from soda ash use are dependent upon a type of end-use processes involved. Specific information characterizing the emissions from particular end-use process is not available. Therefore, uncertainties are related primarily to the accuracy of the emission factor.

Uncertainty estimate associated with default emission factors amounts 30 percent, based on expert judgements. Uncertainty estimate associated with activity data amounts 7.5 percent (5 to 10 percent), based on expert judgements since statistics and manufacturers have not been particularly assessed the uncertainties.

Emissions from Soda Ash Use have been calculated using the same method for every year in the time series. Data sets are different for the period 1990-1995 in relation to the period 1996-2005. As abovementioned, in the period 1990-1995 national classification of activities distinguished soda ash use in glass and ceramic manufacture and in the production of soap and detergents. After this period national classification of activities did not distinguish soda ash use in abovementioned activities and because of that, AD was collected by survey of manufacturer. Some of these activities (from the period 1990-1995) were halted in the meantime, and there is no possibility to collect AD by the same data sets, for entire period.

#### **4.2.4.4. Source-specific QA/QC and verification**

During the preparation of the inventory submission activities related to quality control were mainly focused on completeness and consistency of emission estimates and on proper use of notation keys in the CRF tables. After preparation of final draft of this chapter an audit was carried out to check selected activities from Tier 1 General inventory level QC procedures.

#### **4.2.4.5. Source specific recalculations**

There are no source-specific recalculations in sub-sector Soda Ash Use.

#### **4.2.4.6. Source-specific planned improvements**

For the purpose of accurate calculation of national emission factors, investigation of specific information characterizing the emissions from particular end-use processes is favourably and plans to investigate.

### **4.2.5. PRODUCTION AND USE OF MISCELLANEOUS MINERAL PRODUCTS**

#### **4.2.5.1. Source category description**

There are several mineral production processes which caused emissions of indirect GHGs: Asphalt Roofing Production, Road Paving with Asphalt and Glass Manufacturing.



#### **4.2.5.2. Methodological issues**

##### **Asphalt Roofing Production**

Asphalt roofing production includes production of asphalt roofing and process of asphalt blowing. Emissions of indirect GHGs have been calculated by multiplying annual produced quantities with related emission factor provided by *Revised 1996 IPCC Guidelines*.

For indirect GHGs emissions estimation in the Asphalt Roofing Production the emission factors of 0.049 kg NMVOC/tonne asphalt roofing and 0.0095 kg CO/tonne asphalt roofing have been applied. In the Asphalt Blowing process the emission factor of 2.4 kg NMVOC/tonne asphalt blown has been applied.

The annual produced quantities were extracted from Monthly Industrial Reports published by Central Bureau of Statistics, Department of Manufacturing and Mining. The resulting emissions of indirect GHGs from Asphalt Roofing Production Processes in the period 1990-2005 are presented in the review on indirect GHG emissions from non-energy industrial processes.

##### **Road Paving with Asphalt**

Emissions of indirect GHGs from Road Paving with Asphalt include emissions from asphalt plant, from road surfacing operations and from subsequent road surface. Emissions of indirect GHGs have been calculated by multiplying annual produced quantities of asphalt with related emission factor provided by *Revised 1996 IPCC Guidelines*.

For NMVOC emissions estimation from Asphalt Plant the emission factor of 0.023 kg NMVOC/tonne asphalt has been applied. The emission factor of 320 kg NMVOC/tonne asphalt has been applied for NMVOC emissions estimation from Road Surface.

The annual produced quantities of asphalt were extracted from Monthly Industrial Reports published by Central Bureau of Statistics, Department of Manufacturing and Mining. The resulting emissions of indirect GHGs from Road Paving with Asphalt in the period 1990-2005 are presented in the review on indirect GHG emissions from non-energy industrial processes.

##### **Glass Manufacturing**

Emissions from Container Glass Production and Flat Glass Production have been calculated by multiplying annual produced quantities of container and flat glass with emission factor provided by *Revised 1996 IPCC Guidelines*. The emission factor of 4.5 kg NMVOC/tonne glass has been applied.

The annual produced quantities of glass were extracted from Monthly Industrial Reports published by Central Bureau of Statistics, Department of Manufacturing and Mining. The resulting emissions of NMVOC from Glass Manufacturing in the period 1990-2005 are presented in the review on indirect GHG emissions from non-energy industrial processes.

#### **4.2.5.3. Uncertainties and time-series consistency**

Uncertainties related to emissions of indirect GHGs are related primarily to the accuracy of the emission factor. *Good Practice Guidance* didn't recommend uncertainty estimate associated with default emission factors for Production and Use of Miscellaneous Mineral Products. Uncertainties associated with default emission factors and activity data were not estimated for Production and Use of Miscellaneous Mineral Products.

Emissions from Production and Use of Miscellaneous Mineral Products have been calculated using the same method and data sets for every year in the time series.

#### **4.2.5.4. Source-specific QA/QC and verification**

During the preparation of the inventory submission activities related to quality control were mainly focused on completeness and consistency of emission estimates and on proper use of notation keys in the CRF tables. After preparation of final draft of this chapter an audit was carried out to check selected activities from Tier 1 General inventory level QC procedures.

#### **4.2.5.5. Source specific recalculations**

There are no source-specific recalculations because only NMVOC emissions are calculated in abovementioned sub-sectors. The *IPCC Guidelines* do not provide methodologies for calculation of CO<sub>2</sub> emission from these sources.

#### **4.2.5.6. Source-specific planned improvements**

For the purpose of accurate calculation of national emission factors, analyze and investigation of specific information related to type of asphalt roofing production processes and type of asphalt as well as amounts of diluent which are used in asphalt production plans to achieve.

### 4.3. CHEMICAL INDUSTRY (CRF 2.B.)

#### 4.3.1. AMMONIA PRODUCTION

##### 4.3.1.1. Source category description

Ammonia is produced by catalytic steam reforming of natural gas in which hydrogen is chemically separated from the natural gas and combined with nitrogen to produce ammonia ( $\text{NH}_3$ ). Carbon dioxide which is formed from carbon monoxide in CO shift converter is removed by using two methods: monoethanolamine scrubbing and hot potassium scrubbing. After absorbing the  $\text{CO}_2$ , the amine solution is preheated and regenerated which results in removing the  $\text{CO}_2$  by steam stripping and then by heating. The  $\text{CO}_2$  is either vented to the atmosphere or used as a feedstock in other parts of the plant complex (for production of UREA or dry ice).

##### 4.3.1.2. Methodological issues

Emissions of  $\text{CO}_2$  from ammonia production have been calculated by multiplying annual consumption of natural gas used as a feedstock in process by carbon content of natural gas and molecular weight ratio between  $\text{CO}_2$  and carbon (Tier 1a, *Revised 1996 IPCC Guidelines*).

Data on consumption and composition of natural gas (see Table 4.3-1) used as a feedstock in a process were collected by EKONERG from survey of ammonia manufacturer (Petrokemija d.d Fertilizer Company Kutina) and cross-checked with ammonia production data from Monthly Industrial Reports published by Central Bureau of Statistics, Department of Manufacturing and Mining. Carbon content of gas ( $\text{kg C/m}^3$ ) has been estimated from volume fraction of  $\text{CH}_4$ ,  $\text{C}_2\text{H}_6$ ,  $\text{C}_3\text{H}_8$ ,  $\text{C}_4\text{H}_{10}$ ,  $\text{C}_5\text{H}_{12}$ ,  $\text{CO}_2$  and  $\text{N}_2$  in natural gas.

Table 4.3-1: Consumption and composition of gas in Ammonia Production (1990-2005)

Year	Gas consumption ( $\text{m}^3$ )	Carbon content of gas ( $\text{kg C/m}^3$ )
1990	242905233	0.5519
1991	230492226	0.5579
1992	299567927	0.5524
1993	238269046	0.5395
1994	239717137	0.5401
1995	232773362	0.5423
1996	254116356	0.5395
1997	277311935	0.5372
1998	207973360	0.5373
1999	262772017	0.5388
2000	266433375	0.5377
2001	214441408	0.5416
2002	193045364	0.5421
2003	216859822	0.5431
2004	264367950	0.5391
2005	259004302	0.5383

The resulting emissions of CO<sub>2</sub> from Ammonia Production in the period 1990-2005 are presented in the Figure 4.3-1.




Figure 4.3-1: Emissions of CO<sub>2</sub> from Ammonia Production (1990-2005)

Emissions of CO<sub>2</sub> from ammonia production have been calculated by multiplying annual consumption of natural gas used as a feedstock in process by carbon content of natural gas and molecular weight ratio between CO<sub>2</sub> and carbon (Tier 1a, *Revised 1996 IPCC Guidelines*). Explanation is provided in CRF tables.

#### 4.3.1.3. Uncertainties and time-series consistency

According to *Revised 1996 IPCC Guidelines* the most accurate method of emissions estimation is based on the consumption and composition of natural gas used as a feedstock in the process<sup>2</sup>.

IEF has been calculated according to proposed equation in CRF tables (Table 2(1).A-Gs1). Some fluctuations and inconsistencies between calculated IEF and carbon contents of natural gas, which are registered in the NIR 2006 Review, are caused because IEF has been calculated by data on ammonia production and calculated emissions. As abovementioned, emissions have been calculated by data on consumption of natural gas used as a feedstock in process and carbon content of natural gas (Tier 1a, *Revised 1996 IPCC Guidelines*). Tier 1b method (*Revised 1996 IPCC Guidelines*), which is based on ammonia production, with proposed default IEF of 1.5 t CO<sub>2</sub>/t NH<sub>3</sub> produced, offer different emissions calculations than Tier 1a method. For example, the decrease of 4.1 percent in CO<sub>2</sub> emissions calculation for 1990-1991 by Tier 1a method is not consistent with the increase of 0.7 percent calculated by Tier 1b method. Also, the overall increase in CO<sub>2</sub> emissions calculated by Tier 1a method over the period 1990-2005 which amounts 4.0 percent is not correspond to the increase in CO<sub>2</sub> emissions calculated by Tier 1b method which amounts 15.5 percent. These inconsistencies are resulted by using of different method and consequently Tier 1a method is applied as more accurate than Tier 1b method.

<sup>2</sup> In order to avoid double counting, the quantities and composition of gas used as a feedstock have been separately reported from the quantities used as fuel in the ammonia production process. The latter were reported in the Energy Chapter.

Also, there are some uncertainties concerning to use of CO<sub>2</sub> as a feedstock in downstream manufacturing processes, in the production of urea, dry ice and fertilizer. According to *Revised 1996 IPCC Guidelines* no account should consequently be taken for intermediate binding of CO<sub>2</sub> in production of urea, dry ice and fertilizer.

Uncertainty estimate associated with emission factors amounts 5 percent, accordingly to value recommended in *Good Practice Guidance*. Uncertainty estimate associated with activity data amounts 3 percent (1 to 5 percent), based on expert judgements since statistics and manufacturers have not been particularly assessed the uncertainties.

Emissions from Ammonia Production have been calculated using the same method and data sets for every year in the time series.

#### **4.3.1.4. Source-specific QA/QC and verification**

During the preparation of the inventory submission activities related to quality control were mainly focused on completeness and consistency of emission estimates and on proper use of notation keys in the CRF tables.

After preparation of final draft of this chapter an audit was carried out to check selected activities from Tier 1 General inventory level QC procedures and Tier 2 source-specific QC procedures. Ammonia Production is one of the key source categories in Industrial Processes. Regarding to Tier 2 activities, emission factors and activity data were checked for key source categories. Emissions of CO<sub>2</sub> from ammonia production were estimated using Tier 1a method which is based on gas consumption and could be considered as a *good practice*.

#### **4.3.1.5. Source-specific recalculations**

There are no source-specific recalculations in sub-sector Ammonia Production.

#### **4.3.1.6. Source-specific planned improvements**

Since Ammonia Production is a key source category, more detailed information about use of CO<sub>2</sub> as a feedstock in downstream manufacturing processes, in the production of urea, dry ice and fertilizer are planned to investigate.

### **4.3.2. NITRIC ACID PRODUCTION**

#### **4.3.2.1. Source category description**

There is one manufacturer of nitric acid in Croatia, with dual pressure type of production process, according to the pressure used in the oxidation and absorption stages. Ammonia, which is used as a feedstock, is vaporized, mixed with air and burned over a platinum/rhodium alloy catalyst. Nitrogen monoxide is formed and oxidized to nitrogen dioxide at medium pressures and absorbed in water at high pressure to give nitric acid. During oxidation stage, nitrogen and nitrous oxide are formed as a by-product and released from reactor vents into the

atmosphere. There is no abatement technology installed at the plant. Nitric acid is used in the manufacture of fertilizers.

#### 4.3.2.2. Methodological issues

Emissions of N<sub>2</sub>O from nitric acid production have been calculated by multiplying annual nitric acid production by emission factor which reflects the process type, i.e. dual pressure type. According to *Good Practice Guidance* emission factor given for European designed dual pressure plants is in the range from 8 to 10 kg N<sub>2</sub>O/tonne nitric acid. Emission factor was determined as mean value of estimated range, i.e. 9 kg N<sub>2</sub>O/tonne nitric acid. Data on nitric acid production (see Table 4.3-2) were collected by EKONERG from survey of nitric acid manufacturer and cross-checked with nitric acid production data from Monthly Industrial Reports published by Central Bureau of Statistics, Department of Manufacturing and Mining.

*Table 4.3-2: Nitric acid production (1990-2005)*

Year	Nitric acid production (tonnes)
1990	332459
1991	291997
1992	381797
1993	287805
1994	311236
1995	299297
1996	278683
1997	292892
1998	220509
1999	260198
2000	306201
2001	257534
2002	249992
2003	235645
2004	287567
2005	280746

The resulting emissions of N<sub>2</sub>O from Nitric Acid Production in the period 1990-2005 are presented in the Figure 4.3-2.




Figure 4.3-2: Emissions of N<sub>2</sub>O from Nitric Acid Production (1990-2005)

#### 4.3.2.3. Uncertainties and time-series consistency

The main uncertainties concerning the emissions of N<sub>2</sub>O from nitric acid production are due to applied emission factor.

As mentioned before the process of nitric acid production in Croatia is European designed dual pressure type and because none of the emission factors proposed by *Revised 1996 IPCC Guidelines* correspond to plant type default emission factor was taken from *Good Practice Guidance*<sup>3</sup>.

Uncertainty estimate associated with default emission factors amounts 30 percent, based on expert judgements. Uncertainty estimate associated with activity data amounts 3 percent (1 to 5 percent), based on expert judgements since statistics and manufacturers have not been particularly assessed the uncertainties.

Emissions from Nitric Acid Production have been calculated using the same method and data sets for every year in the time series.

#### 4.3.2.4. Source-specific QA/QC and verification

During the preparation of the inventory submission activities related to quality control were mainly focused on completeness and consistency of emission estimates and on proper use of notation keys in the CRF tables.

After preparation of final draft of this chapter an audit was carried out to check selected activities from Tier 1 General inventory level QC procedures and Tier 2 source-specific QC procedures. Regarding to Tier 2 activities, emission factors and activity data were checked for key source categories. Nitric Acid Production is one of the key source categories in Industrial

<sup>3</sup> *IPCC Guidelines* provide emission factor for medium pressure plants in the range of 6 to 7.5 kg N<sub>2</sub>O/t nitric acid which could be considered as nearest which correspond to plant type. *Good Practice Guidance* provide emission factor for European designed, dual pressure, double absorption plant, in the range of 8 to 10 kg N<sub>2</sub>O/t nitric acid.

Processes. Emissions of N<sub>2</sub>O from nitric acid production were based on default emission factor from IPCC Good Practice Guidance and annual amount of nitric acid production. It is a *good practice* to use direct emission measurement for national emission factor calculation.

#### 4.3.2.5. Source-specific recalculations

There are no source-specific recalculations in sub-sector Nitric Acid Production.

#### 4.3.2.6. Source-specific planned improvements

Since Nitric Acid Production is a key source category, more detailed information about using of direct emission measurement for calculation of national emission factor are planned to investigate. Furthermore, this data are not available since CEM system is not installed and manufacturer is not obliged yet to conduct spot measurement according to relevant regulation.

### 4.3.3. PRODUCTION OF OTHER CHEMICALS

#### 4.3.3.1. Source category description

The production of other chemicals such as carbon black, coke, and some petrochemicals (ethylene, dichlorethylene, and styrene) can be sources of methane emissions. Although most methane sources from industrial processes individually are small, collectively they may be significant.

#### 4.3.3.2. Methodological issues

Emissions of CH<sub>4</sub> from the production of other chemicals have been calculated by multiplying an annual production of each chemical with related emission factor provided by *Revised 1996 IPCC Guidelines*. The annual production of chemicals (see Table 4.3-3) was extracted from Monthly Industrial Reports published by Central Bureau of Statistics, Department of Manufacturing and Mining.

Table 4.3-3: Production of other chemicals (1990-2005)

Year	Carbon black (tonnes)	Ethylene (tonnes)	Dichloro- ethylene (tonnes)	Styrene (tonnes)	Coke (tonnes)
1990	30624	72631	72653	8923	556084
1991	18783	66871	68325	6376	441584
1992	13479	68318	92089	1381	409371
1993	17123	68634	79608	0	420676
1994	21468	65285	97528	0	276854
1995	27185	67547	84374	0	0
1996	26735	64782	48630	0	0
1997	24214	63554	26264	0	0
1998	22165	60148	31308	0	0
1999	17589	60295	47686	0	0



Table 4.3-3: Production of other chemicals (1990-2005), cont.

Year	Carbon black (tonnes)	Ethylene (tonnes)	Dichloro-ethylene (tonnes)	Styrene (tonnes)	Coke (tonnes)
2000	20029	38918	71364	0	0
2001	21180	46632	64442	0	0
2002	19385	43554	0	0	0
2003	21497	41252	0	0	0
2004	20271	49886	0	0	0
2005	18498	50263	0	0	0

The resulting emissions of CH<sub>4</sub> from Production of Other Chemicals in the period 1990-2005 are reported in Table 4.3-4.

Table 4.3-4: Emissions of CH<sub>4</sub> from Production of Other Chemicals (1990-2005)

Year	Emissions of CH <sub>4</sub> from production of other chemicals (Gg)				
	Carbon black	Ethylene	Dichloro-ethylene	Styrene	Coke
1990	0.34	0.07	0.03	0.04	0.28
1991	0.21	0.07	0.03	0.03	0.22
1992	0.15	0.07	0.04	0.01	0.20
1993	0.19	0.07	0.03	0.00	0.21
1994	0.24	0.07	0.04	0.00	0.14
1995	0.30	0.07	0.03	0.00	0.00
1996	0.29	0.06	0.02	0.00	0.00
1997	0.27	0.06	0.01	0.00	0.00
1998	0.24	0.06	0.01	0.00	0.00
1999	0.19	0.06	0.02	0.00	0.00
2000	0.22	0.04	0.03	0.00	0.00
2001	0.23	0.05	0.03	0.00	0.00
2002	0.21	0.04	0.00	0.00	0.00
2003	0.24	0.04	0.00	0.00	0.00
2004	0.22	0.05	0.00	0.00	0.00
2005	0.20	0.05	0.00	0.00	0.00

Emissions of indirect GHGs from Production of Other Chemicals have been calculated by multiplying an annual production of each chemical with related emission factor provided by *Revised 1996 IPCC Guidelines*.

The resulting emissions of indirect GHGs from Production from Other Chemicals in the period 1990-2005 are presented in the review on indirect GHG emissions from non-energy industrial processes.

#### 4.3.3.3. Uncertainties and time-series consistency

Uncertainty estimate associated with default emission factor for CH<sub>4</sub> emissions amounts 30 percent, based on expert judgements. Uncertainty estimate associated with activity data for CH<sub>4</sub>

emissions amounts 7.5 percent (5 to 10 percent), based on expert judgements since statistics and manufacturers have not been particularly assessed the uncertainties.

Emissions from Production from Other Chemicals have been calculated using the same method and data sets for every year in the time series.

#### **4.3.3.4. Source-specific QA/QC and verification**

During the preparation of the inventory submission activities related to quality control were mainly focused on completeness and consistency of emission estimates and on proper use of notation keys in the CRF tables. After preparation of final draft of this chapter an audit was carried out to check selected activities from Tier 1 General inventory level QC procedures.

#### **4.3.3.5. Source-specific recalculations**

In the previous report errors were done during activity data compilation - units of activity data were not in correct values (e.g. kilogramme instead of tonnes). Thereupon, CH<sub>4</sub> emissions are recalculated for the years 2003 and 2004.

#### **4.3.3.6. Source-specific planned improvements**

For the purpose of accurate emission calculations, Croatia planned to analyze specific chemical production processes.

## 4.4. METAL PRODUCTION (CRF 2.C.)

### 4.4.1. IRON AND STEEL PRODUCTION

#### 4.4.1.1. Source category description

Primary production of pig iron in blast furnace was halted in 1992.

Steel production in electric arc furnaces (EAF) are used to produce carbon and alloy steel. The input material to EAFs is 100 percent scrap. Cylindrical lined EAFs are equipped with carbon electrodes. Alloying agents and fluxing materials (limestone) are added. Electric current of opposite polarity electrodes generates heat between the electrodes and through the scrap. The operations which generate emissions during the EAF steelmaking process are melting, refining, charging scrap, tapping steel and dumping slag. During the melting phase carbon electrodes are kept above the steel melt and the electrical arc oxidises the carbon to CO or CO<sub>2</sub>.

#### 4.4.1.2. Methodological issues

##### Pig Iron Production

Emissions of CO<sub>2</sub> have been calculated by multiplying annual production of pig iron by the emission factor proposed by *Revised 1996 IPCC Guidelines* (1.6 tonnes CO<sub>2</sub>/tonne pig iron produced). The emission factor applied was assumed to be applicable to both pig iron production and integrated pig iron and steel production.

The activity data for pig iron were extracted from Monthly Industrial Reports published by Central Bureau of Statistics, Department of Manufacturing and Mining and cross-checked with iron and steel manufacturer<sup>4</sup>.

The resulting emission of CO<sub>2</sub> from Pig Iron Production in 1990 was amounted 335000 tonnes. In 1991 about 111000 tonnes of CO<sub>2</sub> was emitted. CO<sub>2</sub> emissions are not included in Metal Production to avoid double-counting. These emissions are included in Energy sector because Coke Oven Coke used in blast furnace is given in energy balance.

##### Steel Production

Emissions of CO<sub>2</sub> have been calculated by multiplying annual steel production with related emission factor provided by *Revised 1996 IPCC Guidelines*. The carbon emission factor is based on carbon loss from the electrode. Accordingly to value recommended in *Good Practice Guidance* for carbon released from consumed electrodes (roughly 1-1.5 kg carbon/tonne steel), the arithmetic mean has been taken (1.25 kg carbon/tonne steel) and emission factor of 4.58 kg CO<sub>2</sub>/tonne steel has been applied.

---


<sup>4</sup> It should be noticed that blast furnaces were closed at the end of 1991 mainly due to war activities near the location of iron and steel plant.

The activity data for steel production (see Table 4.4-1) were extracted from Monthly Industrial Reports published by Central Bureau of Statistics, Department of Manufacturing and Mining.

*Table 4.4-1: Steel production (1990-2005)*

Year	Steel production (tonnes)
1990	189368
1991	171147
1992	119733
1993	101942
1994	32674
1995	17021
1996	46424
1997	70660
1998	104854
1999	77213
2000	70998
2001	57963
2002	33839
2003	42235
2004	85947
2005	73640

The resulting emissions of CO<sub>2</sub> from Steel Production in the period 1990-2005 are presented in the Figure 4.4-1.



*Figure 4.4-1: Emissions of CO<sub>2</sub> from Steel Production (1990-2005)*

#### 4.4.1.3. Uncertainties and time-series consistency

The main uncertainties concerning the emission of CO<sub>2</sub> from steel production are due to applied emission factor. The use of plant-specific emission factors would minimize uncertainty, but these factors were not available in adequate form.

Uncertainty estimate associated with default emission factors amounts 30 percent, based on expert judgements. Uncertainty estimate associated with activity data amounts 7.5 percent (5 to

10 percent), based on expert judgements since statistics and manufacturers have not been particularly assessed the uncertainties.

Emissions from Steel Production have been calculated using the same method and data sets for every year in the time series.

#### **4.4.1.4. Source specific QA/QC and verification**

During the preparation of the inventory submission activities related to quality control were mainly focused on completeness and consistency of emission estimates and on proper use of notation keys in the CRF tables. After preparation of final draft of this chapter an audit was carried out to check selected activities from Tier 1 General inventory level QC procedures.

#### **4.4.1.5. Source-specific recalculations**

There are no source-specific recalculations in sub-sector Steel Production.

#### **4.4.1.6. Source-specific planned improvements**

For the purpose of accurate calculation of national emission factors, Croatia planned to investigate the plant-specific emission factor to minimize emission calculation uncertainty.

### **4.4.2. FERROALLOYS PRODUCTION**

#### **4.4.2.1. Source category description**

Ferroalloys are alloys of iron and metals such as silicon, manganese and chromium. Similar to emissions from the production of iron and steel, CO<sub>2</sub> is emitted when metallurgical coke is oxidized during a high-temperature reaction with iron and the selected alloying element. Ferroalloys production was halted in 2002.


#### **4.4.2.2. Methodological issues**

Emissions of CO<sub>2</sub> have been calculated by multiplying annual ferroalloys production by material-specific emission factor (1.7 tonnes CO<sub>2</sub>/tonne silicon manganese, 1.6 tonnes CO<sub>2</sub>/tonne ferromanganese and 1.3 tonnes CO<sub>2</sub>/tonne ferrochromium). The activity data for ferroalloys production (see Table 4.4-2) were extracted from Monthly Industrial Reports published by Central Bureau of Statistics, Department of Manufacturing and Mining. Adjustment method proposed by the *Technical Guidance on Methodologies for Adjustments Under Article 5, Paragraph 2, of the Kyoto Protocol*, has been used for calculation of insufficient data for 1999 year. Taking into account the ferroalloys production trend, values for 1998 and 2000 have been used as the pattern for ferroalloys production calculation in 1999 by interpolation method.

Table 4.4-2: Production of ferroalloys (1990-2005)

Year	Ferromanganese (tonnes)	Silicon manganese (tonnes)	Ferrochromium (tonnes)
1990	20535	48561	60859
1991	13053	38365	72845
1992	0	25572	56058
1993	0	8577	28028
1994	562	22071	31704
1995	0	0	26081
1996	0	0	10559
1997	0	0	24231
1998	0	0	11861
1999	0	0	13807
2000	0	0	15753
2001	0	0	361
2002	0	0	0
2003	0	0	0
2004	0	0	0
2005	0	0	0

The resulting emissions of CO<sub>2</sub> from Ferroalloys Production in the period 1990-2005 are presented in the Figure 4.4-2.

Figure 4.4-2: Emissions of CO<sub>2</sub> from Ferroalloys Production (1990-2005)

#### 4.4.2.3. Uncertainties and time-series consistency

Uncertainty estimate associated with default emission factors amounts 30 percent, based on expert judgements. Uncertainty estimate associated with activity data amounts 7.5 percent (5 to 10 percent), based on expert judgements since statistics and manufacturers have not been particularly assessed the uncertainties.

Emissions from Ferroalloys Production have been calculated using the same method and data sets for every year in the time series, except insufficient data for 1999, which was obtained by interpolation method.

#### 4.4.2.4. Source specific QA/QC and verification

During the preparation of the inventory submission activities related to quality control were mainly focused on completeness and consistency of emission estimates and on proper use of notation keys in the CRF tables. After preparation of final draft of this chapter an audit was carried out to check selected activities from Tier 1 General inventory level QC procedures.

#### 4.4.2.5. Source-specific recalculations

In the previous report the production of ferrochromium and associated emissions were insufficient for 1999. In this report data for 1999 was calculated by interpolation method and CO<sub>2</sub> emissions are recalculated.

### 4.4.3. ALUMINIUM PRODUCTION

#### 4.4.3.1. Source category description

Primary aluminium is produced in two steps. First bauxite ore is ground, purified and calcined to produce alumina (Al<sub>2</sub>O<sub>3</sub>). Following this, the alumina is electrically reduced to aluminium by smelting in large pots. This process results in emission of several greenhouse gases including CO<sub>2</sub>, and two PFCs: CF<sub>4</sub> and C<sub>2</sub>F<sub>6</sub>. Primary aluminium production was halted in 1991.

#### 4.4.3.2. Methodological issues

The quantity of CO<sub>2</sub> released was estimated from the production of primary aluminium and the specific consumption of carbon which is oxidized to CO<sub>2</sub> in the process. During alumina reduction using prebaked anodes approximately 1.5 tonnes of CO<sub>2</sub> is emitted for each tonne of primary aluminium produced.

Data on primary aluminium production were collected by EKONERG from voluntary survey of aluminium manufacturer<sup>5</sup>.

The resulting emission of CO<sub>2</sub> from Aluminium Production in 1990 was amounted about 111000 tonnes CO<sub>2</sub>. In 1991 about 76000 tonnes CO<sub>2</sub> was emitted.

PFCs emissions from Aluminium Production could represent a significant source of emissions due to high GWP values. Since only aluminium production statistics were available, emissions of CF<sub>4</sub> and C<sub>2</sub>F<sub>6</sub> were estimated by multiplying annual primary aluminium production with default emission factors provided by *Good Practice Guidance*. Default emission factors equal 1.7 kg/tonne Al for CF<sub>4</sub> and 0.17 kg/tonne Al for C<sub>2</sub>F<sub>6</sub> (Side Worked Prebaked Anodes).

In 1990 about 819000 tonnes CO<sub>2</sub>-eq of CF<sub>4</sub> and 120000 tonnes CO<sub>2</sub>-eq of C<sub>2</sub>F<sub>6</sub> were emitted. In 1991 about 566000 tonnes CO<sub>2</sub>-eq of CF<sub>4</sub> and 83000 tonnes CO<sub>2</sub>-eq of C<sub>2</sub>F<sub>6</sub> were emitted.

---

<sup>5</sup> It should be noticed that primary aluminium production (electrolysis) were closed at the end of 1991 mainly due to war activities near the location of aluminium plant.

Occasionally, sulphur hexafluoride (SF<sub>6</sub>) is also used by the aluminium industry as a cover gas for special foundry products. There are no available data on SF<sub>6</sub> consumption in aluminium industry.

The resulting emissions of indirect GHGs from Aluminium Production in the period 1990-1991 are presented in the review on indirect GHG emissions from non-energy industrial processes.

#### **4.4.3.3. Uncertainties and time-series consistency**

Uncertainties related to calculation of CO<sub>2</sub> emissions are primarily due to applied emission factor. Emissions vary depending on the specific technology used by each plant, however evidence suggests that there is little variation in CO<sub>2</sub> emissions from plants utilising similar technology.

A less uncertain method to calculate CO<sub>2</sub> emissions would be based upon the amount of reducing agent, i.e. amount of prebaked anodes used in a process but this information was not available. Nevertheless, it is very likely that use of the technology-specific emission factor, provided by *Revised 1996 IPCC Guidelines*, along with the correct production data produce accurate estimates.

Uncertainty estimate associated with default emission factor for CO<sub>2</sub> emissions amounts 30 percent, based on expert judgements. Uncertainty estimate associated with activity data for CO<sub>2</sub> emissions amounts 3 percent (1 to 5 percent), based on expert judgements since statistics and manufacturer have not been particularly assessed the uncertainties.

More uncertainties are related to calculation of PFCs emissions because continuous emission monitoring was not carried out, and smelter-specific operating parameters were not available. Default emission factors were therefore applied to calculate PFCs emissions. Uncertainty estimate associated with default emission factor for PFCs emissions amounts 50 percent, based on expert judgements. Uncertainty estimate associated with activity data for PFCs emissions amounts 30 percent, based on expert judgements.

Emissions from Aluminium Production have been calculated using the same method and data sets for every year in the time series.



## **4.5. OTHER PRODUCTION (CRF 2.D.)**

### **4.5.1. PULP AND PAPER**

#### **4.5.1.1. Source category description**

Kraft (sulphate) pulping, acid sulphite pulping and neutral sulphite semi-chemical process are three types of paper production processes. Kraft pulping was used in 1990 and acid sulphite pulping was used until 1994 for paper production. After that, only neutral sulphite semi-chemical process has been used for paper production.

#### **4.5.1.2. Methodological issues**

Emissions of indirect GHGs have been calculated by multiplying annual production quantities with related emission factor provided by *Revised 1996 IPCC Guidelines*. For Kraft Pulping following default emission factors have been used: 7 kg SO<sub>2</sub>/tonne dried pulp, 1.5 kg NO<sub>x</sub>/tonne dried pulp, 3.7 kg NMVOC/tonne dried pulp and 5.6 kg CO/tonne dried pulp. For Acid Sulphite Pulping emission factor of 30 kg SO<sub>2</sub>/tonne dried pulp has been used.

According to *Revised 1996 IPCC Guidelines*, only data for emissions estimation from Kraft and Acid Sulphite Process are available. The methods for emission estimation from Neutral Sulphite Semi-Chemical Process are not available and emissions from that process have therefore not been taken into account.

The activity data for pulp and paper production were extracted from Monthly Industrial Reports published by Central Bureau of Statistics, Department of Manufacturing and Mining.

The resulting emissions of indirect GHGs from Pulp and Paper in the period 1990-1994 are presented in the review on indirect GHG emissions from non-energy industrial processes.

#### **4.5.1.3. Uncertainties and time-series consistency**

Uncertainties associated with default emission factors and activity data were not estimated for Pulp and Paper Production.

Emissions from Pulp and Paper have been calculated using the same method and data sets for every year in the time series.

### **4.5.2. FOOD AND DRINK**

#### **4.5.2.1. Source category description**

Emissions of NMVOC from following types of Food and Drink production processes have been calculated: meat, fish and poultry, sugar, margarine and solid cooking fats, cakes, biscuits and cereals, bread, animal feed, coffee roasting, wine, white wine, beer, spirits and brandy.

#### **4.5.2.2. Methodological issues**

Emissions of indirect GHGs from the production of food and drink have been calculated by multiplying annual production quantities with related emission factors provided by *Revised 1996 IPCC Guidelines*.

The activity data for food and drink production were extracted from Monthly Industrial Reports published by Central Bureau of Statistics, Department of Manufacturing and Mining. The resulting emissions of indirect GHGs from Food and Drink in the period 1990-2005 are presented in the review on indirect GHG emissions from non-energy industrial processes.

#### **4.5.2.3. Uncertainties and time-series consistency**

Uncertainties associated with default emission factors and activity data were not estimated for Food and Drink. Emissions from Food and Drink have been calculated using the same method and data sets for every year in the time series.

## 4.6. CONSUMPTION OF HALOCARBONS AND SF<sub>6</sub> (CRF 2.F.)

### 4.6.1. REFRIGERATION AND AIR CONDITIONING EQUIPMENT

#### 4.6.1.1. Source category description

Hydrofluorocarbons (HFCs), perfluorocarbons (PFCs) and sulphur hexafluoride (SF<sub>6</sub>) are synthetic greenhouse gases whose present contribution to greenhouse effect is relatively small comparing to major greenhouse gases but due to their extremely long lifetime and Global Warming Potentials (GWP) they will continue to accumulate in the atmosphere as long as emissions continue.

Emissions are released by the handling and consumption of synthetic greenhouse gases. HFCs (HFC-32, HFC-125, HFC-134a and HFC-143a) are used as substitutes for cooling gases in refrigerating and air-conditioning systems that deplete the ozone layer. There is no production of HFCs in Croatia, therefore all quantities of HFCs are imported. Minor quantities of some substances are exported.

#### 4.6.1.2. Methodological issues

In order to estimate consumption of HFCs in the period 1990-2005 a questionnaires have been sent to trading, service and manufacturing companies previously identified as possible sources of handling or consumption of these compounds. Several institutions such as Ministry of Environmental Protection, Physical Planning and Construction, Customs Department and Central Bureau of Statistics were contacted and asked to provide information on import and export of HFCs as well as information on consumption of each individual gas at the rather detailed level in order to use Tier 2 method (*Revised 1996 IPCC Guidelines*).

Results of a survey were unable to provide certain data in required extent. Also, National Classification of Activities used by Central Bureau of Statistics, in the same manner, does not particularly mark HFCs, PFCs and SF<sub>6</sub>. Customs Departments Tariff Number does not precisely distinguish these compounds from other fluorinated chemicals which are controlled with Montreal Protocol.

The only useful information is those related to import and export of HFCs provided by Ministry of Environmental Protection, Physical Planning and Construction. According to this information potential HFCs emissions, for the period 1990–2005, were calculated by difference of import and export of these gases (Tier 1a method, *Revised 1996 IPCC Guidelines*). Insufficient data were estimated by interpolation/extrapolation method. Croatia does not yet have the necessary data to calculate actual emissions. According to requirement of Regulation on the Greenhouse Gases Emissions Monitoring in the Republic of Croatia (Official gazette No. 1/07), which will go into force in 2008, each sources of HFCs emissions should report required activity data for more accurate emissions estimation.

Croatia plans to choose 1995 as its base year for HFCs emission estimations. Adjustment method "Extrapolation of emissions based on a driver", proposed by the *Technical Guidance on*

*Methodologies for Adjustments under Article 5, Paragraph 2, of the Kyoto Protocol*, has been used for calculation of insufficient emission estimations for HFC 32 (1990-1999), HFC 125 (1990-1995), HFC 134a (1990-1994) and HFC 143a (1990-1995).

Taking into account the emissions trend, the pattern over three years from 2000 to 2002 has been used for calculation of HFC 32 emissions for the period 1990-1999. The pattern over period from 1996 to 2002 has been used for HFC 125 and HFC 143a emissions calculation for the period 1990-1995. The pattern over period from 1995 to 2002 has been used for HFC 134a emissions calculation for the period 1990-1994. Conservativeness factor, proposed by the *Technical Guidance on Methodologies for Adjustments under Article 5, Paragraph 2, of the Kyoto Protocol*, was used for the base year (1990). Data for 2003 and 2004 have been calculated by interpolation method using the pattern over entire time series and data for 2005, which were obtained by Ministry of Environmental Protection, Physical Planning and Construction.

Annual emissions of HFCs, expressed in Gg CO<sub>2</sub>-eq, in the period 1990-2005, are presented in Table 4.6-1.

*Table 4.6-1: Emissions of HFCs (Gg CO<sub>2</sub>-eq) (1990 – 2005)*

Gas	1990	1991	1992	1993	1994	1995	1996	1997
HFC 32	0.11	0.13	0.13	0.12	0.12	0.11	0.11	0.10
HFC 125	20.05	22.99	21.53	20.06	18.59	17.13	22.20	22.18
HFC 134a	8.36	10.47	10.74	11.02	11.29	7.80	2.34	33.44
HFC 143a	14.85	18.12	18.13	18.14	18.15	18.15	35.61	35.57
<b>Total emission (Gg CO<sub>2</sub>-eq)</b>	<b>43.38</b>	<b>51.71</b>	<b>50.52</b>	<b>49.34</b>	<b>48.15</b>	<b>43.20</b>	<b>60.26</b>	<b>91.29</b>

*Table 4.6-1: Emissions of HFCs (Gg CO<sub>2</sub>-eq) (1990 – 2005), cont.*

Gas	1998	1999	2000	2001	2002	2003	2004	2005
HFC 32	0.10	0.09	0.07	0.12	0.06	1.29	1.58	3.25
HFC 125	1.12	1.76	5.35	12.91	13.29	45.09	52.12	98.76
HFC 134a	14.60	4.63	8.92	14.53	14.32	41.80	47.81	82.71
HFC 143a	1.82	2.70	8.82	21.43	21.64	75.52	87.36	164.46
<b>Total emission (Gg CO<sub>2</sub>-eq)</b>	<b>17.64</b>	<b>9.18</b>	<b>23.16</b>	<b>48.99</b>	<b>49.31</b>	<b>163.71</b>	<b>188.87</b>	<b>349.18</b>

#### 4.6.1.3. Uncertainties and time-series consistency

The main uncertainties of HFCs emissions estimation concerning to activity data. Quantities of HFCs contained in various products imported into or exported from a country were difficult to estimate. Also, the application of abovementioned methodology may lead to underestimation or overestimation of potential emissions, depending on whether the majority of HFC containing products is being imported or exported.

Uncertainty estimate associated with estimation of potential emissions of HFC-32, HFC-125, HFC-134a and HFC-143a amounts 70 percent for emission factor and 70 percent for activity data, based on expert judgements.

Emissions from Consumption of HFCs in Refrigeration and Air Conditioning Equipment have been calculated using the same method for every year in the time series. Two sources of information were used for data sets: data were provided by Ministry of Environmental Protection, Physical Planning and Construction and insufficient data were adjusted by interpolation/extrapolation methods.

#### **4.6.1.4. Source-specific QA/QC and verification**

During the preparation of the inventory submission activities related to quality control were mainly focused on completeness and consistency of emission estimates and on proper use of notation keys in the CRF tables. After preparation of final draft of this chapter an audit was carried out to check selected activities from Tier 1 General inventory level QC procedures and Tier 2 source-specific QC procedures. Consumption of HFCs in Refrigeration and Air Conditioning Equipment is one of the key source categories in Industrial Processes. Regarding to Tier 2 activities, emission factors and activity data were checked for key source categories.

#### **4.6.1.5. Source-specific recalculations**

In the previous report, HFCs emissions for the period 1990–1994 (HFC 134a), 1990–1995 (HFC 125, 143a) and 1990–1999 (HFC 32) have not been calculated because of the difficulty in obtaining the relevant activity data for these estimates. In this report, data have been calculated by extrapolation method and HFCs emissions are recalculated.

#### **4.6.1.6. Source-specific planned improvements**

For the purpose of accurate emission calculations, Croatia planned to examine the quantities of HFCs contained in various products imported into or exported from a country. Also, it is essential to adjust National Classification of Activities used by Central Bureau of Statistics in order to particularly mark HFCs, PFCs and SF<sub>6</sub> and Customs Departments Tariff Number to distinguish these compounds from other fluorinated chemicals which are controlled with Montreal Protocol.

According to requirement of Regulation on the Greenhouse Gases Emissions Monitoring in the Republic of Croatia (Official Gazette No. 1/07), which will go into force in 2008, each sources of HFCs emissions should report required activity data for more accurate emissions estimation (Tier 2 method).

### **4.6.2. OTHER CONSUMPTION OF HFCs, PFCs AND SF<sub>6</sub>**

Emissions from Consumption of HFCs for Foam Blowing, Fire Extinguishers, Aerosols/Metered Dose Inhalers, Solvents and Semiconductor Manufacturing have not been calculated because activity data are not available.

A certain amount of SF<sub>6</sub> is contained in electrical equipment used in Croatian National Electricity and KONCAR Electrical Industries Inc. Equipment manufacturers guarantee annual leakage of less than 1 percent, so this information could be used to determine the SF<sub>6</sub> emissions. However, it is still not included in the inventory because the input data are not available.

According to requirement of Regulation on the Greenhouse Gases Emissions Monitoring in the Republic of Croatia (Official Gazette No. 1/07), which will go into force in 2008, each sources of HFCs and SF<sub>6</sub> emissions should report required activity data for more accurate emissions estimation.

#### **4.7. NON - ENERGY USE (CRF 2.G.)**

Non-energy fuel consumptions (fuels used as feedstock) and appropriate emissions, where one part or even the whole carbon is stored in product for a longer time and the other part oxidizes and goes to atmosphere, are included here. The feedstock use of energy carriers occurs in chemical industry (naphta, lubricants, ethane and other), construction industry (bitumen production), and other products such as motor oil, industrial oil, grease etc.

## 4.8. EMISSION OVERVIEW

### 4.8.1. GHG EMISSIONS

Emissions of GHGs from Industrial Processes in the period 1990-2005 are presented in Table 4.8-1.

Table 4.8-1: Emissions of GHGs from Industrial Processes (1990-2005)

Source	Year	GHG	Emission (Gg)	GWP <sup>1</sup>	Emission (Gg CO <sub>2</sub> -eq)	Percent in Industrial Processes	Percent in Total Country Emission
<b>Cement production</b>	1990	CO <sub>2</sub>	1047.47	1	1047.47	26.20	3.32
	1991		655.12		655.12	21.50	2.64
	1992		791.08		791.08	29.08	3.47
	1993		657.36		657.36	30.93	2.86
	1994		811.71		811.71	34.06	3.67
	1995		592.40		592.40	28.71	2.62
	1996		623.43		623.43	29.90	2.68
	1997		750.86		750.86	31.77	3.03
	1998		816.75		816.75	40.68	3.27
	1999		1055.78		1055.78	43.00	4.06
	2000		1233.24		1233.24	43.94	4.77
	2001		1395.03		1395.03	50.53	5.17
	2002		1361.15		1361.15	50.73	4.83
	2003		1371.54		1371.54	48.67	4.59
	2004		1458.69		1458.69	45.86	4.86
	2005		1495.01		1495.01	44.36	4.90
<b>Lime production</b>	1990	CO <sub>2</sub>	159.78	1	159.78	4.00	0.51
	1991		96.15		96.15	3.16	0.39
	1992		54.49		54.49	2.00	0.24
	1993		60.25		60.25	2.83	0.26
	1994		59.65		59.65	2.50	0.27
	1995		62.27		62.27	3.02	0.28
	1996		79.15		79.15	3.80	0.34
	1997		101.63		101.63	4.30	0.41
	1998		105.77		105.77	5.27	0.42
	1999		102.57		102.57	4.18	0.39
	2000		124.25		124.25	4.43	0.48
	2001		143.48		143.48	5.20	0.53
	2002		163.97		163.97	6.11	0.58
	2003		160.99		160.99	5.71	0.54
	2004		174.34		174.34	5.48	0.58
	2005		196.42		196.42	5.83	0.64

Table 4.8-1: Emissions of GHGs from Industrial Processes (1990-2005), cont.

Source	Year	GHG	Emission (Gg)	GWP <sup>1</sup>	Emission (Gg CO <sub>2</sub> -eq)	Percent in Industrial Processes	Percent in Total Country Emission
<b>Limestone and dolomite use</b>	1990	CO <sub>2</sub>	43.22	1	43.22	1.08	0.14
	1991		24.59		24.59	0.81	0.10
	1992		10.54		10.54	0.39	0.05
	1993		9.60		9.60	0.45	0.04
	1994		15.50		15.50	0.65	0.07
	1995		11.19		11.19	0.54	0.05
	1996		8.50		8.50	0.41	0.04
	1997		7.25		7.25	0.31	0.03
	1998		8.60		8.60	0.43	0.03
	1999		7.95		7.95	0.32	0.03
	2000		8.41		8.41	0.30	0.03
	2001		9.24		9.24	0.33	0.03
	2002		9.62		9.62	0.36	0.03
	2003		11.78		11.78	0.42	0.04
	2004		11.52		11.52	0.36	0.04
	2005		12.05		12.05	0.36	0.04
<b>Soda ash production and use</b>	1990	CO <sub>2</sub>	25.74	1	25.74	0.64	0.08
	1991		21.75		21.75	0.71	0.09
	1992		14.68		14.68	0.54	0.06
	1993		12.53		12.53	0.59	0.05
	1994		15.21		15.21	0.64	0.07
	1995		14.39		14.39	0.70	0.06
	1996		11.41		11.41	0.55	0.05
	1997		9.68		9.68	0.41	0.04
	1998		11.49		11.49	0.57	0.05
	1999		10.60		10.60	0.43	0.04
	2000		11.01		11.01	0.39	0.04
	2001		12.37		12.37	0.45	0.05
	2002		12.22		12.22	0.46	0.04
	2003		14.66		14.66	0.52	0.05
	2004		16.53		16.53	0.52	0.06
	2005		17.22		17.22	0.51	0.06
<b>Ammonia production</b>	1990	CO <sub>2</sub>	491.55	1	491.55	12.30	1.56
	1991		471.50		471.50	15.47	1.90
	1992		606.76		606.76	22.31	2.66
	1993		471.34		471.34	22.18	2.05
	1994		474.73		474.73	19.92	2.15
	1995		462.85		462.85	22.43	2.05
	1996		502.68		502.68	24.11	2.16
	1997		546.23		546.23	23.12	2.20
	1998		409.73		409.73	20.41	1.64
	1999		519.13		519.13	21.14	2.00
	2000		525.29		525.29	18.72	2.03
	2001		425.85		425.85	15.43	1.58
	2002		383.72		383.72	14.30	1.36
	2003		431.85		431.85	15.32	1.45
	2004		522.58		522.58	16.43	1.74
	2005		511.21		511.21	15.17	1.68



Table 4.8-1: Emissions of GHGs from Industrial Processes (1990-2005), cont.

Source	Year	GHG	Emission (Gg)	GWP <sup>1</sup>	Emission (Gg CO <sub>2</sub> -eq)	Percent in Industrial Processes	Percent in Total Country Emission
Nitric acid production	1990	N <sub>2</sub> O	2.99	310	927.56	23.20	2.94
	1991		2.63		814.67	26.73	3.28
	1992		3.44		1,065.21	39.16	4.67
	1993		2.59		802.98	37.78	3.49
	1994		2.80		868.35	36.44	3.93
	1995		2.69		835.04	40.46	3.70
	1996		2.51		777.53	37.29	3.34
	1997		2.64		817.17	34.58	3.30
	1998		1.98		615.22	30.64	2.46
	1999		2.34		725.95	29.57	2.79
	2000		2.76		854.30	30.44	3.31
	2001		2.32		718.52	26.03	2.66
	2002		2.25		697.48	26.00	2.47
	2003		2.12		657.45	23.33	2.20
	2004		2.59		802.31	25.22	2.67
	2005		2.53		783.28	23.25	2.57
Production of other chemicals	1990	CH <sub>4</sub>	0.75	21	15.80	0.40	0.05
	1991		0.55		11.49	0.38	0.05
	1992		0.46		9.74	0.36	0.04
	1993		0.50		10.48	0.49	0.05
	1994		0.48		10.06	0.42	0.05
	1995		0.40		8.41	0.41	0.04
	1996		0.38		7.94	0.38	0.03
	1997		0.34		7.15	0.30	0.03
	1998		0.32		6.65	0.33	0.03
	1999		0.27		5.73	0.23	0.02
	2000		0.29		6.04	0.22	0.02
	2001		0.31		6.41	0.23	0.02
	2002		0.26		5.39	0.20	0.02
	2003		0.28		5.83	0.21	0.02
	2004		0.27		5.73	0.18	0.02
	2005		0.25		5.33	0.16	0.02
Steel production	1990	CO <sub>2</sub>	0.87	1	0.87	0.02	0.003
	1991		0.78		0.78	0.03	0.003
	1992		0.55		0.55	0.02	0.002
	1993		0.47		0.47	0.02	0.002
	1994		0.15		0.15	0.01	0.001
	1995		0.08		0.08	0.004	0.0003
	1996		0.21		0.21	0.01	0.001
	1997		0.32		0.32	0.01	0.001
	1998		0.48		0.48	0.02	0.002
	1999		0.35		0.35	0.01	0.001
	2000		0.33		0.33	0.01	0.001
	2001		0.27		0.27	0.01	0.001
	2002		0.15		0.15	0.01	0.001
	2003		0.19		0.19	0.01	0.001
	2004		0.39		0.39	0.01	0.001
	2005		0.34		0.34	0.01	0.001

Table 4.8-1: Emissions of GHGs from Industrial Processes (1990-2005), cont.

Source	Year	GHG	Emission (Gg)	GWP <sup>1</sup>	Emission (Gg CO <sub>2</sub> -eq)	Percent in Industrial Processes	Percent in Total Country Emission
<b>Ferroalloys production</b>	1990	CO <sub>2</sub>	194.53	1	194.53	4.87	0.62
	1991		180.80		180.80	5.93	0.73
	1992		116.35		116.35	4.28	0.51
	1993		51.02		51.02	2.40	0.22
	1994		79.64		79.64	3.34	0.36
	1995		33.91		33.91	1.64	0.15
	1996		13.73		13.73	0.66	0.06
	1997		31.50		31.50	1.33	0.13
	1998		15.42		15.42	0.77	0.06
	1999		17.95		17.95	0.73	0.07
	2000		20.48		20.48	0.73	0.08
	2001		0.47		0.47	0.02	0.002
	2002		0.00		0.00	0.00	0.00
	2003		0.00		0.00	0.00	0.00
	2004		0.00		0.00	0.00	0.00
	2005		0.00		0.00	0.00	0.00
<b>Aluminium production</b>	1990	CO <sub>2</sub>	111.37	1	111.37	3.64	0.35
	1991		76.40		76.40	3.16	0.31
	1992		0.00		0.00	0.00	0.00
	1993		0.00		0.00	0.00	0.00
	1994		0.00		0.00	0.00	0.00
	1995		0.00		0.00	0.00	0.00
	1996		0.00		0.00	0.00	0.00
	1997		0.00		0.00	0.00	0.00
	1998		0.00		0.00	0.00	0.00
	1999		0.00		0.00	0.00	0.00
	2000		0.00		0.00	0.00	0.00
	2001		0.00		0.00	0.00	0.00
	2002		0.00		0.00	0.00	0.00
	2003		0.00		0.00	0.00	0.00
	2004		0.00		0.00	0.00	0.00
	2005		0.00		0.00	0.00	0.00
	1990	CF <sub>4</sub>	0.13	6500	820.44	21.76	2.60
	1991		0.09		562.79	19.44	2.27
	1992		0.00		0.00	0.00	0.00
	1993		0.00		0.00	0.00	0.00
	1994		0.00		0.00	0.00	0.00
	1995		0.00		0.00	0.00	0.00
	1996		0.00		0.00	0.00	0.00
	1997		0.00		0.00	0.00	0.00
	1998		0.00		0.00	0.00	0.00
	1999		0.00		0.00	0.00	0.00
	2000		0.00		0.00	0.00	0.00
	2001		0.00		0.00	0.00	0.00
	2002		0.00		0.00	0.00	0.00
	2003		0.00		0.00	0.00	0.00
	2004		0.00		0.00	0.00	0.00
	2005		0.00		0.00	0.00	0.00

Table 4.8-1: Emissions of GHGs from Industrial Processes (1990-2005), cont.

Source	Year	GHG	Emission (Gg)	GWP <sup>1</sup>	Emission (Gg CO <sub>2</sub> -eq)	Percent in Industrial Processes	Percent in Total Country Emission
Aluminium production	1990	C <sub>2</sub> F <sub>6</sub>	0.013	9200	116.12	3.79	0.37
	1991		0.009		79.66	3.31	0.32
	1992		0.00		0.00	0.00	0.00
	1993		0.00		0.00	0.00	0.00
	1994		0.00		0.00	0.00	0.00
	1995		0.00		0.00	0.00	0.00
	1996		0.00		0.00	0.00	0.00
	1997		0.00		0.00	0.00	0.00
	1998		0.00		0.00	0.00	0.00
	1999		0.00		0.00	0.00	0.00
	2000		0.00		0.00	0.00	0.00
	2001		0.00		0.00	0.00	0.00
	2002		0.00		0.00	0.00	0.00
	2003		0.00		0.00	0.00	0.00
	2004		0.00		0.00	0.00	0.00
	2005		0.00		0.00	0.00	0.00
Consumption of HFCs, PFCs and SF <sub>6</sub> <sup>2</sup>	1990	HFC	<sup>3</sup>	<sup>3</sup>	43.38	1.09	0.14
	1991		<sup>3</sup>		51.71	1.70	0.21
	1992		<sup>3</sup>		50.52	1.86	0.22
	1993		<sup>3</sup>		49.34	2.32	0.21
	1994		<sup>3</sup>		48.15	2.02	0.22
	1995		<sup>3</sup>		43.20	2.09	0.19
	1996		<sup>3</sup>		60.26	2.89	0.26
	1997		<sup>3</sup>		91.29	3.86	0.37
	1998		<sup>3</sup>		17.64	0.88	0.07
	1999		<sup>3</sup>		9.18	0.37	0.04
	2000		<sup>3</sup>		23.16	0.83	0.09
	2001		<sup>3</sup>		48.99	1.77	0.18
	2002		<sup>3</sup>		49.31	1.84	0.17
	2003		<sup>3</sup>		163.71	5.81	0.55
	2004		<sup>3</sup>		188.87	5.94	0.63
	2005		<sup>3</sup>		349.18	10.36	1.15

<sup>1</sup> Time horizon chosen for GWP values is 100 years

<sup>2</sup> consumption of SF<sub>6</sub> is not included because data are not available

<sup>3</sup> HFC 32 (GWP=650); HFC 125 (GWP=2800); HFC 134a (GWP=1300); HFC 143a (GWP=3800) - total emissions of HFCs are presented

#### 4.8.2. INDIRECT GHG EMISSIONS

Many non-energy industrial processes generate emissions of ozone and aerosol precursor gases including carbon monoxide (CO), nitrogen oxides (NO<sub>x</sub>), non-methane volatile organic compounds (NMVOC) and sulphur dioxide (SO<sub>2</sub>) (see Table 4.8-2).

Table 4.8-2: Gases generated from different non-energy industrial process

Gas	Industrial Process
SO <sub>2</sub>	Cement Production
	Production of other chemicals
	Pulp and paper production
NO <sub>x</sub>	Nitric acid production
	Production of other chemicals
	Pulp and paper production
CO	Asphalt Roofing Production
	Ammonia production
	Production of other chemicals
	Pulp and paper production
NMVOC	Asphalt Roofing Production
	Road paving with asphalt
	Glass production
	Production of other chemicals
	Pulp and paper production
	Alcoholic beverage production
	Bread and other food production

Total annual emissions of indirect GHGs in the period 1990-2005 are reported in table 4.8-3.

Table 4.8-3: Emissions of indirect GHGs from Industrial Processes (1990-2005)

Year	SO <sub>2</sub> (Gg)	NO <sub>x</sub> (Gg)	CO (Gg)	NMVOC (Gg)
1990	5.25	0.37	3.11	81.64
1991	3.86	0.30	2.93	60.19
1992	5.47	0.39	3.50	22.58
1993	3.66	0.29	2.90	19.32
1994	4.28	0.32	2.98	89.59
1995	4.63	0.31	3.25	95.49
1996	4.47	0.29	3.22	118.22
1997	4.18	0.30	3.42	172.33
1998	3.59	0.23	2.61	168.74
1999	4.19	0.27	3.23	183.15
2000	4.64	0.31	3.32	164.74
2001	3.48	0.27	2.70	130.60
2002	3.63	0.26	2.45	245.42
2003	3.46	0.24	2.76	372.94
2004	4.63	0.30	3.40	441.19
2005	5.24	0.29	3.33	398.14

## 4.9. REFERENCES

Central Bureau of Statistics, Department of Manufacturing and Mining, *Monthly Industrial Reports (1990 – 2005)*, Zagreb

Central Bureau of Statistics, *Statistical Yearbooks (1990-2006)*, Zagreb

IPCC (2000) *Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories*, Japan

IPCC/UNEP/OECD/IEA (1997) *Greenhouse Gas Inventory Workbook*, Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories, Volume 2, United Kingdom

IPCC/UNEP/OECD/IEA (1997) *Greenhouse Gas Inventory Reference Manual*, Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories, Volume 3, United Kingdom

Ministry of Environmental Protection, Physical Planning and Construction (2006) *National Inventory Report 2006, Croatian greenhouse gas inventory for the period 1990 – 2004*, EKONERG, Zagreb

Ministry of Environmental Protection, Physical Planning and Construction (2006) *Second, Third and Fourth National Communication of the Republic of Croatia under the United Nations Framework Convention on Climate Change (UNFCCC)*, Zagreb

## 5. SOLVENT AND OTHER PRODUCT USE (CRF sector 3)

### 5.1. SOLVENT AND OTHER PRODUCT USE

#### 5.1.1. SOURCE CATEGORY DESCRIPTION

The most significant emissions in this category are the emissions of non-methane volatile organic compounds (NMVOCs). The use of solvents is the cause of less than 15 percent of anthropogenic national emissions of NMVOC. The emissions of NMVOC is caused by use of solvent based paint and varnish, degreasing of metal and dry cleaning, in production of chemicals, in printing industry, by use of glue, by use of solvents in households and by all other activities where solvents are used. NMVOC emissions oxidize in the atmosphere and CO<sub>2</sub> emissions are generated as a consequence of this oxidation.

NMVOC and CO<sub>2</sub> emissions are included in emissions estimates in this sector. N<sub>2</sub>O emissions from medical uses and other possible sources are not estimated because activity data are not available.

#### 5.1.2. METHODOLOGICAL ISSUES

Estimation of NMVOC emissions from Solvent and Other Product Use (provided by *EMEP-CORINAIR Emission Inventory Guidebook*) has been carried out by estimating the amount of solvent containing products consumed. The NMVOC emissions from Solvent and Other Product Use have been calculated by using simpler methodology. Default emission factor (*EMEP-CORINAIR Emission Inventory Guidebook*) has been applied for each source category. For several source categories (degreasing and dry cleaning, pharmaceutical products manufacturing and domestic solvent use) the NMVOC emissions calculation is based on population data. The activity data for the other sources were extracted from Monthly Industrial Reports published by Central Bureau of Statistics, Department of Manufacturing and Mining. The contribution of group of activities to NMVOC emissions is given in the Figure 5.1-1.




Figure 5.1-1: Emissions of NMVOC from Solvent and Other Product Use (1990-2005)

Activity data, NMVOC emissions and average emission factors are shown in Table 5.1-1.

Table 5.1-1: NMVOC emissions from Solvent and Other Product Use (Gg) (1990-2005)

Source and Sink Categories		Activity Data								NMVOC Emission								Emission Factor
		1990	1991	1992	1993	1994	1995	1996	1997	1990	1991	1992	1993	1994	1995	1996	1997	1990-2005
		Gg (1000 capita*)								Gg								t/Gg (cap)
<b>3</b>	<b>Total – Solvent Use</b>									<b>27.376</b>	<b>29.984</b>	<b>23.960</b>	<b>25.621</b>	<b>27.516</b>	<b>27.358</b>	<b>29.900</b>	<b>25.577</b>	
<b>3A</b>	<b>Paint Application</b>									<b>3.184</b>	<b>2.011</b>	<b>1.376</b>	<b>1.314</b>	<b>1.566</b>	<b>1.562</b>	<b>2.020</b>	<b>2.175</b>	
	Use of Solvent Base Paint	6.367	4.023	2.753	2.629	3.131	3.124	4.041	4.351	3.184	2.011	1.376	1.314	1.566	1.562	2.020	2.175	500
<b>3B</b>	<b>Degreasing and Dry Cleaning</b>									<b>5.256</b>	<b>4.965</b>	<b>4.917</b>	<b>5.105</b>	<b>5.114</b>	<b>5.136</b>	<b>4.943</b>	<b>5.029</b>	
	Metal Degreasing *	4778	4514	4470	4641	4649	4669	4494	4572	4.061	3.837	3.800	3.945	3.952	3.969	3.820	3.886	0.85
	Dry Cleaning *	4778	4514	4470	4641	4649	4669	4494	4572	1.195	1.129	1.118	1.160	1.162	1.167	1.124	1.143	0.25
<b>3C</b>	<b>Chemical Products</b>									<b>5.830</b>	<b>5.546</b>	<b>4.302</b>	<b>3.275</b>	<b>4.697</b>	<b>5.140</b>	<b>3.488</b>	<b>2.561</b>	
	Polyurethane – rigid foam	0.147	0.081	0.016	0.021	0.035	0.029	0.022	0.044	0.002	0.001	0	0	0.001	0	0	0.001	15
	Polyurethane – soft foam	3.616	2.717	1.660	2.025	2.427	2.880	1.800	1.710	0.090	0.068	0.042	0.051	0.061	0.072	0.045	0.043	25
	Polyester Resins	6.047	4.159	3.523	2.570	2.546	2.225	3.367	7.022	0.242	0.166	0.141	0.103	0.102	0.089	0.135	0.281	40
	Polystyrene Foam	39.069	26.383	57.045	57.666	58.215	49.356	56.513	50.894	0.586	0.396	0.856	0.865	0.873	0.740	0.848	0.763	15
	Polyvinylchloride	104.60	69.357	70.969	44.259	78.331	93.352	44.565	23.094	4.184	4.184	2.839	1.770	3.133	3.734	1.783	0.924	40
	Rubber Processing	5.739	5.442	2.439	2.477	2.338	2.285	1.279	0.026	0.086	0.082	0.037	0.037	0.035	0.034	0.019	0.0004	15
	Pharmaceutical Products Manufacturing*	4778	4514	4470	4641	4649	4669	4494	4572	0.067	0.063	0.063	0.065	0.065	0.065	0.063	0.064	0.014
	Paint and Varnish Manufacturing	21.956	13.872	9.493	9.064	10.797	10.773	13.933	15.002	0.329	0.208	0.142	0.136	0.162	0.162	0.209	0.225	15
	Ink Manufacturing	4.672	3.626	1.343	0.985	1.416	1.367	1.420	1.430	0.140	0.109	0.040	0.030	0.042	0.041	0.043	0.043	30
	Glue Manufacturing	5.139	13.451	7.151	10.910	11.166	10.076	17.197	10.874	0.103	0.269	0.143	0.218	0.223	0.202	0.344	0.217	20
<b>3D</b>	<b>Other Use of Solvent</b>									<b>13.107</b>	<b>17.461</b>	<b>13.365</b>	<b>15.927</b>	<b>16.139</b>	<b>15.520</b>	<b>19.448</b>	<b>15.811</b>	
	Printing Industry	4.672	3.626	1.343	0.985	1.416	1.367	1.420	1.430	0.467	0.363	0.134	0.099	0.142	0.137	0.142	0.143	100
	Application of Glue	5.139	13.451	7.151	10.910	11.166	10.076	17.197	10.874	3.083	8.071	4.291	6.546	6.700	6.046	10.318	6.524	600
	Domestic Solvent Use*	4778	4514	4470	4641	4649	4669	4494	4572	9.556	9.028	8.940	9.282	9.298	9.338	8.988	9.144	2

\* - Activity Data is Number of Inhabitants in Croatia (1000 capita)

Table 5.1-1: NMVOC emissions from Solvent and Other Product Use (Gg) (1990-2005), cont.

Source and Sink Categories		Activity Data								NMVOC Emission								Emission Factor
		1998	1999	2000	2001	2002	2003	2004	2005	1998	1999	2000	2001	2002	2003	2004	2005	1990-2005
		Gg (1000 capita*)								Gg								t/Gg (cap)
<b>3</b>	<b>Total – Solvent Use</b>									<b>25.581</b>	<b>22.992</b>	<b>23.439</b>	<b>25.477</b>	<b>33.836</b>	<b>36.694</b>	<b>46.058</b>	<b>52.918</b>	
<b>3A</b>	<b>Paint Application</b>									<b>2.244</b>	<b>2.203</b>	<b>2.191</b>	<b>2.435</b>	<b>2.200</b>	<b>2.223</b>	<b>2.299</b>	<b>2.377</b>	
	Use of Solvent Base Paint	4.487	4.406	4.381	4.870	4.400	4.446	4.599	4.754	2.244	2.203	2.191	2.435	2.200	2.223	2.299	2.377	500
<b>3B</b>	<b>Degreasing and Dry Cleaning</b>									<b>4.951</b>	<b>5.009</b>	<b>4.819</b>	<b>4.881</b>	<b>4.887</b>	<b>4.886</b>	<b>4.883</b>	<b>4.886</b>	
	Metal Degreasing *	4501	4554	4381	4437	4443	4442	4439	4442	3.826	3.871	3.724	3.771	3.777	3.776	3.773	3.776	0.85
	Dry Cleaning *	4501	4554	4381	4437	4443	4442	4439	4442	1.125	1.139	1.095	1.109	1.111	1.111	1.110	1.111	0.25
<b>3C</b>	<b>Chemical Products</b>									<b>3.050</b>	<b>1.669</b>	<b>1.363</b>	<b>1.774</b>	<b>2.265</b>	<b>2.098</b>	<b>2.259</b>	<b>2.761</b>	
	Polyurethane – rigid foam	0.039	0.060	0.060	0.095	0.180	0.070	0.060	0.120	0.001	0.001	0.001	0.001	0.003	0.001	0.001	0.002	15
	Polyurethane – soft foam	1.790	1.770	1.800	2.655	5.431	2.855	2.424	2.799	0.045	0.044	0.045	0.066	0.136	0.071	0.061	0.070	25
	Polyester Resins	8.258	5.609	12.848	9.661	14.693	9.704	10.948	10.886	0.330	0.224	0.514	0.386	0.588	0.388	0.438	0.435	40
	Polystyrene Foam	54.240	53.047	16.518	47.146	45.439	46.361	34.311	52.933	0.814	0.796	0.248	0.707	0.682	0.695	0.515	0.794	15
	Polyvinylchloride	33.134	3.085	0.811	0.640	0.617	0.206	0.055	0	1.325	0.123	0.032	0.026	0.025	0.008	0.002	0	40
	Rubber Processing	0.017	0.020	0.021	0.021	0.015	0.006	0.011	0.004	0.0003	0.0003	0.003	0.0003	0.0002	0.0001	0.0002	0.0001	15
	Pharmaceutical Products Manufacturing*	4501	4554	4381	4437	4443	4442	4439	4442	0.063	0.064	0.061	0.062	0.062	0.062	0.062	0.062	0.014
	Paint and Varnish Manufacturing	15.473	15.194	15.107	16.794	15.174	15.332	15.857	16.393	0.232	0.228	0.227	0.252	0.228	0.230	0.238	0.246	15
	Ink Manufacturing	1.071	0.797	0.916	0.822	0.863	0.789	0.673	0.665	0.032	0.024	0.027	0.025	0.026	0.024	0.020	0.020	30
	Glue Manufacturing	10.379	8.206	10.355	12.385	25.851	30.873	46.119	56.577	0.208	0.164	0.207	0.248	0.517	0.617	0.922	1.131	20
<b>3D</b>	<b>Other Use of Solvent</b>									<b>15.337</b>	<b>14.111</b>	<b>15.067</b>	<b>16.387</b>	<b>24.483</b>	<b>27.487</b>	<b>36.617</b>	<b>42.894</b>	
	Printing Industry	1.071	0.797	0.916	0.822	0.863	0.789	0.673	0.665	0.107	0.080	0.092	0.082	0.086	0.079	0.067	0.067	100
	Application of Glue	10.379	8.206	10.355	12.385	25.851	30.873	46.119	56.573	6.227	4.924	6.213	7.431	15.511	18.524	27.671	33.944	600
	Domestic Solvent Use*	4501	4554	4381	4437	4443	4442	4439	4442	9.002	9.108	8.762	8.874	8.886	8.884	8.878	8.884	2

\* - Activity Data is Number of Inhabitants in Croatia (1000 capita)



IPCC Guidelines do not provide methodology for calculation of CO<sub>2</sub> emissions from Solvent and Other Product Use. CO<sub>2</sub> emissions are calculated using conversion factor which contains ratio C/NMVOC = 0.8 and recalculation ratio of C to CO<sub>2</sub> equal to 44/12. The overall conversion factor has value of 2.93.

The resulting emissions of CO<sub>2</sub> from Solvent and Other Product Use in the period 1990-2005 are presented in the Figure 5.1-2.




Figure 5.1-2: Emissions of CO<sub>2</sub> from Solvent and Other Product Use (1990-2005)

### 5.1.3. UNCERTAINTIES AND TIME-SERIES CONSISTENCY

Uncertainties in CO<sub>2</sub> emissions estimates are mainly due to the accuracy of used conversion factor (C/NMVOC) and reliability of calculation is very low.

Uncertainty estimates are based on expert judgement. Uncertainty estimate associated with emission factors amounts 50 percent. Uncertainty estimate associated with activity data amounts 50 percent.

Emissions from Solvent and Other Product Use have been calculated using the same method and data sets for every year in the time series.

### 5.1.4. SOURCE-SPECIFIC QA/QC AND VERIFICATION

During the preparation of the inventory submission activities related to quality control were mainly focused on completeness and consistency of emission estimates and on proper use of notation keys in the CRF tables. After preparation of final draft of this chapter an audit was carried out to check selected activities from Tier 1 General inventory level QC and Tier 2 source-specific QC procedures. Solvent and Other Product Use is key source category. Regarding to Tier 2 activities, emission factors and activity data were checked for key source categories.

#### **5.1.5. SOURCE-SPECIFIC RECALCULATIONS**

In the previous report, CO<sub>2</sub> emissions were not calculated because IPCC Guidelines do not provide methodology for calculation of CO<sub>2</sub> emissions from Solvent and Other Product Use. In this report, CO<sub>2</sub> emissions are calculated by means of conversion factor which has value of 2.93. Thereupon, CO<sub>2</sub> emissions are recalculated for the period 1990-2004.

#### **5.1.6. SOURCE-SPECIFIC PLANNED IMPROVEMENTS**

For the purpose of accurate emission calculations, Croatia plan to investigate source category degreasing and dry cleaning, pharmaceutical products manufacturing and domestic solvent use. The NMVOC emissions calculation in these categories is based on population data.

N<sub>2</sub>O emissions from medical uses and other possible sources are not estimated because input data are not available. According to requirement of Regulation on the Greenhouse Gases Emissions Monitoring in the Republic of Croatia (Official Gazette No. 1/07), which will go into force in 2008, each sources of CO<sub>2</sub> and N<sub>2</sub>O emissions from Solvent and Other Product Use should report required activity data for more accurate emissions estimation.

## 5.2. REFERENCES

Central Bureau of Statistics, Department of Manufacturing and Mining, *Monthly Industrial Reports (1990 – 2005)*, Zagreb

Central Bureau of Statistics, *Statistical Yearbooks (1990-2006)*, Zagreb

EMEP/CORINAIR (1996) *Atmospheric Emission Inventory Guidebook*, European Environmental Agency, Denmark

Ministry of Environmental Protection, Physical Planning and Construction (2006) *National Inventory Report 2006, Croatian greenhouse gas inventory for the period 1990 – 2004*, EKONERG, Zagreb

Ministry of Environmental Protection, Physical Planning and Construction (2006) *Second, Third and Fourth National Communication of the Republic of Croatia under the United Nations Framework Convention on Climate Change (UNFCCC)*, Zagreb

## 6. AGRICULTURE (CRF sector 4)

### 6.1. OVERVIEW OF SECTOR

The agricultural activities contribute directly to the emission of greenhouse gases through various processes. The following sources have been identified to make a more complete break down in the emission calculation:

- Livestock: enteric fermentation ( $\text{CH}_4$ ) and manure management ( $\text{CH}_4$ ,  $\text{N}_2\text{O}$ )
- Agricultural soils ( $\text{N}_2\text{O}$ )

The total emissions in 2005 produced by the agricultural activities were 3495 Gg  $\text{CO}_2$ -eq, which represents 11.5 percent of the emission of the total emission inventory. The methane ( $\text{CH}_4$ ) and nitrous oxide ( $\text{N}_2\text{O}$ ) are primary greenhouse gases discharged as a consequence of agricultural activities (Figure 6.1-1). Of all the ruminants, the dairy cattle are the largest source of methane ( $\text{CH}_4$ ) emission. The results of the agricultural soil management, manure management, and the agricultural engineering in cultivation of some crops are relatively high emissions of nitrous oxide ( $\text{N}_2\text{O}$ ). The emission generated by burning the agricultural residues was not included in calculation because its activity is prohibited by law in Republic of Croatia. There are no ecosystems in Republic of Croatia that could be considered natural savannas or rice fields; consequently, no greenhouse gas emissions therefore exist for this sub-category.




Figure 6.1-1: Agriculture GHG Sources (year 2005)

Tables 6.1-1 and 6.1-2 show the total emission from agriculture by gases and emission sources for the period 1990-2005. The emission in table 6.1-2 is given in the equivalents of  $\text{CO}_2$ .

Table 6.1-1: Emission of greenhouse gases from agriculture (Gg)

Gas/Source	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
<b>CH<sub>4</sub></b>	<b>74.82</b>	<b>71.78</b>	<b>63.62</b>	<b>55.55</b>	<b>50.69</b>	<b>48.00</b>	<b>45.32</b>	<b>44.49</b>	<b>43.59</b>	<b>44.48</b>	<b>43.02</b>
Enteric Fermentation	63.99	60.99	54.97	47.16	42.29	40.43	37.86	37.17	36.36	36.48	35.60
Manure management	10.83	10.79	8.66	8.39	8.39	7.57	7.46	7.32	7.23	8.00	7.42
<b>N<sub>2</sub>O</b>	<b>9.33</b>	<b>9.45</b>	<b>7.33</b>	<b>7.26</b>	<b>7.04</b>	<b>6.75</b>	<b>6.73</b>	<b>7.65</b>	<b>6.85</b>	<b>7.06</b>	<b>7.18</b>
Manure management	1.22	1.17	1.09	0.91	0.84	0.80	0.74	0.73	0.71	0.73	0.71
Agricultural soil	8.12	8.29	6.24	6.35	6.21	5.96	5.99	6.92	6.13	6.33	6.48

Table 6.1-1: Emission of greenhouse gases from agriculture (Gg) (cont.)

Gas/Source	2001	2002	2003	2004	2005
<b>CH<sub>4</sub></b>	<b>43.61</b>	<b>42.70</b>	<b>48.58</b>	<b>47.44</b>	<b>46.64</b>
Enteric Fermentation	36.14	35.11	40.38	38.84	39.16
Manure management	7.47	7.59	8.20	8.60	7.48
<b>N<sub>2</sub>O</b>	<b>7.49</b>	<b>7.68</b>	<b>7.42</b>	<b>8.27</b>	<b>8.11</b>
Manure management	0.71	0.70	0.73	0.76	0.76
Agricultural soil	6.77	6.98	6.68	7.51	7.35

Table 6.1-2: Emission of greenhouse gases from agriculture CO<sub>2</sub>-eq (Gg)

Gas/Source	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
<b>CH<sub>4</sub></b>	<b>1571.26</b>	<b>1507.46</b>	<b>1336.09</b>	<b>1166.52</b>	<b>1064.39</b>	<b>1007.99</b>	<b>951.75</b>	<b>934.30</b>	<b>915.31</b>	<b>933.99</b>	<b>903.42</b>
Enteric Fermentation	1343.85	1280.81	1154.33	990.30	888.17	849.05	795.06	780.53	763.54	766.03	747.50
Manure management	227.41	226.65	181.76	176.22	176.22	158.94	156.69	153.77	151.77	167.96	155.92
<b>N<sub>2</sub>O</b>	<b>2892.74</b>	<b>2930.99</b>	<b>2272.93</b>	<b>2251.19</b>	<b>2182.81</b>	<b>2093.29</b>	<b>2086.11</b>	<b>2371.16</b>	<b>2123.02</b>	<b>2188.95</b>	<b>2226.66</b>
Manure management	376.71	361.27	338.01	281.22	259.10	246.87	230.44	226.09	221.53	225.91	218.86
Agricultural soil	2516.03	2569.72	1934.93	1969.97	1923.71	1846.42	1855.67	2145.07	1901.49	1963.04	2007.80

Table 6.1-2: Emission of greenhouse gases from agriculture CO<sub>2</sub>-eq (Gg) (cont.)

Gas/Source	2001	2002	2003	2004	2005
<b>CH<sub>4</sub></b>	<b>915.77</b>	<b>896.69</b>	<b>1020.28</b>	<b>996.19</b>	<b>979.51</b>
Enteric Fermentation	758.97	737.37	847.98	815.64	822.42
Manure management	156.79	159.32	172.30	180.55	157.09
<b>N<sub>2</sub>O</b>	<b>2320.42</b>	<b>2379.78</b>	<b>2299.52</b>	<b>2564.35</b>	<b>2515.32</b>
Manure management	221.09	216.73	227.75	235.19	236.67
Agricultural soil	2099.33	2163.05	2071.76	2329.16	2278.65

Below there is a review of the greenhouse gas emission calculation according to previously stated sources.

## 6.2. CH<sub>4</sub> EMISSIONS FROM ENTERIC FERMENTATION (CRF 4.A.)

### 6.2.1. SOURCE CATEGORY DESCRIPTION

The methane is a direct product of animal metabolism generated during the digestion process. The greatest producers of methane are ruminants (cows, cattle, and sheep). The amount of methane produced and excreted depends on the animal digestive system and the amount and type of the animal feed. Figure 6.2-1 shows the emission of methane from enteric fermentation for the period from 1990-2005. The estimates in this inventory include only emissions in farm animals. Buffalo, camels, and lamas do not occur in the Republic of Croatia. Emissions from wild animals and semi domesticated game are not quantified and neither are emissions from humans or pet animals. CH<sub>4</sub> emission from enteric fermentation is a key source, both by level and trend. Dairy cattle is the single major source of emissions representing 56 percent of total CH<sub>4</sub> emissions from enteric fermentation, followed by non dairy cattle representing 28 percent of total CH<sub>4</sub> from enteric fermentation. Jointly, cattle are responsible for 84 percent of total CH<sub>4</sub> emission from enteric fermentation. No methodology for calculating CH<sub>4</sub> emission from poultry is available in IPCC guidelines.




Figure 6.2-1: CH<sub>4</sub> emission from Enteric fermentation (Gg)

### 6.2.2. METHODOLOGICAL ISSUES

The IPCC methodology has been used to calculate the methane emission from enteric fermentation. IPCC methodology provides two different methods for estimating the quantity of methane from enteric fermentation. Tier I (simplified method) has been used as well as default EF specific for the animal type, the climate zone (cool), geographic region (eastern Europe) and the degree of the region development (developing countries), (IPCC, 1996). Three year average livestock population data for all livestock types for 1990 year were obtained from Croatian Statistical Report (1988, 1989 and 1990). For period 1991-2003 data from FAO Statistics and Croatian statistical yearbooks were taken. For 2004 data were taken from Statistical Yearbook 2004, Agricultural report 2004 and Inventory of agriculture in Croatia 2004. For 2005 data were taken from FAO Statistics and Statistical Yearbook 2005. All sources are official. The numbers of animals 1990-2005 are reported in Figure 6.2-2 and Figure 6.2-3. Tier 2 method was not applied to dairy cattle because accurate activity data was not available.




Figure 6.2-2: Number of dairy cattle, cattle, swine sheep, goats and horses in thousands




Figure 6.2-3: Number of poultry in thousands

### 6.2.3. UNCERTAINTIES AND TIME-SERIES CONSISTENCY

Uncertainty estimates based on expert judgement. Uncertainty of activity data amounts 30 percent. Uncertainty of emission factors amounts 40 percent.

Data sources in the category differ because none of above mentioned sources contains data for all animal types.

### 6.2.4. SOURCE SPECIFIC RECALCULATIONS

There are no source-specific recalculations in sub-sector CH<sub>4</sub> Emissions from Enteric Fermentation.

### 6.3. MANURE MANAGEMENT – CH<sub>4</sub> EMISSIONS (CRF 4.B.)

#### 6.3.1. SOURCE CATEGORY DESCRIPTION

The management of livestock manure produces both methane (CH<sub>4</sub>) and nitrous oxide (N<sub>2</sub>O) emissions. The methane is generated under the conditions of anaerobic decomposition of manure. The storing methods of the manure in which the anaerobic conditions prevail (liquid animal manure in septic pits) are favourable for anaerobic decomposition of organic substance and release of methane. The storing of solid animal manure results in aerobic decomposition and very low production of methane. The methane emission from manure management for the period from 1990 to 2005 is given on the Figure 6.3-1.




Figure 6.3-1: CH<sub>4</sub> emission from Manure Management (Gg)

#### 6.3.2. METHODOLOGICAL ISSUES

The IPCC methodology (Tier 1) has been used to calculate the methane emission from enteric fermentation and manure management. The basic input is the head of cattle (dairy cattle, cattle, sheep, horses, pigs, poultry, goats, mules and asses). The emission factors specific for the animal type, the climate zone (cool), geographic region (Eastern Europe), and the degree of the region development (developing countries) were used for the calculation of the emission. Three year average livestock population data for all livestock types for 1990 year were obtained from Croatian Statistical Report (1988, 1989 and 1990). For period 1991-2003 data from FAO Statistics and Croatian statistical yearbooks were taken. For 2004 data were taken from Statistical Yearbook 2004, Agricultural report 2004 and Inventory of agriculture in Croatia 2004. For 2005 data were taken from FAO Statistics and Statistical Yearbook 2005. All sources are official. The emission factors have been taken from the *Revised 1996 IPCC Reference Manual*.

#### 6.3.3. UNCERTAINTIES AND TIME-SERIES CONSISTENCY

Uncertainty estimates are based on expert judgement. Uncertainty of activity data amounts 30 percent. Uncertainty of emission factors amounts 40 percent.

Data sources in the category differ because none of above mentioned sources contains data for all animal types.



### 6.3.4. SOURCE SPECIFIC RECALCULATIONS

There are no source-specific recalculations in sub-sector CH<sub>4</sub> Emissions from Manure Management.

## 6.4. N<sub>2</sub>O EMISSIONS FROM MANURE MANAGEMENT (CRF 4.B.)

### 6.4.1. SOURCE CATEGORY DESCRIPTION

The emissions of nitrous oxide (N<sub>2</sub>O) from all Animal Waste Management Systems are estimated here. A considerable amount of nitrous oxide evolves during storage of animal waste and is attributed to livestock breeding. This includes emissions from anaerobic lagoons, liquid systems, solid storage, dry lot, and other systems. The N<sub>2</sub>O emissions from pasture range and paddock are reported under Agricultural soils. Farm animals emit directly very little nitrous oxide and have not been considered in estimating GHG. In Republic of Croatia manure is not used for fuel.

### 6.4.2. METHODOLOGICAL ISSUES

The IPCC calculation methodology (Tier 1) has been used. The emission factors are taken from the *Revised 1996 IPCC Reference Manual*. The nitrous oxide (N<sub>2</sub>O) emission is calculated according to the following equations:

$$Nex_{(AWMS)} = \sum_{(T)} [N_{(T)} \times Nex_{(T)} \times AMWS_{(T)}]$$

Where:

$Nex_{(AWMS)}$  - N excretion per Animal Waste Management System

$N_{(T)}$  - numbers of animals of type

$Nex_{(T)}$  - N excretion of animals of type

$AMWS_{(T)}$  - fraction of  $Nex_{(T)}$  that is managed in one of the different distinguished animal waste management systems

T - type of animal category

and

$$N_2O_{(AWMS)} = \sum [Nex_{(AWMS)} \times EF_3]$$

where:

$N_2O_{(AWMS)}$  - N<sub>2</sub>O emissions from all Animal Waste Management Systems (kg N/yr)

$Nex_{(AWMS)}$  - N excretion per Animal Waste Management System (kg/yr)

$EF_3$  - emission factor

The nitrous oxide (N<sub>2</sub>O) emissions from manure management for the period from 1990 to 2005 are shown on figure 6.4-1.




Figure 6.4-1: N<sub>2</sub>O Emissions from Manure Management (Gg)

Three year average livestock population data for all livestock types for 1990 year were obtained from Croatian Statistical Report (1988, 1989 and 1990). For period 1991-2003 data from FAO Statistics and Croatian statistical yearbooks were taken. For 2004 data were taken from Statistical Yearbook 2004, Agricultural report 2004 and Inventory of agriculture in Croatia 2004. For 2005 data were taken from FAO Statistics and Statistical Yearbook 2005. All sources are official. The nitrogen excretion for each manure management system and the emission factors were taken from the *Revised 1996 IPCC Reference Manual* (Table 5.1.7).

#### 6.4.3. UNCERTAINTIES AND TIME-SERIES CONSISTENCY

Uncertainty estimates based on expert judgement. Uncertainty of activity data amounts 30 percent. Uncertainty of emission factors amounts 60 percent.

Data sources in the category differ because none of above mentioned sources contains data for all animal types.

#### 6.4.4. SOURCE SPECIFIC RECALCULATIONS

Data for goat and mule stocks were included in calculations of N<sub>2</sub>O Emissions from Manure Management according to Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories. Hence, N<sub>2</sub>O Emissions from Manure Management were recalculated for the period from 1990 to 2005.

## 6.5. AGRICULTURAL SOILS (CRF 4.D.)

A number of agricultural activities add nitrogen to soils, thereby increasing the amount of nitrogen available for nitrification and denitrification, and ultimately the amount of N<sub>2</sub>O emitted. Three sources of nitrous oxide emissions are distinguished in methodology:

- Direct emissions of N<sub>2</sub>O from agricultural soils
- Direct soil emissions of N<sub>2</sub>O from animal production
- Indirect emissions of N<sub>2</sub>O conditioned by agricultural activities

The highest among the above stated emission comes directly from the agricultural soils by cultivation of soil and crops. The activities stated include the use of synthetic and organic fertilizers, growing of leguminous plants and soybean (nitrogen fixation), nitrogen from the agricultural residues, and the treatment of histosols.




Figure 6.5-1: Total N<sub>2</sub>O Emissions from Agricultural Soils (Gg)

### 6.5.1. DIRECT EMISSION FROM AGRICULTURAL SOILS

#### 6.5.1.1. Source category description

Direct emissions of N<sub>2</sub>O from agricultural soils includes total amount of nitrogen applied to soils through cropping practices. These practices includes application of synthetic fertilizer, nitrogen from animal waste, production of nitrogen – fixing crops, nitrogen from crop residue mineralization and soil nitrogen mineralization due to cultivation of histosols. The input data required for this part of the calculation are: annual quantity of the synthetic fertilizer used, the quantity of organic fertilizer, the head of animals by its category, the biomass of leguminous plants and soybean, and the surface of histosols. The direct emission from agricultural soils is calculated by the following equation:

$$N_2O_{\text{DIRECT}} \text{ (kg N/yr)} = (F_{\text{SN}} + F_{\text{AW}} + F_{\text{CR}} + F_{\text{BN}}) \times EF_1 + F_{\text{OS}} \times EF_2$$

where:

N<sub>2</sub>O<sub>DIRECT</sub> - direct N<sub>2</sub>O emission from agricultural soils (kg N/yr)

F<sub>SN</sub> - nitrogen from synthetic fertilizer excluding emissions of NH<sub>3</sub> and NO<sub>x</sub> (kg N/yr)

- $F_{AW}$  - nitrogen from animal waste (kg N/yr)  
 $F_{CR}$  - nitrogen from crop residues (kg N/yr)  
 $F_{BN}$  - nitrogen from N-fixing crops (kg N/yr)  
 $EF_1, EF_2$  - emission factors  
 $F_{OS}$  - nitrogen from histosols, (kg N/yr)

Figure 6.5-2 shows direct emission of nitrous oxide from agricultural soils (1990-2005).




Figure 6.5-2: Direct  $N_2O$  Emissions from Agricultural Soils (Gg)

#### 6.5.1.2. Methodological issues

For the emission from agricultural soils the IPCC methodology (Tier 1) has been used. The emission factors have been taken from the *Revised 1996 IPCC Reference Manual*.

#### Nitrous oxide from mineral fertilisers

This estimate is based on the amount of N in mineral fertiliser that is annually consumed in the Republic of Croatia. The consumption of nitrogen from mineral fertilisers that are annually consumed in Croatia is obtained from Fertiliser Company Kutina, Statistical Yearbook and expert judgment. Of the total estimated quantity of emitted nitrogen N, which is dispersed into atmosphere in the form of ammonia and  $NO_x$  (10 percent; IPCC, 1996) was subtracted. The emission of nitrous oxide was then calculated by multiplying the quantity of the remaining N with emission factor 0.0125 kg  $N_2O$ -N/kg N (IPCC, 1996).

#### Nitrous oxide from animal manure and liquid/slurry

The estimate is based on the amount of N in solid manure and liquid manure/slurry, which is annually used for fertilizing crops. Of the total estimated quantity of emitted N, the N that is emitted on the pasture (24 percent; country specific), and N that is dispersed into the atmosphere in the form of ammonia and  $NO_x$  (20 percent; IPCC, 1996) was subtracted. The emission of nitrous oxide was then calculated by multiplying the quantity of the remaining N with emission factor 0.0125 kg  $N_2O$ -N/kg N (IPCC, 1996).

### **Nitrous oxide from biological fixation of N**

The estimate is based on the amount of pulses and soybeans (Statistical Yearbook) produced in country as a dry biomass. According to IPCC 1996, dry biomass of N-fixing crops was multiplied by factor 2 and by fraction of nitrogen in N-fixing crops (3 percent; IPCC, 1996). The emission of nitrous oxide was then calculated by multiplying the quantity of the remaining N with emission factor 0.0125 kg N<sub>2</sub>O-N/kg N (IPCC, 1996).

### **Emissions of nitrous oxide from crop residue**

The estimate is based on IPCC (1996) methodology which is based on the assumption that the total biomass produced is approximately twice the amount of the produced edible parts of crops, which means that 45 percent of the produced total biomass is removed from agricultural soils. Dry biomass production of pulses and soybeans and dry biomass production of other crops are the basic data for the calculation (Statistical Yearbook and expert judgment). Dry biomass is calculated by applying factor to account for crop water content. Fraction of N-fixing crops (3 percent), fraction of nitrogen in non N-fixing crops (1.5 percent), fraction of crop residue that is removed from the field as crop (45 percent) and fraction of crop residue that is burned rather than left on field (10 percent) were obtained from IPCC (1996). The emission of nitrous oxide was then calculated by multiplying the quantity of the remaining N with emission factor 0.0125 kg N<sub>2</sub>O-N/kg N (IPCC, 1996).

### **Emissions of nitrous oxide due to cultivation of organic soils**

Cultivation of soils with high contents of organic materials causes a release of a long term bound N. The area of organic soil in Republic of Croatia has been obtained from expert judgment. The emission of nitrous oxide due to cultivation of histosols was then calculated by multiplying the area of histosols with emission factor 5 kg N/ha/yr (IPCC, 1996).

#### **6.5.1.3. Uncertainties and time-series consistency**

Uncertainty estimates based on expert judgement. Uncertainty of activity data amounts 30 percent. Uncertainty of emission factors amounts 40 percent.

Direct N<sub>2</sub>O emissions from agricultural soils have been calculated using the same data sets for every year in the time series except for the calculations of N<sub>2</sub>O emissions from animal manure and liquid/slurry because none of the sources contains livestock data for all animal types.

#### **6.5.1.4. Source specific recalculations**

In the previous report insufficient data was estimated as average value for 2001, 2002 and 2003. In this report adjustment method "Extrapolation of emissions based on a driver", proposed by the *Technical Guidance on Methodologies for Adjustments under Article 5, Paragraph 2, of the Kyoto Protocol*, has been used for calculation of insufficient data. The pattern over the period 1992-2003 has been used for the estimation of pulses and soybeans and non-N-fixing crop production for 2004 and 2005.

## 6.5.2. DIRECT EMISSION OF N<sub>2</sub>O FROM ANIMALS

### 6.5.2.1. Methodological issues

Estimates of N<sub>2</sub>O emissions from animals were based on animal waste deposited directly on soils by animals in pasture, range and paddock. N<sub>2</sub>O emissions from animals can be calculated as follows:

$$N_2O_{ANIMALS} = N_2O_{(AWMS)} = \sum_{(T)} [N_{(T)} \times Nex_{(T)} \times AWMS_{(T)} \times EF_{3(AWMS)}]$$

where:

- $N_2O_{ANIMALS}$  - N<sub>2</sub>O emissions from animal production (kg N/yr)
- $N_2O_{(AWMS)}$  - N<sub>2</sub>O emissions from Animal Waste Management Systems (kg N/yr)
- $N_{(T)}$  - number of animals of type T
- $Nex_{(T)}$  - N excretion of animals of type T (kg N/animal/yr)
- $AWMS_{(T)}$  - fraction of  $Nex_{(T)}$  that is managed in one of the different distinguished animal waste management systems for animals of type T
- $EF_{3(AWMS)}$  - emission factor

The same emission factor (0.02 kg N<sub>2</sub>O-N/kg of emitted N) suggested by IPCC (1996) was used for all grazing animals, regardless of their species and climatic conditions.

Figure 6.5-3 shows direct emission of nitrous oxide from animals (1990-2005).




Figure 6.5-3: Direct N<sub>2</sub>O Emissions from Animals (Gg)

### 6.5.2.2. Uncertainties and time-series consistency

Uncertainty estimates based on expert judgement. Uncertainty of activity data amounts 30 percent. Uncertainty of emission factors amounts 40 percent.

Data sources in the category differ because none of the sources contains livestock data for all animal types.

### 6.5.2.3. Source specific recalculations

Data for goat and mule stocks were included in calculations of direct N<sub>2</sub>O emissions from animals according to *Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories*. Hence, direct N<sub>2</sub>O emissions from animals were recalculated for the period from 1990 to 2005.

## 6.5.3. INDIRECT N<sub>2</sub>O EMISSIONS FROM NITROGEN USED IN AGRICULTURE

### 6.5.3.1. Source category description

Calculations of indirect N<sub>2</sub>O emissions from nitrogen used in agriculture are based on two pathways. These are: volatilization and subsequent atmospheric deposition of NH<sub>3</sub> and NO<sub>x</sub> (originating from the application of fertilizers and animal manure), and leaching and runoff of the N that is applied to, or deposited on soils. These two indirect emission pathways are treated separately, although the activity data used are identical. The indirect emission of N<sub>2</sub>O from the agriculture is calculated by the following equation:

$$N_2O_{\text{INDIRECT}} = N_2O_{(G)} + N_2O_{(L)}$$

Where:

- N<sub>2</sub>O<sub>INDIRECT</sub> - indirect N<sub>2</sub>O emissions (kg N/yr)
- N<sub>2</sub>O<sub>(G)</sub> - N<sub>2</sub>O emissions due to atmospheric deposition of NH<sub>3</sub> and NO<sub>x</sub> (kg N/yr)
- N<sub>2</sub>O<sub>(L)</sub> - N<sub>2</sub>O emissions due to nitrogen leaching and runoff (kg N/yr)

The emissions of N<sub>2</sub>O produced from the discharge of human sewage N into rivers are reported under the sector Waste.

Figure 6.5-4 shows the indirect emission of nitrous oxide from agriculture (1990-2005).




Figure 6.5-4: Indirect N<sub>2</sub>O Emissions from Agricultural (Gg)

### 6.5.3.2. Methodological issues

#### **Nitrous oxide arising due to volatilization of ammonia (NH<sub>3</sub>) and nitrogen oxides (NO<sub>x</sub>)**

In fertilizing agricultural soils with nitrogen fertilizers, some N volatilises in form of ammonia and nitrogen oxides (N<sub>2</sub>O). This nitrogen is deposited by precipitation and particulate matter on agricultural soil, in forests and waters and thus indirectly contributes to emissions of N<sub>2</sub>O. Emissions are attributed to the place of origin of ammonia and NO<sub>x</sub>, not to the place where N is re-deposited, causing N<sub>2</sub>O emissions.

#### **Emissions from mineral fertilizers**

Indirect emissions of nitrous oxide from mineral fertilizers depend to a large extent on the fraction of N that volatilises during fertilizing. The amount of volatilised N depends very strongly on the type of fertilizer as well as on weather conditions and the manner of application. It has been considered that 10 percent of N from mineral fertilizers volatilises (IPCC, 1996). For calculating indirect emissions of nitrous oxide, the emission factor 0.01 kg N<sub>2</sub>O-N/kg NH<sub>3</sub> and NO<sub>x</sub>-N (IPCC, 1996) has been considered.

#### **Emissions from animal manure**

Numerous factors influence the fraction of volatilised N in form of ammonia and nitrogen oxides, such as: the ratio between N excreted in dung and N excreted in urine, the manner of slurry storage, the manner of slurry application etc. Generic IPCC (1996) emission factor (20 percent) of the excreted N is supposed to volatilise in form of ammonia and nitrogen oxides. Emissions of nitrous oxide have been calculated by multiplying the estimated quantities of volatilised N with emission factor 0.01 kg N<sub>2</sub>O-N/kg NH<sub>3</sub>-N and NO<sub>x</sub>-N (IPCC, 1996).

#### **Nitrous oxide from leaching and runoff of nitrogen compounds into surface waters, groundwater, and watercourses**

Surface runoff and leaching of N into groundwater, surface waters, and watercourses due to mineral fertilisers:

It has been considered that 30 percent of N from mineral fertilizers is lost through surface runoff and leaching into the groundwater and watercourses. In calculating emissions of nitrous oxide, it has been considered that for every kg of leached/run-off nitrogen, 0.025 kg of N<sub>2</sub>O-N is emitted (IPCC, 1996).

Nitrogen leaching and runoff into groundwater, surface waters, and watercourses due to animal manure:

It has been considered that for every kg of N, which is excreted by farm animals, 0.3 kg of N is lost through surface runoff to watercourses and groundwater (IPCC, 1996). In calculating emissions of nitrous oxide, the same emission factors has been considered as in the case of nitrogen leaching/runoff due to mineral fertilizer (0.025 kg N<sub>2</sub>O-N/kg of leached/run-off N)

### 6.5.3.3. Uncertainty and time-series consistency

The uncertainty of the calculation is conditioned by the use of the emission factors recommended by the methodology and the unreliability of the input data. According to the



bibliography, the uncertainty of the recommended emission factors is high. Uncertainty of activity data amounts 30 percent. Uncertainty of emission factors amounts 60 percent. Data sources for mineral fertilizer use were the same for every year in the time series while livestock data were obtained from different data sources because none of the sources contains data for all animal types.

#### **6.5.3.4. Source specific recalculations**

Data for goat and mule stocks were included in calculations of indirect N<sub>2</sub>O emissions from nitrogen used in agriculture according to recommendation by *Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories*. Hence, indirect N<sub>2</sub>O emissions were recalculated for the period from 1990 to 2005.

### **6.6. SOURCE SPECIFIC QA/QC AND VERIFICATION**

During the preparation of the inventory submission activities related to quality control were mainly focused on completeness and consistency of emission estimates and on proper use of notation keys in the CRF tables. After preparation of final draft of this chapter an audit was carried out to check selected activities from Tier 1 General inventory level QC procedures which revealed that most of the activities were correctly carried out during preparation of the inventory despite the fact that formal QC procedures were not prepared.

Regarding to Tier 2 activities, emission factors and activity data were checked for key source categories. In Agriculture four source categories represent key source category: enteric fermentation in domestic livestock, manure management, direct emission from agricultural soils and indirect emissions from nitrogen use in agriculture.

### **6.7. SOURCE SPECIFIC PLANNED IMPROVEMENT**

Emission of CH<sub>4</sub> from enteric fermentation in domestic livestock were estimated using Tier 1 method since detailed data on livestock population is not available at the moment and a comprehensive research is required in the future to provide these data. The availability and consistency of activity data is still a major problem in other key source categories within this sector and application of higher tier methodologies will be possible in the future after detailed research and adjustments of statistical methods for data collection. Moreover, national emission factors should be developed to increase the calculation quality.

### **6.8. REFERENCES**

Bašić, F., O. Nestroy, I. Kisić, A. Butorac, M. Mesić (1988) *Zaštita od erozije-ključna uloga obrade tla u aktualnim i izmijenjenim klimatskim prilikama*, HAZU – Znanstveni skup s međunarodnim sudjelovanjem "Prilagodba poljoprivrede i šumarstva klimi i njenim promjenama", str 115-125, Zagreb

- Bogunović, M., Vidaček, Ž., Racz, Z., Husnjak, S., Sraka, M. (1997) *Namjenska pedološka karta Republike Hrvatske i njena uporaba*, Agronomski glasnik 5-6, str. 369-399, Zagreb
- Butorac, A. (1963) *Toplinsko stanje tla i najkarakterističniji pragovi temperature tla za neka mjesta u glavnim poljoprivrednim rajonima Hrvatske*, Agr. glasnik 12/63
- Central Bureau of Statistics, *Statistical Yearbooks (1999-2005)*, Zagreb
- Central Bureau of Statistics (2003) *Report on agricultural production*, Zagreb
- Central Bureau of Statistics (2003) *Inventory of agriculture in the Republic of Croatia*, Zagreb
- IPCC (2000) *Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories*, Japan
- IPCC/UNEP/OECD/IEA (1997) *Greenhouse Gas Inventory Workbook*, Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories, Volume 2, United Kingdom
- IPCC/UNEP/OECD/IEA (1997) *Greenhouse Gas Inventory Reference Manual*, Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories, Volume 3, United Kingdom
- FAO (1976) *A framework for land evaluation*, Soil Bull. No.32. FAO, Rome and ILRI, Wageningen, Publ. No.22.
- FAO DATA BASE [www.fao.org](http://www.fao.org)
- Kisić, I., Mesić, F. Bašić, A. Butorac (1999) *Kalcifikacija i gnojidba tala gorske Hrvatske (Projekt financiran od strane MPŠ RH Vijeća za istraživanje u poljoprivredi – Razvoj službi za potporu obiteljskim poljoprivrednim gospodarstvima)*, Fond str. dok. Zavoda za OPB Agronomskog fakulteta u Zagrebu, str. 16., Zagreb
- Mesić, M., A. Butorac, F. Bašić, I. Kisić, I. Gašpar (1999) *Utjecaj različitih količina mineralnog dušika primijenjenih za uljnu repicu na količinu dušika u vodi iz drenskih cijevi*, XXXV. Znanstveni skup hrvatskih agronoma, str. 53-54, Opatija
- Penzar, B., Makjanić, B. (1980) *Osnovna statistička obrada podataka u klimatologiji*, Sveučilište u Zagrebu, 159 str., Zagreb
- Ministry of Environmental Protection, Physical Planning and Construction (2006) *National Inventory Report 2006, Croatian greenhouse gas inventory for the period 1990 – 2004*, EKONERG, Zagreb
- Ministry of Environmental Protection, Physical Planning and Construction (2006) *Second, Third and Fourth National Communication of the Republic of Croatia under the United Nations Framework Convention on Climate Change (UNFCCC)*, Zagreb
- Racz, Z. (1992) *Svjetski i domaći trendovi zaštite tala i poljoprivredne proizvodnje od Stocholma do Rio de Jeneira*, Socijalna ekologija, Vol. 1, No 3, 399-405, Zagreb
- Racz, Z. (1992) *Značaj tla u prirodnim i agroekosustavima*, Socijalna ekologija Vol. 1, No 1, 105-118, Zagreb
- Znaor, D. and Karoglan Todorović, S. (2005) *Analysis of yields, nitrogen content and N-fixation of Croatian agricultural production in 2002-2003*, University of Essex, Colchester, Unpublished

## 7. LAND-USE, LAND USE CHANGE AND FORESTRY (CRF sector 5)

### 7.1. OVERVIEW OF SECTOR

Forests and woodland in the Republic of Croatia are goods of a general interest and are under special protection of the state. Forest is an area of land of minimum 0.1 ha covered with trees (*Forestry Act, Official Gazette No. 140/05, 82/06*). The terms and the way of their use have been prescribed in Forestry Act. Ministry of Agriculture, Forestry and Water Management is authorized institution for collecting data about state of forest land. Moreover, Regulation on the Greenhouse Gases Emissions Monitoring in the Republic of Croatia (Official Gazette No. 1/07), which will come into force in 2008, prescribes obligation and procedure for emissions monitoring. Among others, the regulation prescribes monitoring of areas within different land use categories, such as forest area, agricultural area, grasslands, wetlands, settlements and other land. Based on the Forest Management Area Plan of the Republic of Croatia (1996-2005), the forests and the forest land cover 43.5 percent of the total surface area. By its origin, approximately 95 percent of the forests in Croatia were formed by natural regeneration and the 5 percent of the forests are grown artificially. In the Republic of Croatia 81 percent of the forest are state owned and 19 percent are private. Out of the total surface area occupied by forests 2089607 ha (84 percent) is the forest-covered area, 327630 ha (13 percent) is non forest land, and 74063 ha (3 percent) is bare unproductive and unfertile forestland. The basic principles of the Croatian forestry are sustainable forest management along with the preservation of the natural structure and diversity of the forests, as well as the permanent enhancement of the stability and quality of the forests commercial and welfare functions.

The total growing stock in the Croatian forests is around 342 million m<sup>3</sup>. It consists of approximately 84 percent of deciduous trees and 16 percent of evergreen trees. The most frequent species are beech (*Fagus sylvatica*), common fir (*Abies alba*), sessile oak (*Quercus petraea*), and other types of deciduous and evergreen trees. The average growing stock in the state-owned forests is 202 m<sup>3</sup>/ha and in the privately owned forests 82 m<sup>3</sup>/ha. The annual increment in Croatia forests is around 9643000 m<sup>3</sup> of wood. The increment is an increase in the forest timber stock over a specific period and it is calculated as an annual, periodical and average increment. The check method or the method of bore-spills is most often used in Croatia to identify the increment. The quality and quantity of increment can be improved by different methods of forest cultivation. The annual cut is a part of the forest timber stock planned for commercial harvesting for a certain period (1 year, 10 years, 20 years) expressed in timber stock (m<sup>3</sup>, m<sup>3</sup>/ha) or by the surface area. To satisfy the basic principles of the sustainable forest management (continuous management), the annual cut must not be larger than the increment value.

According to the methodology proposed by *IPCC Good Practice Guidance for LULUCF* (GPG 2003) the top-level categories for greenhouse gas (GHG) reporting are:

- Forest land
- Cropland
- Grassland
- Wetlands
- Settlements
- Other land

The Republic of Croatia only reports data for Forest land category. Data needed for calculations of emissions/removals for other land categories are partly available but not enough adequate, consistent and complete. Working group for calculations emissions/removals from LULUCF sector was established by Ministry of Agriculture, Forestry and Water Management in the May 2005, with the objective to improve the quality of inventory and data collection. During the year 2005, Working group worked on harmonisation of national definition referring to GPG 2003 definition and internationally accepted definition to minimise gaps and overlaps.

## 7.2 SOURCE CATEGORY

### 7.2.1. SOURCE CATEGORY DESCRIPTION

The carbon in forests is bound in trees, underbrush, soil and dead wood. As a result of biological processes in forests and anthropogenic activities the carbon is in a constant cycling process. Deforestation, among all anthropogenic activities, has the greatest impact on the change of carbon stock in the existing forests. The problem of deforestation in Croatia does not exist. According to the current data total forest area in Croatia has not decreased over the last 100 years.

### 7.2.2. METHODOLOGICAL ISSUES

The IPCC methodology (GPG 2003) has been used for calculation of CO<sub>2</sub> emissions and removals from LULUCF sector. GHG inventory for the land-use category Forest Land Remaining Forest Land (FF) is reported using Tier 1 method. All emission factors were used according to GPG 2003. The Forest Management Area Plan of the Republic of Croatia for the period from 1986 to 1996 and from 1996 to 2005 is the main source for the data on the forest land and the annual increment (Table 7.2-1.). The data on commercial harvesting and wood fuel are obtained from Statistical yearbooks of the Republic of Croatia (1986-1989 and 1991-1996). Data on commercial harvest including wood for fuel for the year 1990 and years 1997-2005 are obtained from experts preparing data for UNECE. The criteria in choosing data were the following: continuity, quality, comparability as well as accessibility of sources. Contemporarily National Forestry Inventory Project (CRONFI) is ongoing and will be completed in year 2008. The law prohibits the renewal of forests by clear cutting, and the natural rejuvenation is the principal method for renewal of all natural forests.

*Table 7.2-1: Forest areas and annual increment of forests in Croatia according to Forest Management Plan (1986-1996; 1996-2006)*

Year	Forest area managed (ha)	Annual volume increment (m <sup>3</sup> )
1986 - 1996	2061509	9643000
1996 - 2006	2089606	9643000

#### 7.2.2.1. Forest Land Remaining Forest Land

According to GPG 2003, Tier 1 method, GHG inventory for the Forest land remaining forest land (FF) is estimated only for aboveground and belowground biomass. Other carbon pools are not taken in consideration since Tier I method is applied and due to lack of activity data.

Change in carbon stocks in living biomass is calculated by multiplying difference in oven dry weight of biomass increments and losses with appropriate carbon fraction. Method 1 (default method) is applied for estimating carbon stock changes in biomass. Method 1 required the biomass carbon loss ( $\Delta C_{FFL}$ ) to be subtracted from the biomass carbon increment ( $\Delta C_{FFG}$ ) for the reporting year (GPG 2003 Equation 3.2.2.)

Annual Increase in Carbon Stock due to biomass increment ( $\Delta C_{FFG}$ ) in FF is estimated according to Equation 3.2.4., GPG 2003. Estimation of annual increase in carbon stock due to

biomass increment required estimates of area and annual increment of total biomass for each forest type (coniferous, deciduous) ( $G_{TOTAL}$ ) and climatic zone (temperate) in Croatia. The carbon fraction of biomass (CF) used is default value of 0.5.

$G_{TOTAL}$  is the expansion of annual increment rate of aboveground biomass (Gw) to include belowground part involving multiplication by the ratio of belowground biomass to aboveground biomass (root to shoot ratio) that applies to increments. Since Gw data are not available directly the increment in volume (lv) was used with biomass expansion factor for conversion of annual net increment to aboveground increment.

Average annual increment in biomass (Gw) is calculated according to Equation 3.2.5. GPG 2003, using data on:

- lv = average annual net increment in volume suitable for industrial processing,  $m^3 ha^{-1} yr^{-1}$  (Forestry Management Plan 1986-2006)
- D = basic wood density, tonnes d.m.  $m^{-3}$ , (GPG 2003)
- BEF1 = biomass expansion factor for conversion of annual net increment (including bark) to aboveground tree biomass, dimensionless (GPG 2003)
- R = root to shoot ratio, dimensionless; GPG 2003, Table 3A.1.8
- CF = carbon fraction of dry matter (default = 0.5), tonnes C (tonne d. m. )  $^{-1}$

Average Increment in Biomass ( $G_{TOTAL}$ ) is calculated multiplying average increment in biomass (Gw) per root to shoot ratio (R) appropriate to increment, dimensionless (GPG 2003, Table 3A.1.8.)

Annual Decrease in Carbon Stock Due to Biomass Loss in FF ( $\Delta C_{FFL}$ ) is calculated as a sum of losses from commercial roundwood feelings ( $L_{fellings}$ ) and fuelwood gathering ( $L_{fuelwood}$ ) (GPG 2003, Equation 3.2.6). Other losses are not included due to lack of accurate data available.

Annual Carbon Loss due to Commercial Feelings ( $L_{fellings}$ ) is calculated according to Equation 3.2.7, GPG 2003., using input data on:

- H = annual extracted volume, roundwood,  $m^3 yr^{-1}$  (Statistical Yearbook, 1986-1996, UNECE 1996-2005)
- D = basic wood density, tonnes d.m.  $m^{-3}$ , (GPG 2003)
- BEF2 = biomass expansion factor for conversion volumes of extracted roundwood to total aboveground (including bark) biomass, dimensionless (GPG 2003)
- $f_{BL}$  = fraction of biomass left to decay in forest
- CF = carbon fraction of dry matter (default = 0.5), tonnes C (tonne d. m. )  $^{-1}$

In applying above mentioned equation,  $f_{bl}$  is set to 0 according to assumption that total biomass associated with volume of the extracted roundwood is considered as an immediate emission.

Annual Carbon Loss due to Fuelwood gathering is estimated according to Equation 3.2.8., GPG 2003 using input data on:

- FG = annual volume of fuelwood gathering, tonnes C  $yr^{-1}$  (Statistical Yearbook, 1986-1996, UNECE 1996-2005)
- D = basic wood density, tonnes d.m.  $m^{-3}$ , (GPG 2003)
- BEF2 = biomass expansion factor for conversion volumes of extracted roundwood to total aboveground (including bark) biomass, dimensionless (GPG 2003)
- CF = carbon fraction of dry matter (default = 0.5), tonnes C (tonne d. m. )  $^{-1}$

Other carbon losses that include losses from disturbances such as pest outbreaks, fires are not estimated due to lack of accurate data availability and are planned to be in future reports. More reliable data will be available upon completion of CRONFI Project. Moreover, Regulation on the Greenhouse Gases Emissions Monitoring in the Republic of Croatia (Official Gazette No. 1/07), which will come into force in 2008, prescribes monitoring of disturbances such as fires, pests. Table 7.2-2 provides information on factors used in estimations.

*Table 7.2-2: Emission factors used in estimations*

Forest type	D	BEF1	R	BEF2	CF
Coniferous	0.4	1.15	0.23	1.3	0.5
Deciduous	0.588	1.2	0.24	1.4	0.5

- D = basic wood density, calculated from table 3A.1.9 (GPG 2003) according to major tree species in growing stock in Croatia
- BEF1 = biomass expansion factor for conversion of annual net increment (including bark) to aboveground tree biomass, dimensionless (GPG 2003)
- R = root to shoot ratio, dimensionless; (GPG 2003, Table 3A.1.8)
- BEF2 = biomass expansion factor for conversion volumes of extracted roundwood to total aboveground (including bark) biomass, dimensionless (GPG 2003)
- CF = carbon fraction of dry matter (default = 0.5), tonnes C (tonne d. m. )<sup>-1</sup>

Table 7.2-3 provides information on Annual change in Carbon Stock in living biomass in Forest land remaining forest land.

*Table 7.2-3: Annual change in Carbon Stock in living biomass in Forest Land Remaining Forest Land (Gg CO<sub>2</sub>)*

Year	Annual increase in carbon stocks (Gg CO <sub>2</sub> )	Annual decrease in carbon due to carbon loss (Gg CO <sub>2</sub> )	Annual change in carbon stock in living biomass (Gg CO <sub>2</sub> )
1990	14855.32	8574.09	6281.22
1991	14855.32	4059.16	10796.16
1992	14855.32	4021.75	10833.57
1993	14855.32	3860.74	10994.58
1994	14855.32	4116.51	10738.81
1995	14855.32	4226.76	10628.56
1996	14855.32	4886.29	9969.03
1997	14855.32	5883.96	8971.36
1998	14855.32	6532.69	8322.63
1999	14855.32	6628.64	8226.67
2000	14855.32	6843.51	8011.81
2001	14855.32	5975.51	8879.81
2002	14855.32	6318.16	8537.16
2003	14855.32	6977.67	7877.64
2004	14855.32	6906.99	7948.33
2005	14855.32	7076.34	7778.98

### 7.2.3. UNCERTAINTIES AND TIME-SERIES CONSISTENCY

The uncertainty of the input data was estimated at 40 to 50 percent, while uncertainty of using wood density and BEFs was estimated at 30 percent.

Emissions from sub-sector Forest Land Remaining Forest Land have been calculated using the same data source for every year in the time series.

### 7.2.4. SOURCE-SPECIFIC QA/QC AND VERIFICATION

During the preparation of the inventory submission activities related to quality control were mainly focused on completeness and consistency of emission estimates and on proper use of notation keys in the CRF tables. After preparation of final draft of this chapter an audit was carried out to check selected activities from Tier 1 General inventory level QC procedures which revealed that most of the activities were correctly carried out during preparation of the inventory despite the fact that formal QC procedures were not prepared.

### 7.2.5. SOURCE-SPECIFIC RECALCULATIONS

Data sets for the period from 1990 to 2005 were included in the calculation of emissions and removals from LULUCF instead of incorrectly reported period in the previous report starting with 1986. New values for root-to-shoot ratios (R), biomass expansion factors (BEF1, BEF2) and basic wood densities (D), were included in the calculations of emissions and removals from LULUCF according to GPG 2003 and recalculations for the period 1990-2005 were done.

### 7.2.6. SOURCE-SPECIFIC PLANNED IMPROVEMENTS

Major areas for improvement:

- Development of land use database needed for greenhouse gas inventories with aim to collect more quality data from existing databases and use of complete land inventories. Regulation on the Greenhouse Gases Emissions Monitoring in the Republic of Croatia (Official Gazette No. 1/07), prescribes obligation and procedure for emissions monitoring. Among others, the regulation prescribes monitoring of areas within different land use categories, such as forest area, agricultural area, grasslands, wetlands, settlements and other land.
- Development of country specific factors (BEFs).
- Obtaining more accurate data for Forest Land Remaining Forest Land, especially for annual increment of biomass. Regulation on the Greenhouse Gases Emissions Monitoring in the Republic of Croatia (Official Gazette No. 1/07) prescribes monitoring of forest area, annual biomass increment, annual fellings, annual fuelwood gathering and other disturbances including fires, pests etc.



### 7.3. REFERENCES

Central Bureau of Statistics, *Statistical Yearbooks (1986-2005)*, Zagreb

Hrvatske šume (1996) *Šumskogospodarska osnova područja Republike Hrvatske*, Javno poduzeće "Hrvatske šume", Zagreb

IPCC (2003) *Good Practice Guidelines for Land Use Land-Use Change and Forestry*, IGES, Japan

IPCC (2000) *Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories*, Japan

IPCC/UNEP/OECD/IEA (1997) *Greenhouse Gas Inventory Workbook*, Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories, Volume 2, United Kingdom

IPCC/UNEP/OECD/IEA (1997) *Greenhouse Gas Inventory Reference Manual*, Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories, Volume 3, United Kingdom

Klepac D. (1998) *Hrvatsko šumarstvo u XIX. I XX. Stoljeću, Sveučilišna šumarska nastava u Hrvatskoj 1898-1998, knjiga II*, Šumarski fakultet Sveučilišta u Zagrebu, 9-31, Zagreb

Matić S. (1999) *The forests of Croatia – country report, Virgin forests and forest reserves in Central and East european countries*, Development of forestry and renewable forest resources – Biotechnical faculty, 17-24, Ljubljana

Matić S., Meštrović Š., Vukelić J. (1997) *Gospodarenje šumama i šumskim prostorom na području grada Zagreba i Županije Zagrebačke*, Šumarski fakultet Sveučilišta u Zagrebu, 213 str., Zagreb

Ministry of Environmental Protection, Physical Planning and Construction (2006) *National Inventory Report 2006, Croatian greenhouse gas inventory for the period 1990 – 2004*, EKONERG, Zagreb

Ministry of Environmental Protection, Physical Planning and Construction (2006) *Second, Third and Fourth National Communication of the Republic of Croatia under the United Nations Framework Convention on Climate Change (UNFCCC)*, Zagreb

Sabadi, R. (1994) *Pregled šumarstva i drvoprerađivačkog sektora u Republici Hrvatskoj*, Ministarstvo poljoprivrede i šumarstva Republike Hrvatske i "Hrvatske šume" p.o. Zagreb, 120 str., Zagreb

UNECE data base ; [www.unece.com](http://www.unece.com)

## 8. WASTE (CRF sector 6)

### 8.1. OVERVIEW OF SECTOR

Waste management activities such as disposal and treatment of municipal solid waste and wastewaters can produce emissions of greenhouse gases (GHGs) including methane ( $\text{CH}_4$ ), carbon dioxide ( $\text{CO}_2$ ) and nitrous oxide ( $\text{N}_2\text{O}$ ).

$\text{CH}_4$  emissions as a result of disposal and treatment of municipal solid waste,  $\text{CH}_4$  emissions from disposal of domestic and commercial wastewater in septic tanks, indirect  $\text{N}_2\text{O}$  emissions from human sewage and  $\text{CO}_2$  emissions resulting from incineration of hazardous and clinical waste (without energy recovery) are included in emissions estimates in this sector.

The methodology used to estimate emissions from waste management activities requires country-specific knowledge on waste generation, composition and management practice. The fact that waste management activities in Croatia are generally inadequately organized and implemented results in the lack and inconsistency of data. Therefore, effort was done in order to evaluate and compile data coming from different sources and adjust them to recommended Intergovernmental Panel on Climate Change (IPCC) methodology which is used for GHGs emissions estimation.

Regulation on the Greenhouse Gases Emissions Monitoring in the Republic of Croatia (Official Gazette No. 1/07), which will go into force in 2008, prescribes obligation and procedure for emissions monitoring, which comprise estimation and/or reporting of all anthropogenic emissions and removals. According to requirement, sources of abovementioned greenhouse gases should report required activity data for more accurate emissions estimation.

The total annual emissions of GHGs, expressed in Gg  $\text{CO}_2$ -eq, from waste management in the period 1990-2005 are presented in the Figure 8.1-1.




Figure 8.1-1: Emissions of GHGs from Waste (1990-2005)

## 8.2. SOLID WASTE DISPOSAL ON LAND (CRF 6.A.)

### 8.2.1. SOURCE CATEGORY DESCRIPTION

Landfill gas consists of approximately 50 percent CO<sub>2</sub> and 50 percent CH<sub>4</sub> by volume. Anaerobic decomposition of organic matter in Solid Waste Disposal Sites (SWDSs) results in the release of CH<sub>4</sub> to the atmosphere. The composition of waste is one of the main factors influencing the amount and the extent of CH<sub>4</sub> production within SWDSs. Temperature, moisture content and pH are important physical factors influencing fermentation of degradable organic substances and gas production.

### 8.2.2. METHODOLOGICAL ISSUES

A method used to calculate CH<sub>4</sub> emissions according to *Revised 1996 IPCC Guidelines* is First Order Decay (FOD) method.

The quantity of CH<sub>4</sub> emitted during decomposition process is directly proportional to the fraction of degradable organic carbon (DOC), which is defined as the carbon content of different types of organic biodegradable wastes such as paper and textiles, garden and park waste, food waste, wood and straw waste. DOC was estimated by using country-specific data on waste composition and quantities based on compiled data from Potočnik, V. (2000), *Report: The basis for methane emissions estimation in Croatia 1990-1998, B. Data on Municipal Solid Waste in Croatia 1990-1998*. Country-specific composition of waste is presented in Table 8.2-1.

Table 8.2-1: Country-specific composition of waste

Waste stream	Percent in the MSW	Percent DOC
Paper and textiles	21 - 22	40
Garden and park waste	18 - 19	17
Food waste	23 - 24	15
Wood and straw waste	3	30

The country-specific fraction of DOC in municipal solid waste (MSW), according to data from Table 8.2-1, was estimated to be 0.17 in the period 1990-2004 and 0.16 in 2005.

The decomposition of DOC does not occur completely and some of the potentially degradable materials always remain in the site over a long period of time. According to *Good Practice Guidance* approximately 50-60 percent of total DOC actually degrades<sup>6</sup> and converts to landfill gas. A mean value, i.e. 55 percent, was taken into account for the purpose of CH<sub>4</sub> emissions estimation from SWDSs.

The methodology provides a classification of SWDSs into “managed” and “unmanaged” sites through knowledge of site activities carried out. Unmanaged sites are further divided as deep ( $\geq 5$ m depth) or shallow ( $< 5$ m depth). The classification is used to apply a methane correction factor (MCF) to account for the methane generation potential of the site.

<sup>6</sup> The *Revised 1996 IPCC Guidelines* provide a default value of 77 percent for DOC that is converted to landfill gas, but this value, according to review of recent literature, is too high.

Land disposal is the only method of MSW management in Croatia. Quality and composition of disposed MSW and the main characteristic of SWDSs have been evaluated for the entire time series. Historical data for the total amount of generated waste and disposed MSW for the period 1970-1990 have been estimated based on national rate for waste generation and fraction of MSW disposed at different types of SWDSs. Extrapolation/interpolation methods has been used to obtain insufficient data. Total annual MSW disposed to SWDSs for the period 1990-1998 has been evaluated from available relevant data compiled into Report; Fundurulja, D., Mužinić, M. (2000) *Estimation of the Quantities of Municipal Solid Waste in the Republic of Croatia in the period 1990 – 1998 and 1998 – 2010*, Zagreb. Data for the quantity of disposed MSW in 1999 were evaluated by interpolation method. Data for the quantity of disposed MSW in 2000 were obtained from *Report of Environment Condition*, Ministry of Environmental Protection, Physical Planning and Construction. Data for the quantity of disposed MSW in 2004 were obtained from *The State of the Environment Report*, Croatian Environment Agency. Taking into account the pattern over 2000 and 2004 (total quantity of disposed MSW), quantity of MSW disposed to different types of SWDSs and the main characteristic of SWDSs for the period 2000 to 2004 were evaluated by interpolation/extrapolation method. Data for the quantity of disposed MSW in 2005 were obtained from *Waste Management Plan in the Republic of Croatia (2007-2015)*.

The most of managed SWDSs are not covered with aerated material and because of that default value for oxidation factor (OX), which equals zero, has been used.

The total annual MSW disposed to different types of SWDSs in the period 1990-2005 and related MCF are reported in Table 8.2-2.

*Table 8.2-2: Total annual MSW disposed to SWDSs and related MCF (1990-2005)*

Year	Managed SWDS (Gg)	Unmanaged SWDS ( $\geq 5m$ ) (Gg)	Unmanaged SWDS ( $< 5m$ ) (Gg)	MCF (fraction)
1990	18	277	295	0.606
1991	19	280	300	0.606
1992	20	284	309	0.605
1993	22	297	324	0.606
1994	26	322	329	0.613
1995	31	364	342	0.623
1996	35	392	361	0.625
1997	40	433	375	0.632
1998	45	470	398	0.636
1999*	54	538	383	0.654
2000	60	618	260	0.702
2001*	62	631	273	0.700
2002*	67	672	238	0.716
2003*	73	718	209	0.731
2004*	79	769	201	0.738
2005*	101	970	215	0.749

\* data on the annual MSW disposed to different types of SWDSs were obtained by interpolation/extrapolation method.

The resulting annual emissions of CH<sub>4</sub> from land disposal of MSW in the period 1990-2005 are presented in the Figure 8.2-1.




Figure 8.2-1: Emissions of CH<sub>4</sub> from Solid Waste Disposal on Land (1990-2005)

### 8.2.3. UNCERTAINTIES AND TIME-SERIES CONSISTENCY

The uncertainties contained in CH<sub>4</sub> emissions estimates are related primarily to assessment of historical data for quantity of MSW disposed to different types of SWDSs and the main characteristic of SWDSs as well as the usage of default IPCC methane generation rate constant ( $k=0.05$ ).

In addition, SWDSs in Croatia are classified into several categories, according to applied waste management activities, legality, volume (capacity and quantity of disposed MSW) as well as status. Municipal solid waste which is disposed to "Official" SWDSs is in most cases collected in an organized manner by registered companies. "Official" SWDSs do not necessarily fall under managed SWDSs category as defined by IPCC (site management activities carried out in "Official" SWDSs in most cases do not meet requirements to be characterized as managed). "Unofficial" SWDSs can be described as locations where all sorts of waste are dumped uncontrollably without any site management activities carried out. In order to adjust country-specific to IPCC SWDSs classification it was proposed that all "Unofficial" SWDSs fall under unmanaged shallow sites (<5m), whereas "Official" SWDSs fall under all three IPCC categories depending on management activities and dimensions of waste disposal sites. It is obvious that this distribution represents additional uncertainty in the estimation of country-specific MCF.

Another uncertainty is related to estimation of degradable organic carbon (DOC) in MSW. There were only few sorting of waste in Croatia, and in consequence of that these results were compared and adjusted to relevant data in similar countries.

Uncertainty estimate associated with emission factor amounts 50 percent, accordingly to provided uncertainty assessment in *Good Practice Guidance*. Uncertainty estimate associated with activity data amounts 50 percent, based on expert judgements.

Emissions from Solid waste Disposal on Land have been calculated using the same method for every year in the time series. Different source of information were used for data sets.

#### **8.2.4. SOURCE-SPECIFIC QA/QC AND VERIFICATION**

During the preparation of the inventory submission activities related to quality control were mainly focused on completeness and consistency of emission estimates and on proper use of notation keys in the CRF tables.

After preparation of final draft of this chapter an audit was carried out to check selected activities from Tier 1 General inventory level QC procedures and Tier 2 source-specific QC procedures. Regarding to Tier 2 activities, emission factors and activity data were checked for key source categories. Solid waste disposal on land represent key source category in Waste sector. CH<sub>4</sub> emissions from solid waste disposal on land were estimated using Tier 2 method which is a *good practice*. The uncertainty of activity data is very high due to high discrepancy between various data sources.

#### **8.2.5. SOURCE SPECIFIC RECALCULATIONS**

There are no source-specific recalculations in sub-sector Solid Waste Disposal on Land.

In the previous report, CH<sub>4</sub> emission estimations, calculated according to First Order Decay (FOD method), were recalculated for entire time series. Application of more complex model led to significantly lower CH<sub>4</sub> emissions from the Waste sector during the all observed period, as it was expected. Default IPCC Tier 1 method, which was previously used, was not provide an accurate trend, because it give a reasonable annual estimate of actual emissions if the amount of waste disposed have been constant. Because of consequence that amount of waste varying over time, emission calculations were overestimated. FOD method more accurately reflects the emissions trend over time. The results of the FOD model show that the emissions from this sector continuously increases mainly as a consequence of the larger amounts of waste disposed at the landfills and absence of secondary measures for emission reduction at the landfills (flaring, electricity production from the landfill gas).

#### **8.2.6. SOURCE-SPECIFIC PLANNED IMPROVEMENTS**

According to National Environmental Action Plan (NEAP) (Official Gazette No. 46/02), Croatian Waste Management Strategy (Official Gazette No. 130/05) and *Waste Management Plan in the Republic of Croatia (2007-2015)* (Official Gazette No. 85/07), infrastructure development for integral system of waste management has been emphasized, respectively, conditions for effectively waste management activities are created. Consequently, more accurate data for CH<sub>4</sub> emission calculations should be available.

##### **8.2.6.1. Activity data improvement**

By-law on Cadastre of Emission to Environment (Official Gazette No. 36/96) and The Waste Law (Official Gazette No. 178/04, 111/06) define administration commitments of manufacturers and all entities which contributed in waste management. The base for systematic gathering and

saving activity data was created by establishment of the Revision of Cadastre of Waste Disposal Sites (KEO). This will present part of new KEO software which is developed as a electronic managed data base with georeferent application (*Geographical Information System, GIS*) and access to the data base through web site of Croatian Environment Agency. By means of data base in GIS-tools, assessment and quantitative categorization of waste disposal sites are provided.

For the purposes of improvement activity data gathering from solid waste disposal activities it is necessary to improve quality of existing data:

- equipping the major landfills with automatic weigh-bridges in order to accurately estimate the quantities of delivered MSW;
- providing methodology to determine country-specific MSW composition;
- periodic analysis of waste composition at major landfills according to provided methodology;
- modification of Cadastre of Emission to Environment (KEO) Reporting Forms regarding to MSW with additional information on waste quantities and composition.

#### **8.2.6.2. Emission factor and methodology improvement**

For the purposes of emission inventory improvement it is necessary to adjust country-specific to IPCC SWDSs classification, in order to accurately estimate the MCF. Due to lack of adequate information, extrapolation method has been applied for estimation of waste and landfills characteristics over a long period of time. For the purposes of emission inventory improvement it is necessary to improve the quality of existing data and to reconstruct historical data. It is also necessary to apply a unique methodology to determinate waste quantity and composition.

By-law on Conditions for Waste Treatment (Official Gazette No. 123/97, 112/01) as well as By-law on Waste Management (Official Gazette No. 23/07) defines priority for improvement and organization of disposal sites and waste disposal on managed disposal sites.

## 8.3. WASTEWATER HANDLING (CRF 6.B.)

### 8.3.1. SOURCE CATEGORY DESCRIPTION

Aerobic biological process is used mostly in wastewater treatment. Anaerobic process is applied in some industrial wastewater treatment. Total amount of gas is flared in these treatments, and therefore all methane from gas is oxidized to carbon dioxide and water vapour.

Disposal of domestic and commercial wastewater, particularly in rural areas where systems such as septic tanks are used, are partly anaerobic without flaring, which results with CH<sub>4</sub> emissions. CH<sub>4</sub> emissions from these systems are estimated only for 2004 and 2005, because activity data for CH<sub>4</sub> emission calculations for the period 1990-2003 are not available.

Indirect N<sub>2</sub>O emissions from human sewage are included in emission estimates.

### 8.3.2. METHODOLOGICAL ISSUES

#### 8.3.2.1. Domestic and commercial wastewater

Methane emissions from domestic and commercial wastewater have been calculated using the methodology proposed by *Revised 1996 IPCC Guidelines*, by multiplying the total domestic organic wastewater in kg BOD/yr and emission factor which was obtained using default value for maximum methane producing capacity (0.25 kg CH<sub>4</sub>/kg BOD).

Data for calculation of degradable organic component in kg BOD/1000 person/yr were obtained by state company Croatian Water Resources Management (Hrvatske vode). Data, which are based on the Croatian Water Management Strategy, are available only for 2004 and 2005.

#### 8.3.2.2. Human sewage

Indirect nitrous oxide (N<sub>2</sub>O) emissions from human sewage have been calculated using the methodology proposed by *Revised 1996 IPCC Guidelines*, by multiplying annual per capita protein intake, fraction of nitrogen in protein, number of people in country and default emission factor which equals 0.01 kg N<sub>2</sub>O-N/kg sewage N produced.


The population estimate of the Republic of Croatia for the period 1990-2005 were taken from Statistical Yearbook 2006. Croatian data on the annual per capita Protein intake value (PIV), for the period 1992-2003, were obtained by the FAOSTAT Statistical Database. Adjustment method "Extrapolation of emissions based on a driver" has been used for calculation of insufficient data. Taking into account the PIV trend, the pattern over three years from 1992 to 1994 has been used for calculation of N<sub>2</sub>O emissions in 1990 and 1991. Conservativeness factor, proposed by the *Technical Guidance on Methodologies for Adjustments under Article 5, Paragraph 2, of the Kyoto Protocol*, was used for the base year (1990). Emissions values for 2002 and 2003 have been used as the pattern for N<sub>2</sub>O emission calculation in 2004 and 2005 by extrapolation method. Data for N<sub>2</sub>O emission calculation from Human Sewage for the period 1990-2005 are presented in the Table 8.3.1.



Table 8.3-1: Data for N<sub>2</sub>O emission calculation from Human Sewage (1990-2005)

Year	Protein intake (kg/person/yr)	Population
1990	20.71	4778000
1991	21.53	4513000
1992	22.16	4470000
1993	21.86	4641000
1994	22.96	4649000
1995	25.00	4669000
1996	24.78	4494000
1997	24.38	4572000
1998	23.98	4501000
1999	24.86	4554000
2000	24.67	4381000
2001	26.1	4437000
2002	27.52	4443000
2003	26.94	4442000
2004	26.38	4439000
2005	25.80	4442000

The resulting annual emissions of N<sub>2</sub>O from Human Sewage in the period 1990-2005 are presented in the Figure 8.3-1.

Figure 8.3-1: Emissions of N<sub>2</sub>O from Human Sewage (1990-2005)

### 8.3.3. UNCERTAINTIES AND TIME-SERIES CONSISTENCY

The uncertainties contained in N<sub>2</sub>O emissions estimates are related primarily to applied default emission factor and extrapolated values for protein intake.

Uncertainty estimate associated with emission factor amounts 30 percent, accordingly to provided uncertainty assessment in *Good Practice Guidance*. Uncertainty estimate associated with activity data amounts 10 percent, based on expert judgements.

Emissions from Human Sewage have been calculated using the same method and data sets for every year in the time series, except insufficient data which have been calculated by extrapolation method.

#### **8.3.4. SOURCE-SPECIFIC QA/QC AND VERIFICATION**

During the preparation of the inventory submission activities related to quality control were mainly focused on completeness and consistency of emission estimates and on proper use of notation keys in the CRF tables. After preparation of final draft of this chapter an audit was carried out to check selected activities from Tier 1 General inventory level QC procedures.

#### **8.3.5. SOURCE SPECIFIC RECALCULATIONS**

There are no source-specific recalculations in sub-sector Wastewater Handling.

#### **8.3.6. SOURCE-SPECIFIC PLANNED IMPROVEMENTS**

Improvements in the sub-sector Disposal of domestic and commercial wastewater are related primarily to aggregation of data for CH<sub>4</sub> emission calculations from systems such as septic tanks which are used in rural areas. CH<sub>4</sub> emissions from these systems are estimated only for 2004 and 2005, because activity data for CH<sub>4</sub> emission calculations for the period 1990-2003 are not available. For the purpose of completeness of inventory, it is necessary to collect and assess data for entire time series.

In order to accurate calculation of N<sub>2</sub>O emissions from Human Sewage, Croatia planned to analyze the influence of tourism on the population influx due to summer months, as well as fact that nearly 25 percent of the Croatian population lives close to the sea, which has influence on the emission factor.

## **8.4. WASTE INCINERATION (CRF 6.C.)**

### **8.4.1. SOURCE CATEGORY DESCRIPTION**

Incineration of waste produces emissions of CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O. According to *Revised 1996 IPCC Guidelines* only CO<sub>2</sub> emissions resulting from incineration of carbon in waste of fossil origin (e.g. plastics, textiles, rubber, liquid solvents and waste oil) without energy recovery, should be included in emissions estimates from Waste sector. Emissions from incineration with energy recovery should be reported in the Energy sector.

An incinerator of hazardous waste was functioning in Croatia between 1998 and 2002. CO<sub>2</sub> emissions from incineration of hazardous waste are not estimated because data for categorisation of waste types is lacking.

There is also incineration of clinical waste. Emissions from incineration of clinical waste are estimated only for 2004 and 2005, because activity data for CO<sub>2</sub> emission calculations for the period 1990-2003 are not available.

### **8.4.2. METHODOLOGICAL ISSUES**

CO<sub>2</sub> emissions from incineration of clinical waste have been calculated using the methodology proposed by *Revised 1996 IPCC Guidelines*, by multiplying the total incinerated clinical waste with default values for fraction of carbon content, fraction of fossil carbon and burn out efficiency of combustion.

Data for quantity of incinerated waste were obtained by Croatian Environment Agency. Data are available only for 2004 and 2005 and are accepted from Cadastre of Emission to Environment (KEO) Reporting Forms regarding to hazardous waste.

### **8.4.3. UNCERTAINTIES AND TIME-SERIES CONSISTENCY**

The uncertainties contained in CO<sub>2</sub> emissions estimates from incineration of clinical waste are related primarily to applied default emission factor.

Uncertainty estimate associated with emission factor amounts 30 percent, accordingly to provided uncertainty assessment in *Good Practice Guidance*. Uncertainty estimate associated with activity data amounts 10 percent, based on expert judgements.

### **8.4.4. SOURCE-SPECIFIC QA/QC AND VERIFICATION**

During the preparation of the inventory submission activities related to quality control were mainly focused on completeness and consistency of emission estimates and on proper use of notation keys in the CRF tables. After preparation of final draft of this chapter an audit was carried out to check selected activities from Tier 1 General inventory level QC procedures.

#### **8.4.5. SOURCE SPECIFIC RECALCULATIONS**

There are no source-specific recalculations in sub-sector Waste Incineration.

#### **8.4.6. SOURCE-SPECIFIC PLANNED IMPROVEMENTS**

Improvements in the sub-sector Waste Incineration are related primarily to aggregation of data for CO<sub>2</sub> emission calculations from incineration of hazardous and clinical waste. CO<sub>2</sub> emissions from incineration of clinical waste are estimated only for 2004 and 2005, because activity data for emission calculations for the period 1990-2003 are not available. For the purpose of inventory completeness, it is necessary to collect and assess data for entire time series.

## 8.5. EMISSION OVERVIEW

Emissions of GHGs from Waste in the period 1990-2005 are presented in Table 8.5-1.

Table 8.5-1: Emissions from Waste (1990-2005)

Source	Year	GHG	Emission (Gg)	GWP <sup>1</sup>	Emission (Gg CO <sub>2</sub> -eq)	Percent in Waste	Percentage in Total Country Emission
<b>Solid Waste Disposal on Land</b>	1990	CH <sub>4</sub>	10.53	21	221.21	74.15	0.70
	1991		11.12		233.57	75.51	0.94
	1992		11.71		245.84	76.10	1.08
	1993		12.32		258.72	76.59	1.12
	1994		12.98		272.60	76.62	1.23
	1995		13.74		288.59	76.03	1.28
	1996		14.57		305.92	77.90	1.31
	1997		15.48		325.17	78.92	1.31
	1998		16.45		345.38	80.41	1.38
	1999		17.53		368.16	80.67	1.41
	2000		18.62		391.10	82.28	1.51
	2001		19.71		414.01	82.10	1.53
	2002		20.82		437.24	82.10	1.55
	2003		21.96		461.25	83.18	1.55
	2004		23.18		486.81	75.85	1.62
	2005		22.36		469.50	80.56	1.54
<b>Domestic and Commercial Wastewater</b>	1990-2003	CH <sub>4</sub>	NE	21	-	-	-
	2004		3.03		63.65	9.92	0.21
	2005		1.14		23.95	4.11	0.08
<b>Human Sewage</b>	1990	N <sub>2</sub> O	0.25	310	77.12	25.85	0.24
	1991		0.24		75.73	24.49	0.30
	1992		0.25		77.21	23.90	0.34
	1993		0.26		79.07	23.41	0.34
	1994		0.27		83.20	23.38	0.38
	1995		0.29		90.98	23.97	0.40
	1996		0.28		86.80	22.10	0.37
	1997		0.28		86.88	21.08	0.35
	1998		0.27		84.13	19.59	0.34
	1999		0.28		88.24	19.33	0.34
	2000		0.27		84.24	17.72	0.33
	2001		0.29		90.26	17.90	0.33
	2002		0.31		95.30	17.90	0.34
	2003		0.30		93.27	16.82	0.31
	2004		0.29		91.27	14.22	0.30
	2005		0.29		89.33	15.33	0.29
<b>Waste Incineration</b>	1990-2003	CO <sub>2</sub>	NE	1	-	-	-
	2004		0.08		0.08	0.01	0.0003
	2005		0.03		0.03	0.01	0.0001

<sup>1</sup> Time horizon chosen for GWP values is 100 years

NE - emission is not estimated

## 8.6. REFERENCES

- APO (2005) *Revision of Cadastre of Waste Disposal Sites, Final Draft*, Zagreb
- Central Bureau of Statistics, *Statistical Yearbooks (1990-2006)*
- Diaz, L.F. et al. (1996) *Solid Waste Management for Economically Developing Countries*, ISWA
- Government of the Republic of Croatia (2005) *Croatian Waste Management Strategy*
- FAOSTAT: Statistical Database, <http://www.fao.org>
- Fundurulja, D., Mužinić, M. (2000) *Estimation of the Quantities of Municipal Solid Waste in the Republic of Croatia in the period 1990 – 1998 and 1998 – 2010*, Zagreb
- IPCC (2000) *Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories*, Japan
- IPCC/UNEP/OECD/IEA (1997) *Greenhouse Gas Inventory Workbook*, Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories, Volume 2, United Kingdom
- IPCC/UNEP/OECD/IEA (1997) *Greenhouse Gas Inventory Reference Manual*, Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories, Volume 3, United Kingdom
- Ministry of Environmental Protection, Physical Planning and Construction: *Report of Environment Condition*, <http://www.mzopu.hr/okolis>
- Ministry of Environmental Protection, Physical Planning and Construction (2006) *National Inventory Report 2006, Croatian greenhouse gas inventory for the period 1990 – 2004*, EKONERG, Zagreb
- Ministry of Environmental Protection, Physical Planning and Construction (2006) *Second, Third and Fourth National Communication of the Republic of Croatia under the United Nations Framework Convention on Climate Change (UNFCCC)*, Zagreb
- Ministry of Environmental Protection, Physical Planning and Construction (2007) *Waste Management Plan in the Republic of Croatia (2007-2015)*, Zagreb
- Potočnik, V. (2000) *Report: The basis for methane emission estimation in Croatia 1990-1998, B. Data on Municipal Solid Waste in Croatia 1990-1998*, Zagreb
- Schaller, A. (2000) *Republic of Croatia: First National Communication, Waste Management Review – Waste Disposal Sites*, Zagreb

## 9. RECALCULATIONS AND IMPROVEMENTS

The key differences between the previous and latest submission of CRF tables for the time series 1990-2005 are outlined in this chapter. Detailed description and explanations for recalculations are shown in recalculation sections in the sector chapters, Chapters 3 to 8.

### 9.1. EXPLANATIONS AND JUSTIFICATIONS FOR RECALCULATIONS, INCLUDING IN RESPONSE TO THE REVIEW PROCESS

The recalculations are performed in accordance with:

- 1) decisions of sectoral experts
- 2) suggestions of expert review team

According to suggestions reported in "Report of the individual review of the greenhouse gas inventory of Croatia submitted in 2006" Croatia performed recalculations in the following sectors:

- Energy (Road Transportation, Manufacturing industry and consumption, Other stationary - Non energy fuel combustion, CO<sub>2</sub> emission from scrubbing);
- Industrial Processes (Cement Production, Production of other chemical, Consumption of halocarbons and SF<sub>6</sub>);
- Solvent and other product use;
- N<sub>2</sub>O emission from manure management, Direct emission from agriculture soils, Direct N<sub>2</sub>O emissions from animals, Indirect N<sub>2</sub>O emission from nitrogen used in agriculture;
- Land Use, Land-Use Change and Forestry

In this section, the summary of the recalculations performed and justification is given using the following categories of distinction:

- Changes or refinements in methods (Chapter 9.1.1.)
- Correction of errors (Chapter 9.1.2.)

#### 9.1.1. CHANGES OR REFINEMENTS IN METHODS

The following methodological changes were made for the calculation of greenhouse gases according to:

- Changes in available data;
- Consistency with good practice guidance;
- New methods.

**Changes in available data:****Industrial processes**

**Mineral products (2.A.); Cement Production (2.A.1.)** - in the previous report, two different data source were used across the reporting years. For 1990-2001 years data were obtained from document "Data for drafting baseline scenarios and analysis of GHG emission reductions in the Republic of Croatia" (J. Garilovic), which were cross-checked with clinker and cement production data from Monthly Industrial Reports published by Central Bureau of Statistics, Department of Manufacturing and Mining. For 2002-2004 years data obtained from survey of cement manufacturers, which were cross-checked with data from Monthly Industrial Reports. In this report, clinker and cement production data for the period 1990-2005 were collected from survey of cement manufacturers due to improving comparability and time series consistency. Also, country specific EFs and its CKD correction factor were developed.

**Chemical Industry (2.B.); Production of other chemicals (2.B.5.)** - in the previous report errors were done during activity data compilation for 2003 and 2004 - units of activity data were not in correct values (e.g. kilogramme instead of tonnes).

**Metal Production (2.C.); Ferroalloys Production (2.C.2.)** - in the previous report the production of ferrochromium and associated emissions were insufficient for 1999. In this report data for 1999 was calculated by interpolation method.

**Consumption of Halocarbons and SF<sub>6</sub> (2.F.); Refrigeration and Air Conditioning Equipment (2.F.1.)** - in the previous report, HFCs emissions for the period 1990-1994 (HFC 134a), 1990-1995 (HFC 125, 143a) and 1990-1999 (HFC 32) have not been calculated because of the difficulty in obtaining the relevant activity data for these estimates. In this report, data have been calculated by extrapolation method.

**Agriculture**

**N<sub>2</sub>O Emissions from Manure Management (4.B.); Direct Emission of N<sub>2</sub>O from Animals (4.D.2.); Indirect N<sub>2</sub>O Emissions from Nitrogen Used in Agriculture (4.D.3.)** - data for goat and mule stocks were included in calculations of N<sub>2</sub>O emissions according to *Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories*.

**Direct Emission from Agricultural Soils (4.D.1.)** - in the previous report insufficient data was estimated as average value for 2001, 2002 and 2003. In this report adjustment method "Extrapolation of emissions based on a driver", proposed by the *Technical Guidance on Methodologies for Adjustments under Article 5, Paragraph 2, of the Kyoto Protocol*, has been used for calculation of insufficient data. The pattern over the period 1992-2003 has been used for the estimation of pulses and soyabeans and non-N-fixing crop production for 2004 and 2005.



### **Land-use, Land Use Change and Forestry**

Data sets for the period from 1990 to 2005 were included in the calculation of emissions and removals from LULUCF instead of incorrectly reported period in the previous report starting with 1986. New values for root-to-shoot ratios (R), biomass expansion factors (BEF1, BEF2) and basic wood densities (D), were included in the calculations of emissions and removals from LULUCF according to GPG 2003.

### **Consistency with good practice guidance:**

#### **Energy**

**Road Transportation (1.AA.3.B)** – the justification for the recalculation of CO<sub>2</sub>, N<sub>2</sub>O and CH<sub>4</sub> emissions from Road Transport is trend consistency. Therefore, the Tier 2/3 methodology was applied for the entire period from 1990-2005.

**Manufacturing industry and construction (1.A.2)** - energy consumption of natural gas in petrochemical industry is calculated under 1.A.2.f Manufacturing industries and construction – Other – Petrochemical Industry. Tier 1 methodology for calculation of CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O emissions from Manufacturing industries and construction – Other – Petrochemical Industry was used for calculation of all GHG emissions (in previous submission energy use of fuels in Petrochemical industry wasn't reported).

**Other Stationary - Non energy fuel consumption (1.A.5)** - according to suggestion of review team, reconciliation between the reference and sectoral approaches was made. Natural gas and liquid fuels from non energy fuel consumption is reported in 1.A.d. instead of 1.A.5.a.

**Production and processing (1.B.2.b(ii))** - according to suggestion of review team, CO<sub>2</sub> emission from scrubbing is reported in subcategory 1.B.2.b(ii) Production and processing instead of 1.B.2.d Other, for whole time series, although activity data doesn't exist (country specific issue).

### **New methods:**

#### **Energy**

**Road Transportation (1.A.3.b.)** - according to suggestion of review team, Tier 3 method for CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O emissions calculation is used, because of trend consistency reasons.

**Solvent and other product use (3)** - in the previous report, CO<sub>2</sub> emissions were not calculated because IPCC Guidelines do not provide methodology for calculation of CO<sub>2</sub> emissions from Solvent and Other Product Use. In this report, CO<sub>2</sub> emissions are calculated by means of conversion factor which contains ratio C/NMVOC = 0.8 and recalculation ratio of C to CO<sub>2</sub> equal to 44/12. The overall conversion factor has value of 2.93.

### 9.1.2. CORRECTION OF ERRORS

The majority of recalculations performed were due to errors which can be divided as follows:

- typing errors;
- errors regarding notation keys.

All the explanations for errors of these types are provided directly in the CRF Reporter. Table 8(b) Recalculation – explanatory information doesn't provide information on such errors due to large amount of these types of errors.

## 9.2. THE IMPLICATION OF THE RECALCULATIONS ON THE LEVEL AND TREND, INCLUDING TIME SERIES CONSISTENCY

This section outlines the implications over time for the emission levels as well as the implications for emission trends, including time-series consistency.

Table 9.2-1 shows the differences between the last submission (NIR 2006) and current submission (NIR 2007), on the level of the different greenhouse gases.

*Table 9.2-1: Differences between NIR 2006 and NIR 2007 for 1990-2004 due to recalculations*

Gas	Source	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
CO <sub>2</sub> (Tg) Incl. LULUCF	NIR 2006	8.60	2.02	1.03	1.74	-0.36	-4.29	-3.65	2.81	-1.53	-0.58
	NIR 2007	17.03	6.42	5.60	5.96	5.52	6.25	7.64	9.69	11.24	12.13
	<b>Difference %</b>	<b>98</b>	<b>218</b>	<b>444</b>	<b>243</b>	<b>-1633</b>	<b>-246</b>	<b>-309</b>	<b>245</b>	<b>-835</b>	<b>-2192</b>
CO <sub>2</sub> (Tg) Excl. LULUCF	NIR 2006	23.03	16.74	15.81	16.43	15.69	16.25	16.94	18.02	18.91	19.70
	NIR 2007	23.31	17.22	16.44	16.96	16.26	16.88	17.61	18.66	19.57	20.36
	<b>Difference %</b>	<b>1.2</b>	<b>2.9</b>	<b>4.0</b>	<b>3.2</b>	<b>3.6</b>	<b>3.9</b>	<b>4.0</b>	<b>3.6</b>	<b>3.5</b>	<b>3.4</b>
CH <sub>4</sub> (CO <sub>2</sub> -eq Gg)	NIR 2006	3233	3007	2826	2771	2564	2532	2557	2624	2460	2496
	NIR 2007	3247	3018	2835	2779	2573	2540	2565	2631	2466	2501
	<b>Difference %</b>	<b>0.4</b>	<b>0.4</b>	<b>0.3</b>	<b>0.3</b>	<b>0.4</b>	<b>0.3</b>	<b>0.3</b>	<b>0.3</b>	<b>0.2</b>	<b>0.2</b>
N <sub>2</sub> O (CO <sub>2</sub> -eq Gg)	NIR 2006	3920	3827	3601	3200	3207	3123	3004	3348	2912	3103
	NIR 2007	4017	3908	3498	3221	3224	3116	3058	3398	2955	3148
	<b>Difference %</b>	<b>2.5</b>	<b>2.1</b>	<b>-2.9</b>	<b>0.6</b>	<b>0.5</b>	<b>-0.2</b>	<b>1.8</b>	<b>1.5</b>	<b>1.5</b>	<b>1.4</b>
PFCs (CO <sub>2</sub> -eq Gg)	NIR 2006	937	642	0	0	0	0	0	0	0	0
	NIR 2007	937	642	0	0	0	0	0	0	0	0
	<b>Difference %</b>	<b>0.0</b>	<b>0.1</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>
HFCs (CO <sub>2</sub> -eq Gg)	NIR 2006	0.00	0.00	0.00	0.00	0.00	7.80	60.15	91.18	17.54	9.09
	NIR 2007	43.38	51.71	50.52	49.34	48.15	43.20	60.26	91.28	17.64	9.18
	<b>Difference %</b>						<b>453.8</b>	<b>0.2</b>	<b>0.1</b>	<b>0.6</b>	<b>1.0</b>
SF <sub>6</sub> (CO <sub>2</sub> -eq Gg)	NIR 2006	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	NIR 2007	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	<b>Difference %</b>	<b>0.00</b>	<b>0.00</b>	<b>0.00</b>	<b>0.00</b>	<b>0.00</b>	<b>0.00</b>	<b>0.00</b>	<b>0.00</b>	<b>0.00</b>	<b>0.00</b>
Total (Tg CO <sub>2</sub> -eq) Incl. LULUCF	NIR 2006	16.69	9.49	7.46	7.71	5.41	1.38	1.97	3.26	3.86	5.03
	NIR 2007	25.27	14.04	11.99	12.01	11.36	11.95	13.33	15.81	16.68	17.79
	<b>Difference %</b>	<b>51.4</b>	<b>48.0</b>	<b>60.7</b>	<b>55.8</b>	<b>110.0</b>	<b>765.7</b>	<b>576.4</b>	<b>385.1</b>	<b>332.2</b>	<b>253.7</b>
Total (Tg CO <sub>2</sub> -eq) Excl. LULUCF	NIR 2006	31.12	24.21	22.24	22.40	21.46	21.91	22.56	24.09	24.30	25.31
	NIR 2007	31.55	24.84	22.82	23.01	22.10	22.58	23.29	24.78	25.00	26.02
	<b>Difference %</b>	<b>1.4</b>	<b>2.6</b>	<b>2.6</b>	<b>2.7</b>	<b>3.0</b>	<b>3.0</b>	<b>3.3</b>	<b>2.9</b>	<b>2.9</b>	<b>2.8</b>

Table 9.2-1: Differences between NIR 2006 and NIR 2007 for 1990-2004 due to recalculations (cont.)

Gas	Source	2000	2001	2002	2003	2004
CO <sub>2</sub> (Tg) Incl. LULUCF	NIR 2006	0.13	2.66	4.70	6.23	6.23
	NIR 2007	11.93	12.07	13.49	15.62	15.20
	<b>Difference %</b>	<b>9081</b>	<b>354</b>	<b>187</b>	<b>151</b>	<b>144</b>
CO <sub>2</sub> (Tg) Excl. LULUCF	NIR 2006	19.42	20.43	21.50	22.88	22.55
	NIR 2007	19.95	20.94	22.03	23.50	23.15
	<b>Difference %</b>	<b>2.7</b>	<b>2.5</b>	<b>2.5</b>	<b>2.7</b>	<b>2.6</b>
CH <sub>4</sub> (CO <sub>2</sub> -eq Gg)	NIR 2006	2544	2690	2745	2925	3015
	NIR 2007	2548	2691	2746	2925	3015
	<b>Difference %</b>	<b>0.2</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>
N <sub>2</sub> O (CO <sub>2</sub> -eq Gg)	NIR 2006	3284	3251	3317	3221	3677
	NIR 2007	3317	3289	3358	3263	3680
	<b>Difference %</b>	<b>1.0</b>	<b>1.2</b>	<b>1.2</b>	<b>1.3</b>	<b>0.1</b>
PFCs (CO <sub>2</sub> -eq Gg)	NIR 2006	0	0	0	0	0
	NIR 2007	0	0	0	0	0
	<b>Difference %</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>
HFCs (CO <sub>2</sub> -eq Gg)	NIR 2006	23.15	49.00	49.32	163.7	188.9
	NIR 2007	23.15	49.00	49.32	163.7	188.9
	<b>Difference %</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>
SF <sub>6</sub> (CO <sub>2</sub> -eq Gg)	NIR 2006	0.00	0.00	0.00	0.00	0.00
	NIR 2007	0.00	0.00	0.00	0.00	0.00
	<b>Difference %</b>	<b>0.00</b>	<b>0.00</b>	<b>0.00</b>	<b>0.00</b>	<b>0.00</b>
Total (Tg CO <sub>2</sub> -eq) Incl. LULUCF	NIR 2006	5.98	8.65	10.81	12.54	13.11
	NIR 2007	17.82	18.09	19.65	21.98	22.08
	<b>Difference %</b>	<b>198.0</b>	<b>109.2</b>	<b>81.8</b>	<b>75.2</b>	<b>68.4</b>
Total (Tg CO <sub>2</sub> -eq) Excl. LULUCF	NIR 2006	25.27	26.42	27.61	29.19	29.43
	NIR 2007	25.84	26.97	28.19	29.85	30.03
	<b>Difference %</b>	<b>2.2</b>	<b>2.1</b>	<b>2.1</b>	<b>2.3</b>	<b>2.0</b>

The change in the 1990-2005 trend for the greenhouse gas emissions compared to the previous submission is presented in Table 9.2-2. It can be concluded that the trend in the total national emissions decreased by 8.81 percent including LULUCF and 0.62 percent excluding LULUCF compared to NIR 2006. The largest absolute changes in emission trends are recorded for CO<sub>2</sub>, HFCs and total CO<sub>2</sub>-eq, described in Table 9.2-2.

Table 9.2-2: Differences between NIR 2006 and NIR 2007 for the emission trends 1990-2004

Gas CO <sub>2</sub> -eq (Gg)	Trend (absolute)			Trend (percent)		
	NIR 2006*	NIR 2007**	Difference	NIR 2006*	NIR 2007**	Difference
CO <sub>2</sub> emissions including net CO <sub>2</sub> from LULUCF	-2367.43	-1826.97	-540.46	-27.54	-10.73	-16.81
CO <sub>2</sub> emissions excluding net CO <sub>2</sub> from LULUCF	-483.47	-159.86	-323.61	-2.10	-0.69	-1.41
CH <sub>4</sub>	-217.85	-231.74	13.88	-6.74	-7.14	0.40
N <sub>2</sub> O	-242.66	-337.86	95.21	-6.19	-8.41	2.22
HFCs	188.87	145.49	43.38	100.00	335.42	-235.42
PFCs	-936.56	-936.56	0.00	-100.00	-100.00	0.00
<b>Total (including LULUCF)</b>	<b>-3575.64</b>	<b>-3187.64</b>	<b>-388.00</b>	<b>-21.43</b>	<b>-12.61</b>	<b>-8.81</b>
<b>Total (excluding LULUCF)</b>	<b>-1691.68</b>	<b>-1520.53</b>	<b>-171.15</b>	<b>-5.44</b>	<b>-4.82</b>	<b>-0.62</b>

\* - Difference, in previous submission, between emissions in 2004 and 1990 (absolute and percent)

\*\* - Difference, in latest submission, between emissions in 2004 and 1990 (absolute and percent)

### 9.3. PLANNED IMPROVEMENTS TO THE INVENTORY

The framework for development of Croatian greenhouse gas emissions inventory was established during preparation of the First National Communication to the UNFCCC in 2001. The framework was built upon experiences and lessons learned from the previously established scheme for national reporting and international data exchange through the EEA/ETC-ACC system and reporting under Convention on Long-range Transboundary Air Pollution (CLRTAP). Since then Croatia has submitted National Inventory Reports in 2003 for period 1995-2001, in 2004 for period 1990-2002, in 2005 for period 1990-2003, in 2006 for period 1990-2004 and this latest submission in October 2007.

Generally, Croatia has developed a sound and well-documented greenhouse gas inventory system in only a few years but it still requires continuous improvements in almost all key elements related to compilation and submission of the inventory. In order to fulfill these requirements Croatia has taken strategic approach and as a result a draft of National GHG Inventory Improvement Strategy has been prepared<sup>7</sup>. The purpose of this strategic document is to recognize strengths and weaknesses of the existing national GHG inventory system and to determine a realistic short- and long- term objectives in order to establish cost-effective GHG inventory preparation system that will enable timely, accurate, transparent and consistent international reporting, taking into account national circumstances, resources and available information.

There are several priority tasks for improvements of the inventory system which are outlined in the strategy:

- preparation of Regulation on the Greenhouse Gas Emissions Monitoring in the Republic of Croatia, which is confirmed in January 2007 (Official Gazette No. 1/2007). It should improve existing system of greenhouse gas emission monitoring and reporting in accordance with the requirements of the Kyoto protocol and relevant legislation of the EU (Decision 280/2004/EC) and defines institutional responsibilities and mandates for national inventory compilation. It is expected that the Regulation will enter into force in 2008;
- authorization of appropriate national institution to be in charge of approving the inventory;
- establish national reference centre for air and climate change;
- ensuring sustainable inventory preparation process including establishment of QA/QC system;
- carrying out awareness-raising campaign targeting policy-makers and other stakeholders on importance and benefits of sustainable inventory process;
- improving collection of activity data, emission factors and overall emission calculation for key sources, based on long term inventory preparation program;
- increasing the financial, technical and human resources for inventory preparation, based on long-term inventory program.

<sup>7</sup> National GHG Inventory Improvement Strategy was prepared under UNDP/GEF regional project Capacity Building for Improving the Quality of GHG Inventories (Europe and CIS Region).

Sector specific goals are outlined below:

## **ENERGY**

### Short-term goals (< 1 years)

Generally, the changes from Tier 1 to Tier 2/3 estimation methodologies for Energy key sources are recommended. The priority should be the key sources with high uncertainties of emission estimation. But, significant constraints are availability of activity data, especially for the beginning years of concerned period. Consequently, implementation of more detailed methodology approach (Tier 2/3) for key sources, for entire period, will be very difficult.

COPERT III software (Tier 3) is used for emission estimation from Road Transport. The difficulties lie in gathering of detailed activity data (1990-2000). In any case, the improvement of the emission estimation for entire period, based on COPERT model results, is a short-term goal.

The extensive use of detailed methodology (Tier 2/3) for Energy Industries is also one of the short-term goals. For achievement of abovementioned goal is necessary to ensure delivery of detailed activity data for Energy Industries. The good example is the usage of technology/plant-specific data for sub-sector Thermal power plants and public cogeneration plants (Tier 2) for the last five inventory years.

### Long-term goals (> 1 years)

For estimation of fugitive emissions from oil and natural gas operations, a Tier 1 method was applied. Used emission factors are an average value of the range proposed in the IPCC Manual. However, fugitive emission from natural gas is key source and implementation of rigorous source-specific evaluations approach (Tier 3) is necessary. The Tier 3 approach will generally involve compiling the following types of information:

- detailed inventories of the amount and types of process infrastructure (e.g. wells, field installations and production/processing facilities);
- production disposition analyses oil and gas production, vented, flared and reinjected volumes of gas, and fuel gas consumption;
- accidental releases (i.e. well blow-outs and pipeline ruptures);
- typical design and operating practices and their impact on the overall level of emission control.

The extensive use of plant-specific data collected in the framework of Cadastre of Emissions to Environment (CEE) is recommended ("bottom up" approach). In addition, usage of more source-specific QA/QC procedures will improve the quality of GHG inventory in Energy sector.

## **INDUSTRIAL PROCESSES AND SOLVENT AND OTHER PRODUCT USE**

### Short-term goals (< 1 years)

Uncertainty of emission estimation is mainly caused by implementation of default IPCC emission factors. Consequently, wider use of well documented country-specific (technology-specific and plant-specific) emission factors, in sectors Industrial Processes and Solvent and

Other Product Use, is an important short-term goal. The use of country-specific EFs, where available, as a way to minimize uncertainty, is recommended.

Short-term goals are also improvements of halocarbons and SF<sub>6</sub> emission estimations.

There are gaps in the time series of some productions, provided by statistical institutions. Filling these gaps by using direct surveys and comparison with time series of other related data is recommended.

#### Long-term goals (> 1 years)

As a small country with a small number of plants and good-quality production statistics, Croatia has often adopted higher-tier methodologies for Industrial Processes, based on plant-level information. Croatia considers wider use of source-specific verification procedures, through systematic cross-checking of plant-specific information with production statistics, and also the use other sources of information, such as CEE and the national energy balance.

### **AGRICULTURE**

#### Short-term goals (< 1 years)

The QA/QC procedures should be applied to avoid possible mistakes. Expert group for agriculture should be established in the aim to improve the quality of inventory and data collection. The applicability of higher-tier methods for key sources should be explored, depending on availability of data.

#### Long-term goals (> 1 years)

Source-specific explorations should be done in order to determinate country-specific emission factors and collect detailed set of activity data, which will improve CH<sub>4</sub> and N<sub>2</sub>O emission calculation (Tier 2/3) in key agriculture sources, such as: CH<sub>4</sub> Emission from Enteric Fermentation, N<sub>2</sub>O Emission from Manure Management, Direct N<sub>2</sub>O emission from Agricultural Soils and Animals and N<sub>2</sub>O Emission from Nitrogen Used in Agriculture.

### **LAND-USE CHANGE AND FORESTRY**

#### Short-term goals (< 1 years)

The Republic of Croatia only reports data for changes in the forest and other woody biomass stocks. Expert group for calculations emissions/removals from LULUCF sector was established by Ministry of Agriculture, Forestry and Water Management in the May 2005, with the objective to improve the quality of data collection and inventory preparation. Expert group should in future choose criteria for data collection, finding the model to choose most reliable existing data, respecting continuity, quality and reciprocal comparability, as well as accessibility of sources. Usage of new very detailed CRF tables, as much as possible, is also short-term goal.

### Long-term goals (> 1 years)

The GHG emissions/removals calculation for sub-sectors Forest and Grassland Conversion (5B), Abandonment of Managed Lands (5C) and CO<sub>2</sub> Emissions and Removals from Soil (5D) should be included, as much as possible, depending on availability of activity data. It is planned to complete those sub-sectors in future inventories, and to do inventory according to the Good Practice Guidelines for Land Use, Land Use Change and Forestry.

## **WASTE**

### Short-term goals (< 1 years)

Croatia plans to improve its waste statistics and to carry out sector-specific studies related to Solid Waste Disposal in order to use the Tier 2 method.

By-law on Cadastre of Emission to Environment (Official Gazette No. 36/96) and The Waste Law (Official Gazette 151/03) define administration commitments of manufacturers and all entities which contributed in waste management. The base for systematic gathering and saving activity data was created by establishment of the Revision of Cadastre of Waste Disposal Sites (CEE). This will present part of new CEE software which is developed as an electronic managed data base with georeferent application (*Geographical Information System, GIS*) and access to the data base through web site of Croatian Environment Agency. By means of data base in GIS-tools, assessment and quantitative categorization of waste disposal sites will be provided.

By-law on Conditions for Waste Treatment (Official Gazette No. 123/97, 112/01) as well as By-law on Waste Management (Official Gazette No. 23/07) defines priority for improvement and organization of disposal sites and waste disposal on managed disposal sites.

For the purposes of emission inventory improvement it is necessary to adjust country-specific to IPCC SWDSs classification, in order to accurately estimate the MCF.

Also, it is necessary to apply a unique methodology to determine waste quantity and composition. For the purposes of improvement activity data gathering from solid waste disposal activities it is necessary to improve quality of existing data:

- equipping the major landfills with automatic weigh-bridges in order to accurately estimate the quantities of delivered MSW;
- providing methodology to determine country-specific MSW composition;
- periodic analysis of waste composition at major landfills according to provided methodology;
- modification of Cadastre of Emission to Environment (CEE) Reporting Forms regarding to MSW with additional information on waste quantities and composition.

### Long-term goals (> 1 years)

New waste statistics and sector-specific studies should be used to reconstruct historical activity data in applying the Tier 2 method for key source Solid Waste Disposal on Land.

Emissions from Wastewater Handling and Waste Incineration should be included for the sake of completeness.



## **ANNEX 1**

### **KEY CATEGORIES**

## A1.1. DESCRIPTION OF METHODOLOGY USED FOR IDENTIFYING KEY CATEGORIES

Key categories according to the IPCC Good Practice Guidance (IPCC, 2000) are those found in the accumulative 95% of the total annual emissions in the last reported year or belonging to the total trend, when ranked from contributing the largest to smallest share in annual total and in the trend. As originally designed it applied only to source categories. In addition, *Good Practice Guidance for Land Use, Land-Use Change and Forestry* expands the original approach to enable the identification of key categories that are either sources or sinks, which provides on how to identify key categories for the LULUCF. Therefore, the key category analysis was determined using both approaches:

- excluding LULUCF
- including LULUCF

Following the *Good Practice Guidelines*, Croatia undertook a key category analysis using Tier 1 Level and Trend methods.

The IPCC and *Good Practice Guidance for Land Use, Land-Use Change and Forestry* also recommended which sources should be checked for their key category status, Table A1-1. Additionally, other sources of direct greenhouse gas emissions not listed in above mentioned guidances were added to the list, e.g. CO<sub>2</sub> Emissions from Natural Gas Scrubbing, CO<sub>2</sub> Emissions from Solvent and Other Product Use, CO<sub>2</sub> Emissions from Non energy-use in Industrial Processes reported under 2.G Other non-specified NEU in CRF Reporter.

*Table A1-1: Categories Assessed in Key Category Analysis*

Source Categories Assessed in Key Source Category Analysis	Direct GHG	Special Considerations
<b>ENERGY SECTOR</b>		
CO <sub>2</sub> Emissions from Stationary Combustion - Coal	CO <sub>2</sub>	
CO <sub>2</sub> Emissions from Stationary Combustion - Oil	CO <sub>2</sub>	
CO <sub>2</sub> Emissions from Stationary Combustion - Gas	CO <sub>2</sub>	
Non-CO <sub>2</sub> Emissions from Stationary Combustion	CH <sub>4</sub>	
Non-CO <sub>2</sub> Emissions from Stationary Combustion	N <sub>2</sub> O	
Mobile Combustion - Road Vehicles	CO <sub>2</sub>	
Mobile Combustion - Road Vehicles	CH <sub>4</sub>	
Mobile Combustion - Road Vehicles	N <sub>2</sub> O	
Mobile Combustion: Water-borne Navigation	CO <sub>2</sub>	
Mobile Combustion: Water-borne Navigation	CH <sub>4</sub>	
Mobile Combustion: Water-borne Navigation	N <sub>2</sub> O	
Mobile Combustion: Aircraft	CO <sub>2</sub>	
Mobile Combustion: Aircraft	CH <sub>4</sub>	
Mobile Combustion: Aircraft	N <sub>2</sub> O	
Mobile Combustion - Agriculture/Forestry/Fishing	CO <sub>2</sub>	
Mobile Combustion - Agriculture/Forestry/Fishing	CH <sub>4</sub>	
Mobile Combustion - Agriculture/Forestry/Fishing	N <sub>2</sub> O	
Fugitive Emissions from Coal Mining and Handling	CH <sub>4</sub>	
Fugitive Emissions from Oil and Gas Operations	CH <sub>4</sub>	
CO <sub>2</sub> Emissions from Natural Gas Scrubbing	CO <sub>2</sub>	IPCC doesn't offer methodology for estimating emission of CO <sub>2</sub> scrubbed from natural gas and subsequently emitted into atmosphere. Natural gas produced in Croatian gas fields has a large amount of CO <sub>2</sub> , more than 15 percent. The maximum volume content of CO <sub>2</sub> in commercial natural gas is 3 percent and gas must be cleaned before coming to pipeline and transport to users. Because of that, the Scrubbing Units exist at largest Croatian gas field. The CO <sub>2</sub> , scrubbed from natural gas, is emitted into atmosphere. The emission is estimated by material balance method.
<b>INDUSTRIAL SECTOR</b>		
CO <sub>2</sub> Emissions from Cement Production	CO <sub>2</sub>	
CO <sub>2</sub> Emissions from Lime Production	CO <sub>2</sub>	
CO <sub>2</sub> Emissions from Iron and Steel Production	CO <sub>2</sub>	

Table A1-1: Categories Assessed in Key Category Analysis (cont.)

N <sub>2</sub> O Emissions from Nitric Acid Production	N <sub>2</sub> O	
N <sub>2</sub> O Emissions from Adipic Acid Production	N <sub>2</sub> O	
PFC Emissions from Aluminium production	PFC	
CO <sub>2</sub> Emissions from Ammonia Production	CO <sub>2</sub>	
CO <sub>2</sub> Emissions from Ferroalloys Production	CO <sub>2</sub>	
CO <sub>2</sub> Emissions from Aluminium production	CO <sub>2</sub>	
Sulfur hexafluoride (SF <sub>6</sub> ) from Magnesium Production	SF <sub>6</sub>	
SF <sub>6</sub> Emissions from Electrical Equipment	SF <sub>6</sub>	
SF <sub>6</sub> Emissions from Other Sources of SF <sub>6</sub>	SF <sub>6</sub>	
SF <sub>6</sub> Emissions from Production of SF <sub>6</sub>	SF <sub>6</sub>	
PFC, HFC, SF <sub>6</sub> Emissions from Semiconductor manufacturing		
Emissions from Substitutes for Ozone Depleting Substances (ODS Substitutes)		
HFC-23 Emissions from HCFC-22 Manufacture	HFC-23	
HFC Emissions from Consumption of HFCs, PFCs and SF <sub>6</sub>	HFC	
CO <sub>2</sub> Emissions from Non energy-use in Industrial Processes	CO <sub>2</sub>	
<b>SOLVENT AND OTHER PRODUCT USE</b>	CO <sub>2</sub>	
<b>AGRICULTURE SECTOR</b>		
CH <sub>4</sub> Emissions from Enteric Fermentation in Domestic Livestock	CH <sub>4</sub>	
CH <sub>4</sub> Emissions from Manure Management	CH <sub>4</sub>	
N <sub>2</sub> O Emissions from Manure Management	N <sub>2</sub> O	
CH <sub>4</sub> and N <sub>2</sub> O Emissions from Savanna Burning		
CH <sub>4</sub> and N <sub>2</sub> O Emissions from Agricultural Residue Burning		
Direct N <sub>2</sub> O Emissions from Agricultural Soils	N <sub>2</sub> O	
N <sub>2</sub> O Emissions from Pasture, Range and Paddock Manure	N <sub>2</sub> O	
Indirect N <sub>2</sub> O Emissions from Nitrogen Used in Agriculture	N <sub>2</sub> O	
CH <sub>4</sub> Emissions from Rice Cultivation	CH <sub>4</sub>	
<b>LULUCF</b>		
Forest land remaining forest land	CO <sub>2</sub>	
Forest land remaining forest land	CH <sub>4</sub>	
Forest land remaining forest land	N <sub>2</sub> O	
<b>WASTE SECTOR</b>		
CH <sub>4</sub> Emissions from Solid Waste Disposal Sites	CH <sub>4</sub>	
Emissions from Waste Water Handling	CH <sub>4</sub>	
Emissions from Waste Water Handling	N <sub>2</sub> O	
Emissions from Waste Incineration	CO <sub>2</sub>	
Emissions from Waste Incineration	N <sub>2</sub> O	

The reference to the summary overview for Key Categories 2005 in CRF tables is the Excel file HRV-2007-2005-v1.1, Table 7.

The level of disaggregation is in accordance with the suggested source categories split of the *Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories* and additionally, with the LULUCF category following the *Good Practice Guidance for Land Use, Land-Use Change and Forestry*.

## A1.2. TABLES 7.A1-7.A3 OF THE IPCC GOOD PRACTICE GUIDANCE

Table A1-2: Key categories analysis – Level Assessment - Tier 1 (Excluding LULUCF)

Tier 1 Analysis - Level Assessment – Excluding LULUCF					
IPCC Source Categories	Direct GHG	Base Year (1990) Estimate (Gg eq-CO <sub>2</sub> )	Current Year (2005) Estimate (Gg eq-CO <sub>2</sub> )	Level Assessment	Cumulative Total (%)
CO <sub>2</sub> Emissions from Stationary Combustion: Oil	CO <sub>2</sub>	8424.076	6505.092	0.213	21%
Mobile Combustion: Road Vehicles	CO <sub>2</sub>	3504.033	5286.911	0.173	39%
CO <sub>2</sub> Emissions from Stationary Combustion: Gas	CO <sub>2</sub>	4053.828	4866.249	0.160	55%
CO <sub>2</sub> Emissions from Stationary Combustion: Coal	CO <sub>2</sub>	3141.488	2681.012	0.088	63%
CO <sub>2</sub> Emissions from Cement Production	CO <sub>2</sub>	1047.472	1495.013	0.049	68%
Fugitive Emissions from Oil and Gas Operations	CH <sub>4</sub>	1186.258	1333.922	0.044	73%
Direct N <sub>2</sub> O Emissions from Agricultural Soils	N <sub>2</sub> O	1347.606	1238.253	0.041	77%
CH <sub>4</sub> Emissions from Enteric Fermentation in Domestic Livestock	CH <sub>4</sub>	1343.853	822.423	0.027	79%
Indirect N <sub>2</sub> O Emissions from Nitrogen Used in Agriculture	N <sub>2</sub> O	916.638	816.932	0.027	82%
N <sub>2</sub> O Emissions from Nitric Acid Production	N <sub>2</sub> O	927.561	783.280	0.026	85%
Mobile Combustion: Agriculture/Forestry/Fishing	CO <sub>2</sub>	839.186	709.390	0.023	87%
CO <sub>2</sub> Emissions from Natural Gas Scrubbing	CO <sub>2</sub>	415.949	691.000	0.023	89%
CO <sub>2</sub> Emissions from Ammonia Production	CO <sub>2</sub>	491.551	511.214	0.017	91%
CH <sub>4</sub> Emissions from Solid Waste Disposal Sites	CH <sub>4</sub>	221.208	469.502	0.015	93%
HFC Emissions from Consumption of HFCs	HFC	43.377	349.080	0.011	94%
N <sub>2</sub> O Emissions from Manure Management	N <sub>2</sub> O	376.710	236.670	0.008	94%
N <sub>2</sub> O Emissions from Pasture, Range and Paddock Manure	N <sub>2</sub> O	251.785	223.461	0.007	95%
CO <sub>2</sub> Emissions from Lime Production	CO <sub>2</sub>	159.780	196.424	0.006	96%
Mobile Combustion: Aircraft	CO <sub>2</sub>	295.612	172.642	0.006	96%
Mobile Combustion: Road Vehicles	N <sub>2</sub> O	49.160	165.628	0.005	97%
CH <sub>4</sub> Emissions from Manure Management	CH <sub>4</sub>	227.409	157.086	0.005	97%
CO <sub>2</sub> Emissions from Solvent and Other Product Use	CO <sub>2</sub>	80.211	155.050	0.005	98%
Fuel Combustion: Stationary Sources	CH <sub>4</sub>	172.214	110.036	0.004	98%
Mobile Combustion: Water-borne Navigation	CO <sub>2</sub>	132.980	99.590	0.003	99%
Mobile Combustion: Railways	CO <sub>2</sub>	137.525	95.518	0.003	99%
Emissions from Waste Water Handling	N <sub>2</sub> O	77.117	89.325	0.003	99%
CO <sub>2</sub> Emissions from Other non-specified NEU	CO <sub>2</sub>	208.051	72.191	0.002	100%
Fuel Combustion: Stationary Sources	N <sub>2</sub> O	65.467	51.046	0.002	100%
Mobile Combustion: Road Vehicles	CH <sub>4</sub>	29.876	32.610	0.001	100%
Emissions from Waste Water Handling	CH <sub>4</sub>	-	23.951	0.001	100%
CO <sub>2</sub> Emissions from Soda Ash Production and Use	CO <sub>2</sub>	25.740	17.222	0.001	100%
CO <sub>2</sub> Emissions from Limestone and Dolomite Use	CO <sub>2</sub>	43.218	12.053	0.000	100%
Production of Other Chemicals	CH <sub>4</sub>	15.798	5.329	0.000	100%
Mobile Combustion: Agriculture/Forestry/Fishing	N <sub>2</sub> O	2.038	1.718	0.000	100%
Mobile Combustion: Aircraft	N <sub>2</sub> O	2.589	1.513	0.000	100%
Mobile Combustion: Agriculture/Forestry/Fishing	CH <sub>4</sub>	1.299	1.072	0.000	100%
CO <sub>2</sub> Emissions from Iron and Steel Production	CO <sub>2</sub>	0.867	0.337	0.000	100%
Mobile Combustion: Water-borne Navigation	N <sub>2</sub> O	0.337	0.253	0.000	100%
Mobile Combustion: Railways	N <sub>2</sub> O	0.390	0.242	0.000	100%
Mobile Combustion: Water-borne Navigation	CH <sub>4</sub>	0.190	0.143	0.000	100%
Mobile Combustion: Railways	CH <sub>4</sub>	0.213	0.137	0.000	100%
Emissions from Waste Incineration	CO <sub>2</sub>	-	0.034	0.000	100%
Mobile Combustion: Aircraft	CH <sub>4</sub>	0.044	0.026	0.000	100%
PFC Emissions from Aluminium production	PFC	936.564	-	0.000	100%
CO <sub>2</sub> Emissions from Ferroalloys Production	CO <sub>2</sub>	194.526	-	0.000	100%
CO <sub>2</sub> Emissions from Aluminium Production	CO <sub>2</sub>	111.372	-	0.000	100%
Fugitive Emissions from Coal Mining and Handling	CH <sub>4</sub>	48.757	-	0.000	100%
<b>TOTAL</b>		<b>31551.924</b>	<b>30480.582</b>		

Table A1-3: Key categories analysis – Level Assessment - Tier 1 (Including LULUCF)

Tier 1 Analysis - Level Assessment - Including LULUCF					
IPCC Source Categories	Direct GHG	Base Year (1990) Estimate (Gg eq-CO <sub>2</sub> )	Current Year (2005) Estimate (Gg eq-CO <sub>2</sub> )	Level Assessment	Cumulative Total (%)
Forest land remaining forest land	CO <sub>2</sub>	6281.223	7778.976	0.203	20%
CO <sub>2</sub> Emissions from Stationary Combustion: Oil	CO <sub>2</sub>	8424.076	6505.092	0.170	37%
Mobile Combustion: Road Vehicles	CO <sub>2</sub>	3504.033	5286.911	0.138	51%
CO <sub>2</sub> Emissions from Stationary Combustion: Gas	CO <sub>2</sub>	4053.828	4866.249	0.127	64%
CO <sub>2</sub> Emissions from Stationary Combustion: Coal	CO <sub>2</sub>	3141.488	2681.012	0.070	71%
CO <sub>2</sub> Emissions from Cement Production	CO <sub>2</sub>	1047.472	1495.013	0.039	75%
Fugitive Emissions from Oil and Gas Operations	CH <sub>4</sub>	1186.258	1333.922	0.035	78%
Direct N <sub>2</sub> O Emissions from Agricultural Soils	N <sub>2</sub> O	1347.606	1238.253	0.032	82%
CH <sub>4</sub> Emissions from Enteric Fermentation in Domestic Livestock	CH <sub>4</sub>	1343.853	822.423	0.021	84%
Indirect N <sub>2</sub> O Emissions from Nitrogen Used in Agriculture	N <sub>2</sub> O	916.638	816.932	0.021	86%
N <sub>2</sub> O Emissions from Nitric Acid Production	N <sub>2</sub> O	927.561	783.280	0.020	88%
Mobile Combustion: Agriculture/Forestry/Fishing	CO <sub>2</sub>	839.186	709.390	0.019	90%
CO <sub>2</sub> Emissions from Natural Gas Scrubbing	CO <sub>2</sub>	415.949	691.000	0.018	92%
CO <sub>2</sub> Emissions from Ammonia Production	CO <sub>2</sub>	491.551	511.214	0.013	93%
CH <sub>4</sub> Emissions from Solid Waste Disposal Sites	CH <sub>4</sub>	221.208	469.502	0.012	94%
HFC Emissions from Consumption of HFCs	HFC	43.377	349.080	0.009	95%
N <sub>2</sub> O Emissions from Manure Management	N <sub>2</sub> O	376.710	236.670	0.006	96%
N <sub>2</sub> O Emissions from Pasture, Range and Paddock Manure	N <sub>2</sub> O	251.785	223.461	0.006	96%
CO <sub>2</sub> Emissions from Lime Production	CO <sub>2</sub>	159.780	196.424	0.005	97%
Mobile Combustion: Aircraft	CO <sub>2</sub>	295.612	172.642	0.005	97%
Mobile Combustion: Road Vehicles	N <sub>2</sub> O	49.160	165.628	0.004	98%
CH <sub>4</sub> Emissions from Manure Management	CH <sub>4</sub>	227.409	157.086	0.004	98%
CO <sub>2</sub> Emissions from Solvent and Other Product Use	CO <sub>2</sub>	80.211	155.050	0.004	98%
Fuel Combustion: Stationary Sources	CH <sub>4</sub>	172.214	110.036	0.003	99%
Mobile Combustion: Water-borne Navigation	CO <sub>2</sub>	132.980	99.590	0.003	99%
Mobile Combustion: Railways	CO <sub>2</sub>	137.525	95.518	0.002	99%
N <sub>2</sub> O Emissions from Waste Water Handling	N <sub>2</sub> O	77.117	89.325	0.002	99%
CO <sub>2</sub> Emissions from Other non-specified NEU	CO <sub>2</sub>	208.051	72.191	0.002	100%
Fuel Combustion: Stationary Sources	N <sub>2</sub> O	65.467	51.046	0.001	100%
Mobile Combustion: Road Vehicles	CH <sub>4</sub>	29.876	32.610	0.001	100%
Emissions from Waste Water Handling	CH <sub>4</sub>	-	23.951	0.001	100%
CO <sub>2</sub> Emissions from Soda Ash Production and Use	CO <sub>2</sub>	25.740	17.222	0.000	100%
CO <sub>2</sub> Emissions from Limestone and Dolomite Use	CO <sub>2</sub>	43.218	12.053	0.000	100%
Production of Other Chemicals	CH <sub>4</sub>	15.798	5.329	0.000	100%
Mobile Combustion: Agriculture/Forestry/Fishing	N <sub>2</sub> O	2.038	1.718	0.000	100%
Mobile Combustion: Aircraft	N <sub>2</sub> O	2.589	1.513	0.000	100%
Mobile Combustion: Agriculture/Forestry/Fishing	CH <sub>4</sub>	1.299	1.072	0.000	100%
CO <sub>2</sub> Emissions from Iron and Steel Production	CO <sub>2</sub>	0.867	0.337	0.000	100%
Mobile Combustion: Water-borne Navigation	N <sub>2</sub> O	0.337	0.253	0.000	100%
Mobile Combustion: Railways	N <sub>2</sub> O	0.390	0.242	0.000	100%
Mobile Combustion: Water-borne Navigation	CH <sub>4</sub>	0.190	0.143	0.000	100%
Mobile Combustion: Railways	CH <sub>4</sub>	0.213	0.137	0.000	100%
Emissions from Waste Incineration	CO <sub>2</sub>	-	0.034	0.000	100%
Mobile Combustion: Aircraft	CH <sub>4</sub>	0.044	0.026	0.000	100%
CO <sub>2</sub> Emissions from Ferroalloys Production	CO <sub>2</sub>	194.526	-	0.000	100%
CO <sub>2</sub> Emissions from Aluminium Production	CO <sub>2</sub>	111.372	-	0.000	100%
Fugitive Emissions from Coal Mining and Handling	CH <sub>4</sub>	48.757	-	0.000	100%
PFC Emissions from Aluminium production	PFC	936.564	-	0.000	100%
	<b>ABSOLUTE TOTAL</b>	<b>37833.147</b>	<b>38259.559</b>		

Table A1-4: Key categories analysis – Trend Assessment - Tier 1 (Excluding LULUCF)

Tier 1 Analysis - Trend Assessment – Excluding LULUCF						
IPCC Source Categories	Direct GHG	Base Year (1990) Estimate (Gg eq-CO <sub>2</sub> )	Last Year (2005) Estimate (Gg eq-CO <sub>2</sub> )	Trend Assessment	% Contribution to trend	Cumulative Total of Column F
Mobile Combustion: Road Vehicles	CO <sub>2</sub>	3504.033	5286.911	0.094	22.338%	22%
HFC Emissions from Consumption of HFCs	HFC	43.377	349.080	0.081	19.205%	42%
CO <sub>2</sub> Emissions from Stationary Combustion: Oil	CO <sub>2</sub>	8424.076	6505.092	0.041	9.734%	51%
CO <sub>2</sub> Emissions from Stationary Combustion: Gas	CO <sub>2</sub>	4053.828	4866.249	0.038	8.904%	60%
PFC Emissions from Aluminium production	PFC	936.564	-	0.031	7.275%	67%
CO <sub>2</sub> Emissions from Cement Production	CO <sub>2</sub>	1047.472	1495.013	0.023	5.370%	73%
CH <sub>4</sub> Emissions from Solid Waste Disposal Sites	CH <sub>4</sub>	221.208	469.502	0.018	4.222%	77%
CO <sub>2</sub> Emissions from Natural Gas Scrubbing*	CO <sub>2</sub>	415.949	691.000	0.016	3.738%	81%
Mobile Combustion: Road Vehicles	N <sub>2</sub> O	49.160	165.628	0.013	3.093%	84%
CO <sub>2</sub> Emissions from Stationary Combustion: Coal	CO <sub>2</sub>	3141.488	2681.012	0.010	2.321%	86%
CH <sub>4</sub> Emissions from Enteric Fermentation in Domestic Livestock	CH <sub>4</sub>	1343.853	822.423	0.010	2.254%	88%
Fugitive Emissions from Oil and Gas Operations	CH <sub>4</sub>	1186.258	1333.922	0.007	1.654%	90%
CO <sub>2</sub> Emissions from Ferroalloys Production	CO <sub>2</sub>	194.526	-	0.006	1.511%	92%
CO <sub>2</sub> Emissions from Solvent and Other Product Use	CO <sub>2</sub>	80.211	155.050	0.005	1.166%	93%
CO <sub>2</sub> Emissions from Aluminium Production	CO <sub>2</sub>	111.372	-	0.004	0.865%	94%
N <sub>2</sub> O Emissions from Nitric Acid Production	N <sub>2</sub> O	927.561	783.280	0.003	0.733%	94%
Mobile Combustion: Agriculture/Forestry/Fishing	CO <sub>2</sub>	839.186	709.390	0.003	0.659%	95%
N <sub>2</sub> O Emissions from Manure Management	N <sub>2</sub> O	376.710	236.670	0.003	0.619%	96%
Mobile Combustion: Aircraft	CO <sub>2</sub>	295.612	172.642	0.002	0.511%	96%
Indirect N <sub>2</sub> O Emissions from Nitrogen Used in Agriculture	N <sub>2</sub> O	916.638	816.932	0.002	0.467%	97%
Direct N <sub>2</sub> O Emissions from Agricultural Soils	N <sub>2</sub> O	1347.606	1238.253	0.002	0.442%	97%
CO <sub>2</sub> Emissions from Lime Production	CO <sub>2</sub>	159.780	196.424	0.002	0.404%	97%
Fugitive Emissions from Coal Mining and Handling	CH <sub>4</sub>	48.757	-	0.002	0.379%	98%
CO <sub>2</sub> Emissions from Other non-specified NEU	CO <sub>2</sub>	208.051	72.191	0.001	0.347%	98%
CH <sub>4</sub> Emissions from Manure Management	CH <sub>4</sub>	227.409	157.086	0.001	0.334%	99%
CO <sub>2</sub> Emissions from Ammonia Production	CO <sub>2</sub>	491.551	511.214	0.001	0.298%	99%
Fuel Combustion: Stationary Sources	CH <sub>4</sub>	172.214	110.036	0.001	0.279%	99%
Mobile Combustion: Railways	CO <sub>2</sub>	137.525	95.518	0.001	0.201%	99%
Mobile Combustion: Water-borne Navigation	CO <sub>2</sub>	132.980	99.590	0.001	0.167%	99%
N <sub>2</sub> O Emissions from Pasture, Range and Paddock Manure	N <sub>2</sub> O	251.785	223.461	0.001	0.134%	100%
Emissions from Waste Water Handling	N <sub>2</sub> O	77.117	89.325	0.001	0.134%	100%
Fuel Combustion: Stationary Sources	N <sub>2</sub> O	65.467	51.046	0.0003	0.073%	100%
CO <sub>2</sub> Emissions from Limestone and Dolomite Use	CO <sub>2</sub>	43.218	12.053	0.0003	0.064%	100%
CO <sub>2</sub> Emissions from Soda Ash Production and Use	CO <sub>2</sub>	25.740	17.222	0.0002	0.040%	100%
Mobile Combustion: Road Vehicles	CH <sub>4</sub>	29.876	32.610	0.0001	0.032%	100%
Production of Other Chemicals	CH <sub>4</sub>	15.798	5.329	0.0001	0.026%	100%
Mobile Combustion: Aircraft	N <sub>2</sub> O	2.589	1.513	0.00001	0.005%	100%
Mobile Combustion: Agriculture/Forestry/Fishing	N <sub>2</sub> O	2.038	1.718	0.000007	0.002%	100%
CO <sub>2</sub> Emissions from Iron and Steel Production	CO <sub>2</sub>	0.867	0.337	0.000006	0.002%	100%
Mobile Combustion: Agriculture/Forestry/Fishing	CH <sub>4</sub>	1.299	1.072	0.000005	0.001%	100%
Mobile Combustion: Railways	N <sub>2</sub> O	0.390	0.242	0.000003	0.001%	100%
Mobile Combustion: Water-borne Navigation	N <sub>2</sub> O	0.337	0.253	0.000002	0.0004%	100%
Mobile Combustion: Railways	CH <sub>4</sub>	0.213	0.137	0.000001	0.0003%	100%
Mobile Combustion: Water-borne Navigation	CH <sub>4</sub>	0.190	0.143	0.000001	0.0002%	100%
Mobile Combustion: Aircraft	CH <sub>4</sub>	0.044	0.026	0.0000003	0.0001%	100%
Emissions from Waste Water Handling	CH <sub>4</sub>	-	23.951	0.0000000	0.0000%	100%
Emissions from Waste Incineration	CO <sub>2</sub>	-	0.034	0.0000000	0.0000%	100%
	<b>TOTAL (excluding LULUCF)</b>	<b>31551.9</b>	<b>30480.6</b>			

Table A1-5: Key categories analysis – Trend Assessment - Tier 1 (Including LULUCF)

Tier 1 Analysis - Trend Assessment - Including LULUCF						
IPCC Source/Sink Categories	Direct GHG	Base Year (1990) Estimate (Gg eq-CO <sub>2</sub> )	Last Year (2005) Estimate (Gg eq-CO <sub>2</sub> )	Trend Assessment	% Contribution to trend	Cumulative Total
Mobile Combustion: Road Vehicles	CO <sub>2</sub>	3504.033	5286.911	0.145	20.864%	21%
Forest land remaining forest land	CO <sub>2</sub>	-6281.223	-7778.976	0.120	17.355%	38%
HFC Emissions from Consumption of HFCs	HFC	43.377	349.080	0.110	15.860%	54%
CO <sub>2</sub> Emissions from Stationary Combustion: Gas	CO <sub>2</sub>	4053.828	4866.249	0.067	9.682%	64%
PFC Emissions from Aluminium production	PFC	936.564	-	0.041	5.942%	70%
CO <sub>2</sub> Emissions from Cement Production	CO <sub>2</sub>	1047.472	1495.013	0.036	5.126%	75%
CO <sub>2</sub> Emissions from Stationary Combustion: Oil	CO <sub>2</sub>	8424.076	6505.092	0.033	4.731%	80%
CH <sub>4</sub> Emissions from Solid Waste Disposal Sites	CH <sub>4</sub>	221.208	469.502	0.026	3.681%	83%
CO <sub>2</sub> Emissions from Natural Gas Scrubbing	CO <sub>2</sub>	415.949	691.000	0.024	3.395%	87%
Mobile Combustion: Road Vehicles	N <sub>2</sub> O	49.160	165.628	0.018	2.609%	89%
Fugitive Emissions from Oil and Gas Operations	CH <sub>4</sub>	1186.258	1333.922	0.014	2.011%	91%
CH <sub>4</sub> Emissions from Enteric Fermentation in Domestic Livestock	CH <sub>4</sub>	1343.853	822.423	0.010	1.434%	93%
CO <sub>2</sub> Emissions from Ferroalloys Production	CO <sub>2</sub>	194.526	-	0.009	1.234%	94%
CO <sub>2</sub> Emissions from Solvent and Other Product Use	CO <sub>2</sub>	80.211	155.050	0.007	1.029%	95%
CO <sub>2</sub> Emissions from Aluminium Production	CO <sub>2</sub>	111.372	-	0.005	0.707%	96%
CO <sub>2</sub> Emissions from Stationary Combustion: Coal	CO <sub>2</sub>	3141.488	2681.012	0.004	0.568%	96%
CO <sub>2</sub> Emissions from Ammonia Production	CO <sub>2</sub>	491.551	511.214	0.003	0.497%	97%
CO <sub>2</sub> Emissions from Lime Production	CO <sub>2</sub>	159.780	196.424	0.003	0.427%	97%
N <sub>2</sub> O Emissions from Manure Management	N <sub>2</sub> O	376.710	236.670	0.003	0.388%	98%
Mobile Combustion: Aircraft	CO <sub>2</sub>	295.612	172.642	0.002	0.332%	98%
Fugitive Emissions from Coal Mining and Handling	CH <sub>4</sub>	48.757	-	0.002	0.309%	98%
Direct N <sub>2</sub> O Emissions from Agricultural Soils	N <sub>2</sub> O	1347.606	1238.253	0.002	0.252%	98%
CO <sub>2</sub> Emissions from Other non-specified NEU	CO <sub>2</sub>	208.051	72.191	0.002	0.247%	99%
N <sub>2</sub> O Emissions from Nitric Acid Production	N <sub>2</sub> O	927.561	783.280	0.001	0.211%	99%
CH <sub>4</sub> Emissions from Manure Management	CH <sub>4</sub>	227.409	157.086	0.001	0.195%	99%
Mobile Combustion: Agriculture/Forestry/Fishing	CO <sub>2</sub>	839.186	709.390	0.001	0.187%	99%
Fuel Combustion: Stationary Sources	CH <sub>4</sub>	172.214	110.036	0.001	0.173%	99%
Emissions from Waste Water Handling	N <sub>2</sub> O	77.117	89.325	0.001	0.154%	100%
Mobile Combustion: Railways	CO <sub>2</sub>	137.525	95.518	0.001	0.117%	100%
Mobile Combustion: Water-borne Navigation	CO <sub>2</sub>	132.980	99.590	0.001	0.087%	100%
CO <sub>2</sub> Emissions from Limestone and Dolomite Use	CO <sub>2</sub>	43.218	12.053	0.0003	0.046%	100%
Mobile Combustion: Road Vehicles	CH <sub>4</sub>	29.876	32.610	0.0003	0.042%	100%
Fuel Combustion: Stationary Sources	N <sub>2</sub> O	65.467	51.046	0.0002	0.035%	100%
CO <sub>2</sub> Emissions from Soda Ash Production and Use	CO <sub>2</sub>	25.740	17.222	0.0002	0.024%	100%
Indirect N <sub>2</sub> O Emissions from Nitrogen Used in Agriculture	N <sub>2</sub> O	916.638	816.932	0.0002	0.023%	100%
Production of Other Chemicals	CH <sub>4</sub>	15.798	5.329	0.0001	0.019%	100%
Mobile Combustion: Aircraft	N <sub>2</sub> O	2.589	1.513	0.00002	0.003%	100%
CO <sub>2</sub> Emissions from Iron and Steel Production	CO <sub>2</sub>	0.867	0.337	0.00001	0.001%	100%
N <sub>2</sub> O Emissions from Pasture, Range and Paddock Manure	N <sub>2</sub> O	251.785	223.461	0.00001	0.001%	100%
Mobile Combustion: Agriculture/Forestry/Fishing	N <sub>2</sub> O	2.038	1.718	0.000003	0.0005%	100%
Mobile Combustion: Agriculture/Forestry/Fishing	CH <sub>4</sub>	1.299	1.072	0.000003	0.0004%	100%
Mobile Combustion: Railways	N <sub>2</sub> O	0.390	0.242	0.000002	0.0004%	100%
Mobile Combustion: Water-borne Navigation	N <sub>2</sub> O	0.337	0.253	0.000002	0.0002%	100%
Mobile Combustion: Railways	CH <sub>4</sub>	0.213	0.137	0.000001	0.0002%	100%
Mobile Combustion: Water-borne Navigation	CH <sub>4</sub>	0.190	0.143	0.0000009	0.0001%	100%
Mobile Combustion: Aircraft	CH <sub>4</sub>	0.044	0.026	0.0000003	0.00004%	100%
Emissions from Waste Water Handling	CH <sub>4</sub>	-	23.951	0.0000000	0.000000%	100%
Emissions from Waste Incineration	CO <sub>2</sub>	-	0.034	0.0000000	0.000000%	100%
<b>TOTAL</b>		<b>25270.701</b>	<b>22701.606</b>			

Table A1-6: Key categories for Croatia – summary (Excluding LULUCF)

Tier 1 Analysis – Source Analysis Summary (Croatian Inventory)			
IPCC Source Categories	Direct GHG	Key Source Category Flag	Criteria for Identification
<b>ENERGY SECTOR</b>			
CO <sub>2</sub> Emissions from Stationary Combustion: Coal	CO <sub>2</sub>	Yes	Level, Trend
CO <sub>2</sub> Emissions from Stationary Combustion: Oil	CO <sub>2</sub>	Yes	Level, Trend
CO <sub>2</sub> Emissions from Stationary Combustion: Gas	CO <sub>2</sub>	Yes	Level, Trend
Non-CO <sub>2</sub> Emissions from Stationary Combustion	CH <sub>4</sub>	No	
Non-CO <sub>2</sub> Emissions from Stationary Combustion	N <sub>2</sub> O	No	
Mobile Combustion: Road Vehicles	CO <sub>2</sub>	Yes	Level, Trend
Mobile Combustion: Railways	CO <sub>2</sub>	No	
Mobile Combustion: Domestic Aviation	CO <sub>2</sub>	No	
Mobile Combustion: National Navigation	CO <sub>2</sub>	No	
Mobile Combustion: Agriculture/Forestry/Fishing	CO <sub>2</sub>	Yes	Level, Trend
Mobile Combustion: Road Vehicles	CH <sub>4</sub>	No	
Mobile Combustion: Railways	CH <sub>4</sub>	No	
Mobile Combustion: Domestic Aviation	CH <sub>4</sub>	No	
Mobile Combustion: National Navigation	CH <sub>4</sub>	No	
Mobile Combustion: Agriculture/Forestry/Fishing	CH <sub>4</sub>	No	
Mobile Combustion: Road Vehicles	N <sub>2</sub> O	Yes	Trend
Mobile Combustion: Railways	N <sub>2</sub> O	No	
Mobile Combustion: Domestic Aviation	N <sub>2</sub> O	No	
Mobile Combustion: National Navigation	N <sub>2</sub> O	No	
Mobile Combustion: Agriculture/Forestry/Fishing	N <sub>2</sub> O	No	
Fugitive Emissions from Coal Mining and Handling	CH <sub>4</sub>	No	
Fugitive Emissions from Oil and Gas Operations	CH <sub>4</sub>	Yes	Level, Trend
CO <sub>2</sub> Emissions from Natural Gas Scrubbing	CO <sub>2</sub>	Yes	Level, Trend
<b>INDUSTRIAL SECTOR</b>			
CO <sub>2</sub> Emissions from Cement Production	CO <sub>2</sub>	Yes	Level, Trend
CO <sub>2</sub> Emissions from Lime Production	CO <sub>2</sub>	No	
CO <sub>2</sub> Emissions from Limestone and Dolomite Use	CO <sub>2</sub>	No	
CO <sub>2</sub> Emissions from Soda Ash Production and Use	CO <sub>2</sub>	No	
CO <sub>2</sub> Emissions from Ammonia Production	CO <sub>2</sub>	Yes	Level
CO <sub>2</sub> Emissions from Iron and Steel Production	CO <sub>2</sub>	No	
CO <sub>2</sub> Emissions from Ferroalloys Production	CO <sub>2</sub>	Yes	Trend
CO <sub>2</sub> Emissions from Aluminium Production	CO <sub>2</sub>	Yes	Trend
CH <sub>4</sub> Emissions from Production of Other Chemicals	CH <sub>4</sub>	No	
N <sub>2</sub> O Emissions from Nitric Acid Production	N <sub>2</sub> O	Yes	Level, Trend
HFC Emissions from Consumption of HFCs	HFC	Yes	Level, Trend
PFC Emissions from Aluminium production	PFC	Yes	Trend
CO <sub>2</sub> Emissions from Other non-specified NEU	CO <sub>2</sub>	No	
<b>CO<sub>2</sub> EMISSIONS FROM SOLVENT AND OTHER PRODUCT USE</b>	CO <sub>2</sub>	Yes	Trend
<b>AGRICULTURE SECTOR</b>			
CH <sub>4</sub> Emissions from Enteric Fermentation in Domestic Livestock	CH <sub>4</sub>	Yes	Level, Trend
CH <sub>4</sub> Emissions from Manure Management	CH <sub>4</sub>	No	
CH <sub>4</sub> and N <sub>2</sub> O Emissions from Agricultural Residue Burning	CH <sub>4</sub>	No	
N <sub>2</sub> O Emissions from Manure Management	N <sub>2</sub> O	Yes	Level
Direct N <sub>2</sub> O Emissions from Agricultural Soils and Animals	N <sub>2</sub> O	Yes	Level
N <sub>2</sub> O Emissions from Pasture Range and Paddock Manure	N <sub>2</sub> O	Yes	Level
Indirect N <sub>2</sub> O Emissions from Nitrogen Used in Agriculture	N <sub>2</sub> O	Yes	Level
CH <sub>4</sub> and N <sub>2</sub> O Emissions from Agricultural Residue Burning	N <sub>2</sub> O	No	
<b>WASTE SECTOR</b>			
CH <sub>4</sub> Emissions from Solid Waste Disposal Sites	CH <sub>4</sub>	Yes	Level, Trend
N <sub>2</sub> O Emissions from Human Sewage	N <sub>2</sub> O	No	



Table A1-7: Key categories for Croatia – summary (Including LULUCF)

Tier 1 Analysis – Source Analysis Summary (Croatian Inventory)			
IPCC Source Categories	Direct GHG	Key Source Category Flag	Criteria for Identification
<b>ENERGY SECTOR</b>			
CO <sub>2</sub> Emissions from Stationary Combustion - Coal	CO <sub>2</sub>	Yes	Level
CO <sub>2</sub> Emissions from Stationary Combustion - Oil	CO <sub>2</sub>	Yes	Level, Trend
CO <sub>2</sub> Emissions from Stationary Combustion - Gas	CO <sub>2</sub>	Yes	Level, Trend
Non-CO <sub>2</sub> Emissions from Stationary Combustion	CH <sub>4</sub>	No	
Non-CO <sub>2</sub> Emissions from Stationary Combustion	N <sub>2</sub> O	No	
Mobile Combustion – Road Vehicles	CO <sub>2</sub>	Yes	Level, Trend
Mobile Combustion - Railways	CO <sub>2</sub>	No	
Mobile Combustion - Domestic Aviation	CO <sub>2</sub>	Yes	
Mobile Combustion - National Navigation	CO <sub>2</sub>	No	
Mobile Combustion - Agriculture/Forestry/Fishing	CO <sub>2</sub>	Yes	Level
Mobile Combustion – Road Vehicles	CH <sub>4</sub>	No	
Mobile Combustion - Railways	CH <sub>4</sub>	No	
Mobile Combustion - Domestic Aviation	CH <sub>4</sub>	No	
Mobile Combustion - National Navigation	CH <sub>4</sub>	No	
Mobile Combustion - Agriculture/Forestry/Fishing	CH <sub>4</sub>	No	
Mobile Combustion – Road Vehicles	N <sub>2</sub> O	Yes	Trend
Mobile Combustion - Railways	N <sub>2</sub> O	No	
Mobile Combustion - Domestic Aviation	N <sub>2</sub> O	No	
Mobile Combustion - National Navigation	N <sub>2</sub> O	No	
Mobile Combustion - Agriculture/Forestry/Fishing	N <sub>2</sub> O	No	
Fugitive Emissions from Coal Mining and Handling	CH <sub>4</sub>	No	
Fugitive Emissions from Oil and Gas Operations	CH <sub>4</sub>	Yes	Level, Trend
CO <sub>2</sub> Emissions from Natural Gas Scrubbing	CO <sub>2</sub>	Yes	Level, Trend
<b>CO<sub>2</sub> EMISSIONS FROM SOLVENT AND OTHER PRODUCT USE</b>	CO <sub>2</sub>	Yes	Trend
<b>INDUSTRIAL SECTOR</b>			
CO <sub>2</sub> Emissions from Cement Production	CO <sub>2</sub>	Yes	Level, Trend
CO <sub>2</sub> Emissions from Lime Production	CO <sub>2</sub>	No	
CO <sub>2</sub> Emissions from Limestone and Dolomite Use	CO <sub>2</sub>	No	
CO <sub>2</sub> Emissions from Soda Ash Production and Use	CO <sub>2</sub>	No	
CO <sub>2</sub> Emissions from Ammonia Production	CO <sub>2</sub>	Yes	Level
CO <sub>2</sub> Emissions from Iron and Steel Production	CO <sub>2</sub>	No	
CO <sub>2</sub> Emissions from Ferroalloys Production	CO <sub>2</sub>	Yes	Trend
CO <sub>2</sub> Emissions from Aluminium Production	CO <sub>2</sub>	No	
CH <sub>4</sub> Emissions from Production of Other Chemicals	CH <sub>4</sub>	No	
N <sub>2</sub> O Emissions from Nitric Acid Production	N <sub>2</sub> O	Yes	Level
HFC Emissions from Consumption of HFCs	HFC	Yes	Level, Trend
PFC Emissions from Aluminium production	PFC	No	Trend
CO <sub>2</sub> Emissions from Other non-specified NEU	CO <sub>2</sub>	No	
<b>AGRICULTURE SECTOR</b>			
CH <sub>4</sub> Emissions from Enteric Fermentation in Domestic Livestock	CH <sub>4</sub>	Yes	Level, Trend
CH <sub>4</sub> Emissions from Manure Management	CH <sub>4</sub>	No	
CH <sub>4</sub> and N <sub>2</sub> O Emissions from Agricultural Residue Burning	CH <sub>4</sub>	No	
N <sub>2</sub> O Emissions from Manure Management	N <sub>2</sub> O	No	
Direct N <sub>2</sub> O Emissions from Agricultural Soils and Animals	N <sub>2</sub> O	Yes	Level
N <sub>2</sub> O Emissions from Pasture Range and Paddock Manure	N <sub>2</sub> O	Yes	Level
Indirect N <sub>2</sub> O Emissions from Nitrogen Used in Agriculture	N <sub>2</sub> O	Yes	Level
CH <sub>4</sub> and N <sub>2</sub> O Emissions from Agricultural Residue Burning	N <sub>2</sub> O	No	
<b>LULUCF</b>			
Forest land remaining forest land	CO <sub>2</sub>	Yes	Level, Trend
<b>WASTE SECTOR</b>			
CH <sub>4</sub> Emissions from Solid Waste Disposal Sites	CH <sub>4</sub>	Yes	Level, Trend
N <sub>2</sub> O Emissions from Human Sewage	N <sub>2</sub> O	No	

Table A1-8: Changes in Key categories for Croatia based on the Level and Trend of Emissions

Tier 1 Analysis – Source Analysis Summary (Croatian Inventory)					
IPCC Source Categories	Direct GHG	Criteria for Identification			
		Level		Trend	
		2004	2005	2004	2005
<b>ENERGY SECTOR</b>					
CO <sub>2</sub> Emissions from Stationary Combustion: Coal	CO <sub>2</sub>	Yes	Yes	Yes	Yes
CO <sub>2</sub> Emissions from Stationary Combustion: Oil	CO <sub>2</sub>	Yes	Yes	Yes	Yes
CO <sub>2</sub> Emissions from Stationary Combustion: Gas	CO <sub>2</sub>	Yes	Yes	Yes	Yes
Mobile Combustion: Road Vehicles	CO <sub>2</sub>	Yes	Yes	Yes	Yes
Mobile Combustion: Domestic Aviation	CO <sub>2</sub>	No	No	No	No
Mobile Combustion: Agriculture/Forestry/Fishing	CO <sub>2</sub>	Yes	Yes	No	Yes
Mobile Combustion: Road Vehicles	N <sub>2</sub> O	No	No	Yes	Yes
Fugitive Emissions from Oil and Gas Operations	CH <sub>4</sub>	Yes	Yes	Yes	Yes
CO <sub>2</sub> Emissions from Natural Gas Scrubbing	CO <sub>2</sub>	Yes	Yes	Yes	Yes
<b>INDUSTRIAL SECTOR</b>					
CO <sub>2</sub> Emissions from Cement Production	CO <sub>2</sub>	Yes	Yes	Yes	Yes
CO <sub>2</sub> Emissions from Ammonia Production	CO <sub>2</sub>	Yes	Yes	No	No
CO <sub>2</sub> Emissions from Ferroalloys Production	CO <sub>2</sub>	No	No	Yes	Yes
N <sub>2</sub> O Emissions from Nitric Acid Production	N <sub>2</sub> O	Yes	Yes	No	Yes
PFC Emissions from Aluminium Production	PFC	No	No	Yes	Yes
CO <sub>2</sub> Emissions from Aluminium Production	CO <sub>2</sub>	No	No	Yes*	Yes
HFC Emissions from Consumption of HFCs	HFC	No	Yes	No	Yes
CO <sub>2</sub> Emissions from Other non-specified NEU	CO <sub>2</sub>	Not included in calculation	No	Not included in calculation	No
<b>CO<sub>2</sub> EMISSIONS FROM SOLVENT AND OTHER PRODUCT USE</b>	CO <sub>2</sub>	Not estimated	No	Not estimated	Yes
<b>AGRICULTURE SECTOR</b>					
CH <sub>4</sub> Emissions from Enteric Fermentation in Domestic Livestock	CH <sub>4</sub>	Yes	Yes	Yes	Yes
N <sub>2</sub> O Emissions from Manure Management	N <sub>2</sub> O	Yes	Yes	No	No
Direct N <sub>2</sub> O Emissions from Agricultural Soils and Animals	N <sub>2</sub> O	Yes	Yes	Yes	No
N <sub>2</sub> O Emissions from Pasture, Range and Paddock Manure	N <sub>2</sub> O	Yes*	Yes	No	No
Indirect N <sub>2</sub> O Emissions from Nitrogen Used in Agriculture	N <sub>2</sub> O	Yes	Yes	No	No
<b>LULUCF</b>					
Forest Land Remaining Forest Land	CO <sub>2</sub>	Yes	Yes	Yes	Yes
<b>WASTE SECTOR</b>					
CH <sub>4</sub> Emissions from Solid Waste Disposal Sites	CH <sub>4</sub>	Yes	Yes	Yes	Yes

\* Excluded in submission 2006 due to an error

## **ANNEX 2**

### **DETAILED DISCUSSION OF ACTIVITY DATA AND EMISSION FACTORS FOR ESTIMATING CO<sub>2</sub> EMISSIONS FROM FOSSIL FUEL COMBUSTION**

Table A2-1: The GHG emissions from Thermal Power Plants

	1990	1995	2000	2001	2002	2003	2004	2005
<b>Production</b>								
Electricity product. (GWh)	3760.2	1901.1	3049.4	3530.9	4627.9	4750.1	3326.4	3486.4
<b>Fuel consumption</b>								
Hard coal (1000 t)	253.7	96.2	569.8	627.3	800.4	904.2	852.4	887.5
NCV for hard coal (MJ/kg)	25.1	25.1	26.2	25.6	25.6	24.4	23.8	25.1
Coke gas (1000000 m <sup>3</sup> )	24.5							
NCV for coke gas (MJ/m <sup>3</sup> )	17.9							
Fuel oil (1000 t)	570.7	327.8	283.4	397.5	406.8	559.5	246.5	279.2
NCV for fuel oil (MJ/kg)	40.2	40.2	40.2	40.7	40.8	40.4	40.4	40.4
Extra light oil (1000 t)		24.1	0.2			8.6	0.04	2.9
NCV for ex. light oil (MJ/kg)		42.7	42.7			41.8	41.8	42.3
Natural gas (1000000 m <sup>3</sup> )	201.7	114.1	155.8	165.2	318.8	99.9	130.4	36.5
NCV for nat. gas (MJ/m <sup>3</sup> )	34.0	34.0	34.0	33.3	33.3	33.3	33.3	34.1
<b>Total fuel consumpt. (TJ)</b>	<b>36606.0</b>	<b>20501.5</b>	<b>31595.9</b>	<b>37748.4</b>	<b>47739.7</b>	<b>48355.8</b>	<b>34562.1</b>	<b>34914.5</b>
<b>Emissions</b>								
EF CO <sub>2</sub> – hard coal (t/TJ)	92.7	92.7	92.7	92.7	92.7	92.7	92.7	92.7
EF CO <sub>2</sub> – coke gas (t/TJ)	46.7	46.7	46.7	46.7	46.7	46.7	46.7	46.7
EF CO <sub>2</sub> – fuel oil (t/TJ)	76.6	76.6	76.6	76.6	76.6	76.6	76.6	76.6
EF CO <sub>2</sub> – extra light oil (t/TJ)	73.3	73.3	73.3	73.3	73.3	73.3	73.3	73.3
EF CO <sub>2</sub> – natural gas (t/TJ)	55.8	55.8	55.8	55.8	55.8	55.8	55.8	55.8
<b>CO<sub>2</sub> emission (Gg)</b>	<b>2750.9</b>	<b>1525.3</b>	<b>2550.1</b>	<b>3035.6</b>	<b>3766.5</b>	<b>3988.8</b>	<b>2883.3</b>	<b>3006.9</b>
EF CH <sub>4</sub> – hard coal (kg/TJ)	1.0	1.0	1.0	0.7	0.7	0.7	0.7	0.7
EF CH <sub>4</sub> – coke gas (kg/TJ)	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
EF CH <sub>4</sub> – fuel oil (kg/TJ)	3.0	3.0	3.0	0.9	0.9	0.9	0.9	0.9
EF CH <sub>4</sub> – extra light oil (kg/TJ)	3.0	3.0	3.0	0.9	0.9	0.9	0.9	0.9
EF CH <sub>4</sub> – natural gas (kg/TJ)	1.0	1.0	1.0	0.3	0.9	0.8	0.3	0.3
<b>CH<sub>4</sub> emission (Mg)</b>	<b>82.5</b>	<b>48.9</b>	<b>54.4</b>	<b>27.3</b>	<b>38.6</b>	<b>38.7</b>	<b>24.5</b>	<b>26.0</b>
EF N <sub>2</sub> O – hard coal (kg/TJ)	1.4	1.4	1.4	1.6	1.6	1.6	1.6	1.6
EF N <sub>2</sub> O – coke gas (kg/TJ)	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4
EF N <sub>2</sub> O – fuel oil (kg/TJ)	0.6	0.6	0.6	0.3	0.3	0.3	0.3	0.3
EF N <sub>2</sub> O – extra light oil (kg/TJ)	0.6	0.6	0.6	0.4	0.4	0.4	0.4	0.4
EF N <sub>2</sub> O – natural gas (kg/TJ)	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
<b>N<sub>2</sub>O emission (Mg)</b>	<b>24.0</b>	<b>12.3</b>	<b>28.2</b>	<b>31.1</b>	<b>38.9</b>	<b>42.5</b>	<b>35.8</b>	<b>39.2</b>

Table A2-2: The GHG emissions from Public Cogeneration Plants

	1990	1995	2000	2001	2002	2003	2004	2005
<b>Production</b>								
Electricity production (GWh)	555.6	778.0	836.1	1184.0	1273.7	1952.7	2075.3	1821.6
Heat production (TJ)	11741.3	10446.5	7024.6	8672.1	8668.0	8116.1	7789.5	8764.5
<b>Total (TJ)</b>	<b>13741.5</b>	<b>13247.2</b>	<b>10034.6</b>	<b>12934.5</b>	<b>13253.2</b>	<b>15146.0</b>	<b>15260.4</b>	<b>15322.1</b>
<b>Fuel consumption</b>								
Fuel oil (1000 t)	118.0	337.1	108.6	115.4	92.5	166.2	114.6	162.0
NCV for fuel oil (MJ/kg)	40.2	40.2	40.2	40.4	40.3	40.6	40.6	40.3
Extra light oil (1000 t)		0.9	0.9				0.3	1.5
NCV for extra light oil (MJ/kg)		42.7	42.7				41.8	42.4
Natural gas (1000000 m <sup>3</sup> )	315.5	103.5	363.4	433.1	455.9	521.0	580.1	459.5
NCV for natural gas (MJ/m <sup>3</sup> )	34.0	34.0	34.0	33.3	33.3	33.3	33.3	33.4
<b>Total fuel consumption (TJ)</b>	<b>15469.4</b>	<b>17105.5</b>	<b>16758.7</b>	<b>19101.6</b>	<b>18925.8</b>	<b>24126.6</b>	<b>24007.5</b>	<b>21918.0</b>

Table A2-2: The GHG emissions from Public Cogeneration Plants (cont.)

	1990	1995	2000	2001	2002	2003	2004	2005
<b>Emissions</b>								
EF CO <sub>2</sub> – fuel oil (t/TJ)	76.6	76.6	76.6	76.6	76.6	76.6	76.6	76.6
EF CO <sub>2</sub> – ex.light oil (t/TJ)	73.3	73.3	73.3	73.3	73.3	73.3	73.3	73.3
EF CO <sub>2</sub> – natural gas (t/TJ)	55.8	55.8	55.8	55.8	55.8	55.8	55.8	55.8
<b>CO<sub>2</sub> emission (Gg)</b>	<b>962.0</b>	<b>1236.9</b>	<b>1026.8</b>	<b>1163.1</b>	<b>1133.8</b>	<b>1487.1</b>	<b>1437.0</b>	<b>1360.2</b>
EF CH <sub>4</sub> – fuel oil (kg/TJ)	3.0	3.0	3.0	0.9	0.9	0.9	0.9	0.9
EF CH <sub>4</sub> – ex.light oil (kg/TJ)	3.0	3.0	3.0	0.9	0.9	0.9	0.9	0.9
EF CH <sub>4</sub> – nat. gas (kg/TJ)	1.0	1.0	1.0	3.6	3.9	4.7	4.6	4.6
<b>CH<sub>4</sub> emission (Mg)</b>	<b>25.0</b>	<b>44.3</b>	<b>25.6</b>	<b>56.1</b>	<b>62.0</b>	<b>86.9</b>	<b>93.1</b>	<b>81.8</b>
EF N <sub>2</sub> O – fuel oil (kg/TJ)	0.6	0.6	0.6	0.3	0.3	0.3	0.3	0.3
EF N <sub>2</sub> O – ex.light oil (kg/TJ)	0.6	0.6	0.6	0.4	0.4	0.4	0.4	0.4
EF N <sub>2</sub> O – nat. gas (kg/TJ)	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
<b>N<sub>2</sub>O emission (Mg)</b>	<b>3.9</b>	<b>8.5</b>	<b>3.9</b>	<b>2.8</b>	<b>2.6</b>	<b>3.8</b>	<b>3.3</b>	<b>3.5</b>

Table A2-3: The GHG emissions from Public Heating Plants

	1990	1995	2000	2001	2002	2003	2004	2005
<b>Production</b>								
Heat production (TJ)	0	2493	2708	3338.4	3171.4	3469.7	3303.5	3477.7
<b>Fuel consumption</b>								
Fuel oil (1000 t)	0.0	35.6	37.0	38.6	36.4	38.1	38.6	39
NCV for fuel oil (MJ/kg)	40.2	40.2	40.2	40.2	40.2	40.2	40.2	40.2
Extra light oil (1000 t)	0.0	6.0	4.4	3.6	3.7	4.3	6.8	6.7
NCV for extra light oil (MJ/kg)	42.7	42.7	42.7	42.7	42.7	42.7	42.7	42.7
Natural gas (1000000 m <sup>3</sup> )	0.0	36.2	53.0	70.1	67.2	69.6	64.3	71.3
NCV for natural gas (MJ/m <sup>3</sup> )	34.0	34.0	34.0	34.0	34.0	34.0	34.0	34.0
<b>Total fuel consumption (TJ)</b>	<b>0.0</b>	<b>2917.8</b>	<b>3477.0</b>	<b>4088.5</b>	<b>3905.7</b>	<b>4081.3</b>	<b>4028.0</b>	<b>4277.8</b>
<b>Emissions</b>								
EF CO <sub>2</sub> – fuel oil (t/TJ)	76.6	76.6	76.6	76.6	76.6	76.6	76.6	76.6
EF CO <sub>2</sub> – extra light oil (t/TJ)	73.3	73.3	73.3	73.3	73.3	73.3	73.3	73.3
EF CO <sub>2</sub> – natural gas (t/TJ)	55.8	55.8	55.8	55.8	55.8	55.8	55.8	55.8
<b>CO<sub>2</sub> emission (Gg)</b>	<b>0.0</b>	<b>197.1</b>	<b>228.3</b>	<b>263.1</b>	<b>251.2</b>	<b>262.8</b>	<b>262.1</b>	<b>276.4</b>
EF CH <sub>4</sub> – fuel oil (kg/TJ)	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0
EF CH <sub>4</sub> – extra light oil (kg/TJ)	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0
EF CH <sub>4</sub> – natural gas (kg/TJ)	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
<b>CH<sub>4</sub> emission (Mg)</b>	<b>0.0</b>	<b>6.3</b>	<b>6.8</b>	<b>7.5</b>	<b>7.1</b>	<b>7.5</b>	<b>7.7</b>	<b>8.0</b>
EF N <sub>2</sub> O – fuel oil (kg/TJ)	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
EF N <sub>2</sub> O – extra light oil (kg/TJ)	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
EF N <sub>2</sub> O – natural gas (kg/TJ)	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
<b>N<sub>2</sub>O emission (Mg)</b>	<b>0.0</b>	<b>1.1</b>	<b>1.2</b>	<b>1.3</b>	<b>1.2</b>	<b>1.3</b>	<b>1.3</b>	<b>1.4</b>

The GHG emissions from thermal power plants and public cogeneration plants, for last five years (2001, 2002, 2003, 2004 and 2005), were calculated using more detailed Tier 2 approach. Tier 2 approach is based on bottom-up fuel consumption data from every boiler or gas turbine in plant. There were available data about monthly fuel consumption and detailed fuel characteristics data (net calorific value, sulphur and ash content...). Every plant also has the equipment for continual measurements of SO<sub>2</sub>, NO<sub>x</sub>, CO and particulates emission.

For estimation of CO<sub>2</sub> emissions, default IPCC emission factors were used, while emission factors for CH<sub>4</sub> and N<sub>2</sub>O are based on technology type and configuration (Tier 2). The results of GHG emission calculation, using more detailed approach are presented in tables A2-2 and A2-3 for the last five years, on aggregated level. The GHG emissions on plant level, for the year 2005, are given in the Table A2-5.

*Table A2-4: The GHG emissions from TPPs and PCPs (Tier 2), year 2005*

	TPP Plomin	TPP Rijeka	TPP Sisak	CHP Zagreb – east	CHP Zagreb – west	CHP Osijek	CCGT Jertovec
<b>Production</b>							
Electricity production (GWh)	2265.3	697.2	523.1	1389.0	325.9	106.7	0.8
Heat production (TJ)				4010.4	3512.8	1241.3	
<b>Total (TJ)</b>	<b>8155.1</b>	<b>2509.9</b>	<b>1883.2</b>	<b>9010.9</b>	<b>4685.9</b>	<b>1625.3</b>	<b>3.0</b>
<b>Fuel consumption</b>							
Hard coal (1000 t)	887.5						
NCV for hard coal (MJ/kg)	25.1						
Fuel oil (1000 t)		173.2	105.9	69.2	45.8	47.0	
NCV for fuel oil (MJ/kg)		40.40	40.34	40.19	40.00	40.81	
Extra light oil (1000 t)	1.72	0.9		1.5		0.0	0.29
NCV for extra light oil (MJ/kg)	42.0	42.1		42.0		42.9	42.7
Natural gas (1000000 m <sup>3</sup> )			36.5	324.3	117.2	18.0	
NCV for natural gas (MJ/m <sup>3</sup> )			34.1	33.3	33.3	33.7	
<b>Total fuel consumption (TJ)</b>	<b>22348.5</b>	<b>7034.8</b>	<b>5518.8</b>	<b>13656.3</b>	<b>5737.5</b>	<b>2524.1</b>	<b>12.4</b>
<b>Emissions</b>							
EF CO <sub>2</sub> – hard coal (t/TJ)	92.7	92.7	92.7	92.7	92.7	92.7	92.7
EF CO <sub>2</sub> – fuel oil (t/TJ)	76.6	76.6	76.6	76.6	76.6	76.6	76.6
EF CO <sub>2</sub> – extra light oil (t/TJ)	73.3	73.3	73.3	73.3	73.3	73.3	73.3
EF CO <sub>2</sub> – natural gas (t/TJ)	55.8	55.8	55.8	55.8	55.8	55.8	55.8
<b>CO<sub>2</sub> emission (Gg)</b>	<b>2070.5</b>	<b>538.7</b>	<b>396.8</b>	<b>821.2</b>	<b>358.3</b>	<b>180.8</b>	<b>0.9</b>
EF CH <sub>4</sub> – hard coal (kg/TJ)	0.7	0.7	0.7	0.7	0.7	0.7	0.7
EF CH <sub>4</sub> – fuel oil (kg/TJ)	0.9	0.9	0.9	0.9	0.9	0.9	0.9
EF CH <sub>4</sub> – extra light oil (kg/TJ)	0.9	0.9	0.9	0.9	0.9	0.9	0.9
EF CH <sub>4</sub> – natural gas (kg/TJ)	0.1	0.1	0.1	5.2	4.9	0.4	6.0
<b>CH<sub>4</sub> emission (Mg)</b>	<b>15.7</b>	<b>6.3</b>	<b>4.0</b>	<b>58.4</b>	<b>20.7</b>	<b>1.9</b>	<b>0.0</b>
EF N <sub>2</sub> O – hard coal (kg/TJ)	1.6	1.6	1.6	1.6	1.6	1.6	1.6
EF N <sub>2</sub> O – fuel oil (kg/TJ)	0.3	0.3	0.3	0.3	0.3	0.3	0.3
EF N <sub>2</sub> O – extra light oil (kg/TJ)	0.4	0.4	0.4	0.4	0.4	0.4	0.4
EF N <sub>2</sub> O – natural gas (kg/TJ)	0.1	0.1	0.1	0.1	0.1	0.1	0.1
<b>N<sub>2</sub>O emission (Mg)</b>	<b>35.7</b>	<b>2.1</b>	<b>1.4</b>	<b>1.9</b>	<b>0.9</b>	<b>0.6</b>	<b>0.0</b>

Table A2-5: The GHG emissions from Petroleum refining – own use of energy

	1990	1995	2000	2001	2002	2003	2004	2005
<b>Fuel consumption</b>								
Fuel oil (1000 t)	127.7	101.2	47.8	36.8	57.4	51.7	66.2	70.80
NCV for fuel oil (MJ/kg)	40.2	40.2	40.2	40.2	40.2	40.2	40.2	40.19
LPG (1000 t)		35.0	2.2	4.3	7.7	8.8	5.7	6.60
NCV for LPG (MJ/kg)	46.9	46.9	46.9	46.9	46.9	46.9	46.9	46.89
Petroleum coke (1000 t)	53.7	42.6	63.0	51.5	57.8	59.1	63.4	64.22
NCV for petroleum coke (MJ/kg)	29.3	29.3	31.0	31.0	31.0	31.0	31.0	31.00
Refinery gas (1000 t)	347.5	196.5	221.7	195.3	214.5	229.9	241.7	222.00
NCV for refinery gas (MJ/kg)	48.6	48.6	48.6	48.6	48.6	48.6	48.6	48.57
Natural gas (1000000 m <sup>3</sup> )				3.3	0.3	0.3	0.3	1.20
NCV for natural gas (MJ/m <sup>3</sup> )	34.0	34.0	34.0	34.0	34.0	34.0	34.0	34.00
<b>Total fuel consumption (TJ)</b>	<b>23584.3</b>	<b>16501.0</b>	<b>14745.2</b>	<b>12875.0</b>	<b>14888.2</b>	<b>15499.0</b>	<b>16642.8</b>	<b>15969.1</b>
<b>Emissions</b>								
EF CO <sub>2</sub> – fuel oil (t/TJ)	76.6	76.6	76.6	76.6	76.6	76.6	76.6	76.6
EF CO <sub>2</sub> – LPG (t/TJ)	62.4	62.4	62.4	62.4	62.4	62.4	62.4	62.4
EF CO <sub>2</sub> – petroleum coke (t/TJ)	99.8	99.8	99.8	99.8	99.8	99.8	99.8	99.8
EF CO <sub>2</sub> – refinery gas (t/TJ)	66.1	66.1	66.1	66.1	66.1	66.1	66.1	66.1
EF CO <sub>2</sub> – natural gas (t/TJ)	55.8	55.8	55.8	55.8	55.8	55.8	55.8	55.8
<b>CO<sub>2</sub> emission (Gg)</b>	<b>1665.3</b>	<b>1169.2</b>	<b>1059.9</b>	<b>918.2</b>	<b>1067.0</b>	<b>1106.1</b>	<b>1192.8</b>	<b>1150.6</b>
EF CH <sub>4</sub> – fuel oil (kg/TJ)	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0
EF CH <sub>4</sub> – LPG (kg/TJ)	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0
EF CH <sub>4</sub> – petroleum coke (kg/TJ)	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0
EF CH <sub>4</sub> – refinery gas (kg/TJ)	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0
EF CH <sub>4</sub> – natural gas (kg/TJ)	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
<b>CH<sub>4</sub> emission (Mg)</b>	<b>70.8</b>	<b>49.5</b>	<b>44.2</b>	<b>38.4</b>	<b>44.6</b>	<b>46.5</b>	<b>49.9</b>	<b>47.8</b>
EF N <sub>2</sub> O – fuel oil (kg/TJ)	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
EF N <sub>2</sub> O – LPG (kg/TJ)	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
EF N <sub>2</sub> O – petroleum coke (kg/TJ)	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
EF N <sub>2</sub> O – refinery gas (kg/TJ)	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
EF N <sub>2</sub> O – natural gas (kg/TJ)	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
<b>N<sub>2</sub>O emission (Mg)</b>	<b>14.2</b>	<b>9.9</b>	<b>8.8</b>	<b>7.7</b>	<b>8.9</b>	<b>9.3</b>	<b>10.0</b>	<b>9.6</b>

Table A2-6: The GHG emissions from Petroleum refining – heating/cogeneration plants\*

	1990	1995	2000	2001	2002	2003	2004	2005
<b>Fuel consumption</b>								
Fuel oil (1000 t)	227.2	199.5	193.4	183.8	205.9	212.9	192.6	183.2
NCV for fuel oil (MJ/kg)	40.2	40.2	40.2	40.2	40.2	40.2	40.2	40.19
LPG (1000 t)				6.5			11.6	2.9
NCV for LPG (MJ/kg)	46.9	46.9	46.9	46.9	46.9	46.9	46.9	46.89
Petroleum coke (1000 t)				4.1	6.3	6.9	6.2	6.5
NCV for petroleum coke (MJ/kg)				31.0	31.0	31.0	31.0	31
Refinery gas (1000 t)	58.4	27.7	40.7	25.3	22.7	28.6	22.1	19.1
NCV for refinery gas (MJ/kg)	48.6	48.6	48.6	48.6	48.6	48.6	48.6	48.57
Natural gas (1000000 m <sup>3</sup> )	7.3	7.1	0.2	0.6				
NCV for natural gas (MJ/m <sup>3</sup> )	34.0	34.0	34.0	34.0	34.0	34.0	34.0	34
<b>Total fuel consumption (TJ)</b>	<b>12215.9</b>	<b>9604.7</b>	<b>9756.3</b>	<b>9068.0</b>	<b>9573.0</b>	<b>10159.5</b>	<b>9550.1</b>	<b>8628.0</b>

Table A2-6: The GHG emissions from Petroleum refining – heating/cogeneration plants\* (cont.)

	1990	1995	2000	2001	2002	2003	2004	2005
<b>Emissions</b>								
EF CO <sub>2</sub> – fuel oil (t/TJ)	76.6	76.6	76.6	76.6	76.6	76.6	76.6	76.6
EF CO <sub>2</sub> – LPG (t/TJ)	62.4	62.4	62.4	62.4	62.4	62.4	62.4	62.4
EF CO <sub>2</sub> – petroleum coke (t/TJ)	99.8	99.8	99.8	99.8	99.8	99.8	99.8	99.8
EF CO <sub>2</sub> – refinery gas (t/TJ)	66.1	66.1	66.1	66.1	66.1	66.1	66.1	66.1
EF CO <sub>2</sub> – natural gas (t/TJ)	55.8	55.8	55.8	55.8	55.8	55.8	55.8	55.8
<b>CO<sub>2</sub> emission (Gg)</b>	<b>900.6</b>	<b>716.5</b>	<b>726.3</b>	<b>679.8</b>	<b>726.2</b>	<b>768.5</b>	<b>716.9</b>	<b>653.8</b>
EF CH <sub>4</sub> – fuel oil (kg/TJ)	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0
EF CH <sub>4</sub> – LPG (kg/TJ)	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0
EF CH <sub>4</sub> – petroleum coke (kg/TJ)	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0
EF CH <sub>4</sub> – refinery gas (kg/TJ)	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0
EF CH <sub>4</sub> – natural gas (kg/TJ)	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
<b>CH<sub>4</sub> emission (Mg)</b>	<b>36.2</b>	<b>28.3</b>	<b>29.3</b>	<b>27.2</b>	<b>28.7</b>	<b>30.5</b>	<b>28.7</b>	<b>25.9</b>
EF N <sub>2</sub> O – fuel oil (kg/TJ)	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
EF N <sub>2</sub> O – LPG (kg/TJ)	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
EF N <sub>2</sub> O – petroleum coke (kg/TJ)	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
EF N <sub>2</sub> O – refinery gas (kg/TJ)	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
EF N <sub>2</sub> O – natural gas (kg/TJ)	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
<b>N<sub>2</sub>O emission (Mg)</b>	<b>7.2</b>	<b>5.6</b>	<b>5.9</b>	<b>5.4</b>	<b>5.7</b>	<b>6.1</b>	<b>5.7</b>	<b>5.2</b>



Table A2-7: The GHG emissions from manufacturing of solid fuels and other energy industries

	1990	1995	2000	2001	2002	2003	2004	2005
<b>Fuel consumption</b>								
LPG (1000 t)			1.0					
NCV for LPG (MJ/kg)			46.9					
Coke gas (1000000 m <sup>3</sup> )	107.4							
NCV for coke gas (MJ/m <sup>3</sup> )	17.9							
Extra light oil (1000 t)	0.7	0.8	7.1	6.3	6.4	9.5	6.3	5.5
NCV for ex.light oil (MJ/kg)	42.7	42.7	42.7	42.7	42.7	42.7	42.7	42.71
Natural gas (1000000 m <sup>3</sup> )	238.2	171	140	112.8	130.9	123.2	136.5	175.5
NCV for nat. gas (MJ/m <sup>3</sup> )	34.0	34.0	34.0	34.0	34.0	34.0	34.0	34
<b>Total fuel consumpt. (TJ)</b>	<b>10052.2</b>	<b>5848.2</b>	<b>5110.1</b>	<b>4104.3</b>	<b>4723.9</b>	<b>4594.5</b>	<b>4910.1</b>	<b>6201.9</b>
<b>Emissions</b>								
EF CO <sub>2</sub> – LPG (t/TJ)	62.4	62.4	62.4	62.4	62.4	62.4	62.4	62.4
EF CO <sub>2</sub> – coke gas (t/TJ)	46.7	46.7	46.7	46.7	46.7	46.7	46.7	46.7
EF CO <sub>2</sub> – ex.light oil (t/TJ)	73.3	73.3	73.3	73.3	73.3	73.3	73.3	73.3
EF CO <sub>2</sub> – natural gas (t/TJ)	55.8	55.8	55.8	55.8	55.8	55.8	55.8	55.8
<b>CO<sub>2</sub> emission (Gg)</b>	<b>544.1</b>	<b>327.0</b>	<b>290.9</b>	<b>233.8</b>	<b>268.5</b>	<b>263.6</b>	<b>278.8</b>	<b>350.3</b>
EF CH <sub>4</sub> – hard coal (kg/TJ)	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
EF CH <sub>4</sub> – coke gas (kg/TJ)	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
EF CH <sub>4</sub> – ex.ligh oil (kg/TJ)	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0
EF CH <sub>4</sub> – nat. gas (kg/TJ)	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
<b>CH<sub>4</sub> emission (Mg)</b>	<b>10.1</b>	<b>5.9</b>	<b>5.7</b>	<b>4.6</b>	<b>5.3</b>	<b>5.4</b>	<b>5.4</b>	<b>6.7</b>
EF N <sub>2</sub> O – hard coal (kg/TJ)	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4
EF N <sub>2</sub> O – coke gas (kg/TJ)	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4
EF N <sub>2</sub> O – ex.ligh oil (kg/TJ)	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
EF N <sub>2</sub> O – nat. gas (kg/TJ)	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
<b>N<sub>2</sub>O emission (Mg)</b>	<b>3.5</b>	<b>0.6</b>	<b>0.7</b>	<b>0.5</b>	<b>0.6</b>	<b>0.7</b>	<b>0.6</b>	<b>0.7</b>

Table A2-8: The GHG emissions from Manufacturing Industries and Construction – liquid fuels and natural gas

	1990	1995	2000	2001	2002	2003	2004	2005
<b>Fuel consumption</b>								
Gasoline (1000 t)	0.2	8.5	7.6	7.8	7.6	9.1	10.2	6.9
NCV for gasoline (MJ/kg)	44.6	44.6	44.6	44.6	44.6	44.6	44.6	44.6
Petroleum (1000 t)	0.1							
NCV for petroleum (MJ/kg)	44.0							
Gas/diesel oil (1000 t)	246.6	101.5	130.8	129.9	119.3	145.0	135.9	110.6
NCV for gas/diesel o.(MJ/kg)	42.7	42.7	42.7	42.7	42.7	42.7	42.7	42.7
Fuel oil (1000 t)	419.1	269.7	302.2	325.5	324.4	284.7	230.5	198.6
NCV for fuel oil (MJ/kg)	40.2	40.2	40.2	40.2	40.2	40.2	40.2	40.2
LPG (1000 t)	17.6	17.6	21.0	19.8	19.3	20.9	17.3	22.8
NCV for LPG (MJ/kg)	46.9	46.9	46.9	46.9	46.9	46.9	46.9	46.9
Lubricants (1000 t)	8.6							
NCV for lubricants (MJ/kg)	33.6							
Petroleum coke (1000 t)				16.3	11.9	6.3	68.4	172.3
NCV for petr. coke (MJ/kg)				31.0	31.0	31.0	31.0	31.0
Natural gas (1000000 m <sup>3</sup> )	1099.1	1017.1	1031.5	985.7	921.7	966.0	1038.3	1040.1

*Table A2-8: The GHG emissions from Manufacturing Industries and Construction – liquid fuels and natural gas (cont.)*

	1990	1995	2000	2001	2002	2003	2004	2005
<b>Fuel consumption</b>								
NCV for natural gas (MJ/m <sup>3</sup> )	34.0	34.0	34.0	34.0	34.0	34.0	34.0	34.0
<b>Total fuel consumption (TJ)</b>	<b>65876.0</b>	<b>50963.4</b>	<b>54129.3</b>	<b>53928.6</b>	<b>51085.5</b>	<b>52063.6</b>	<b>53759.8</b>	<b>54787.8</b>
<b>Emissions</b>								
EF CO <sub>2</sub> – gasoline (t/TJ)	68.6	68.6	68.6	68.6	68.6	68.6	68.6	68.6
EF CO <sub>2</sub> – petroleum (t/TJ)	71.1	71.1	71.1	71.1	71.1	71.1	71.1	71.1
EF CO <sub>2</sub> – gas/diesel oil (t/TJ)	73.3	73.3	73.3	73.3	73.3	73.3	73.3	73.3
EF CO <sub>2</sub> – fuel oil (t/TJ)	76.6	76.6	76.6	76.6	76.6	76.6	76.6	76.6
EF CO <sub>2</sub> – LPG (t/TJ)	62.4	62.4	62.4	62.4	62.4	62.4	62.4	62.4
EF CO <sub>2</sub> – lubricants (t/TJ)	72.6	72.6	72.6	72.6	72.6	72.6	72.6	72.6
EF CO <sub>2</sub> – petroleum coke (t/TJ)	99.8	99.8	99.8	99.8	99.8	99.8	99.8	99.8
EF CO <sub>2</sub> – natural gas (t/TJ)	55.8	55.8	55.8	55.8	55.8	55.8	55.8	55.8
<b>CO<sub>2</sub> emission (Gg)</b>	<b>4111.1</b>	<b>3060.7</b>	<b>3182.6</b>	<b>3321.1</b>	<b>3153.4</b>	<b>3181.7</b>	<b>3302.7</b>	<b>3571.2</b>
EF CH <sub>4</sub> – gasoline (kg/TJ)	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0
EF CH <sub>4</sub> – petroleum (kg/TJ)	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0
EF CH <sub>4</sub> – gas/diesel oil (kg/TJ)	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0
EF CH <sub>4</sub> – fuel oil (kg/TJ)	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0
EF CH <sub>4</sub> – LPG (kg/TJ)	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0
EF CH <sub>4</sub> – lubricants (kg/TJ)	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0
EF CH <sub>4</sub> – petroleum coke (kg/TJ)	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0
EF CH <sub>4</sub> – natural gas (kg/TJ)	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0
<b>CH<sub>4</sub> emission (Mg)</b>	<b>236</b>	<b>199</b>	<b>205</b>	<b>202</b>	<b>190</b>	<b>196</b>	<b>207</b>	<b>212</b>
EF N <sub>2</sub> O – gasoline (kg/TJ)	0.6	0.6	0.6	0.6	0.6	0.6	0.6	
EF N <sub>2</sub> O – petroleum (kg/TJ)	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
EF N <sub>2</sub> O – gas/diesel oil (kg/TJ)	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
EF N <sub>2</sub> O – fuel oil (kg/TJ)	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
EF N <sub>2</sub> O – LPG (kg/TJ)	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
EF N <sub>2</sub> O – lubricants (kg/TJ)	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
EF N <sub>2</sub> O – petroleum coke (kg/TJ)	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
EF N <sub>2</sub> O – natural gas (kg/TJ)	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
<b>N<sub>2</sub>O emission (Mg)</b>	<b>21</b>	<b>13</b>	<b>15</b>	<b>15</b>	<b>15</b>	<b>15</b>	<b>14</b>	<b>16</b>

*Table A2-9: The GHG emissions from Manufacturing Industries and Construction – solid fuels*

	1990	1995	2000	2001	2002	2003	2004	2005
<b>Fuel consumption</b>								
Anthracite (1000 t)	107.2	5.0	0.1		7.2		0.4	0.3
NCV for anthracite (MJ/kg)	29.3	29.3	29.3		29.3		29.3	29.3
Hard coal (1000 t)	42.0	41.9	53.2	68.8	70.3	125.7	253.7	169.3
NCV for hard coal (MJ/kg)	25.1	25.1	26.2	25.8	25.8	24.5	24.3	25.1
Brown Coal (1000 t)	261.2	95.8	28.2	42.3	35.5	34.4	59.2	55.9
NCV for brown coal (MJ/kg)	16.7	16.7	17.8	18.2	18.2	18.5	18.3	18.5
Lignite (1000 t)	73.0	56.3	14.4	20.2	25.6	18.4	0.7	0.2
NCV for lignite (MJ/kg)	10.9	10.9	12.0	12.2	12.2	12.3	12.2	12.1
Briquettes (1000 t)	3.3							
NCV for briquettes (MJ/kg)	16.7							
Coke oven coke (1000 t)	199.8	31.4	37.7	18.3	17.1	17.3	19.3	29.3
NCV for coke oven coke (MJ/kg)	29.3	29.3	29.3	29.3	29.3	29.3	29.3	22.8
Gas work gas (1000000 m <sup>3</sup> )	6.1	9.8	7.9	6.9	9.6	4.4	3.0	3.6

Table A2-9: The GHG emissions from Manufacturing Industries and Construction – solid fuels (cont.)

	1990	1995	2000	2001	2002	2003	2004	2005
<b>Fuel consumption</b>								
NCV for gas work gas (MJ/m <sup>3</sup> )	15.8	15.8	19.5	19.5	19.5	22.6	21.5	21.5
Coke oven gas (1000000 m <sup>3</sup> )	29.9							
NCV for coke oven gas (MJ/m <sup>3</sup> )	17.9							
Blast furnace gas (1000000 m <sup>3</sup> )	418.1							
NCV for blast furnace gas (MJ/m <sub>3</sub> )	3.6							
<b>Total fuel consumption (TJ)</b>	<b>17416.7</b>	<b>4492.5</b>	<b>3327.8</b>	<b>3461.2</b>	<b>3670.2</b>	<b>4547.6</b>	<b>7897.9</b>	<b>6040.2</b>
<b>Emissions</b>								
EF CO <sub>2</sub> – anthracite (t/TJ)	96.3	96.3	96.3	96.3	96.3	96.3	96.3	96.3
EF CO <sub>2</sub> – hard coal (t/TJ)	92.7	92.7	92.7	92.7	92.7	92.7	92.7	92.7
EF CO <sub>2</sub> – brown coal (t/TJ)	94.1	94.1	94.1	94.1	94.1	94.1	94.1	94.1
EF CO <sub>2</sub> – lignite (t/TJ)	99.2	99.2	99.2	99.2	99.2	99.2	99.2	99.2
EF CO <sub>2</sub> – briquettes (t/TJ)	92.7	92.7	92.7	92.7	92.7	92.7	92.7	92.7
EF CO <sub>2</sub> – coke oven coke (t/TJ)	106.0	106.0	106.0	106.0	106.0	106.0	106.0	106.0
EF CO <sub>2</sub> – gas work gas (t/TJ)	46.7	46.7	46.7	46.7	46.7	46.7	46.7	46.7
EF CO <sub>2</sub> – coke oven gas (t/TJ)	46.7	46.7	46.7	46.7	46.7	46.7	46.7	46.7
EF CO <sub>2</sub> – blast furnace gas (t/TJ)	237.2	237.2	237.2	237.2	237.2	237.2	237.2	237.2
<b>CO<sub>2</sub> emission (Gg)</b>	<b>1904.2</b>	<b>428.4</b>	<b>318.0</b>	<b>324.5</b>	<b>342.0</b>	<b>426.2</b>	<b>738.4</b>	<b>453.8</b>
EF CH <sub>4</sub> – anthracite (kg/TJ)	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0
EF CH <sub>4</sub> – hard coal (kg/TJ)	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0
EF CH <sub>4</sub> – brown coal (kg/TJ)	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0
EF CH <sub>4</sub> – lignite (kg/TJ)	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0
EF CH <sub>4</sub> – briquettes (kg/TJ)	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0
EF CH <sub>4</sub> – coke oven coke (kg/TJ)	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0
EF CH <sub>4</sub> – gas work gas (kg/TJ)	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0
EF CH <sub>4</sub> – coke oven gas (kg/TJ)	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0
EF CH <sub>4</sub> – blast furnace gas(kg/TJ)	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0
<b>CH<sub>4</sub> emission (Mg)</b>	<b>174.2</b>	<b>44.9</b>	<b>33.3</b>	<b>34.6</b>	<b>36.7</b>	<b>45.5</b>	<b>79.0</b>	<b>60.0</b>
EF N <sub>2</sub> O – anthracite (kg/TJ)	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4
EF N <sub>2</sub> O – hard coal (kg/TJ)	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4
EF N <sub>2</sub> O – brown coal (kg/TJ)	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4
EF N <sub>2</sub> O – lignite (kg/TJ)	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4
EF N <sub>2</sub> O – briquettes (kg/TJ)	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4
EF N <sub>2</sub> O – coke oven coke (kg/TJ)	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4
EF N <sub>2</sub> O – gas work gas (kg/TJ)	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4
EF N <sub>2</sub> O – coke oven gas (kg/TJ)	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4
EF N <sub>2</sub> O – blast furnace gas(kg/TJ)	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4
<b>N<sub>2</sub>O emission (Mg)</b>	<b>24.4</b>	<b>6.3</b>	<b>4.7</b>	<b>4.8</b>	<b>5.1</b>	<b>6.4</b>	<b>11.1</b>	<b>8.0</b>

Table A2-10: The number of road motor vehicles in Croatia

	1990	1995	2000	2001	2002	2003	2004	2005
Passenger Cars	852585	710910	1124825	1195450	1244252	1293421	1337538	1361879
Light and Heavy Duty Vehicles	70477	73497	122516	129497	138743	148275	154790	162565
Buses	6398	3897	4660	4770	4792	4833	4869	6849
Motorcycles	11847	9933	21868	24305	28188	33925	39315	76993
<b>Total</b>	<b>941307</b>	<b>798237</b>	<b>1273869</b>	<b>1354022</b>	<b>1415975</b>	<b>1480454</b>	<b>1536512</b>	<b>1608286</b>

Table A2-11: GHG emissions from Road Transport

	1990	1995	2000	2001	2002	2003	2004	2005
<b>Fuel consumption</b>								
Gasoline (1000 t)	740.0	558.0	764.2	734.9	742.8	739.6	705.2	693.5
NCV for gasoline (MJ/kg)	44.6	44.6	44.6	44.6	44.6	44.6	44.6	44.6
Diesel (1000 t)	366.0	406.0	557.8	601.0	683.6	810.9	888.1	955.6
NCV for diesel (MJ/kg)	42.7	42.7	42.7	42.7	42.7	42.7	42.7	42.7
LPG (1000 t)	0.0	13.7	9.8	12.6	13.2	13.2	16.7	22.1
NCV for LPG (MJ/kg)		46.9	46.9	46.9	46.9	46.9	46.9	46.9
<b>Total fuel consumption (TJ)</b>	<b>49575.0</b>	<b>43107.7</b>	<b>58371.7</b>	<b>59050.4</b>	<b>62945.5</b>	<b>68256.9</b>	<b>70158.7</b>	<b>72773.1</b>
<b>Emissions</b>								
<b>CO<sub>2</sub> emission (Gg)</b>	3504.0	3100.4	4218.8	4271.3	4557.8	4947.0	5092.6	5286.9
<b>CH<sub>4</sub> emission (Mg)</b>	1422.7	1130.1	1480.7	1349.5	1303.3	1243.7	1207.8	1552.9
<b>N<sub>2</sub>O emission (Mg)</b>	158.6	177.7	333.1	370.9	443.1	508.8	540.5	534.3

Table A2-12: The GHG emissions from Domestic Air Transport

	1990	1995	2000	2001	2002	2003	2004	2005
<b>Fuel consumption</b>								
Gasoline (1000 t)			0.1		0.1	0.2	1.1	1.1
NCV for gasoline (MJ/kg)			44.6		44.6	44.6	44.6	44.6
Jet kerosene (1000 t)	95	28	40	52	50	45	50	54
NCV for jet kerosene (MJ/kg)	44.0	44.0	44.0	44.0	44.0	44.0	44.0	44.0
<b>Total fuel consumption (TJ)</b>	<b>4176.2</b>	<b>1230.9</b>	<b>1762.9</b>	<b>2285.9</b>	<b>2202.5</b>	<b>1987.1</b>	<b>2233.9</b>	<b>2440.5</b>
<b>Emissions</b>								
EF CO <sub>2</sub> – gasoline (t/TJ)	68.6	68.6	68.6	68.6	68.6	68.6	68.6	68.6
EF CO <sub>2</sub> – jet kerosene (t/TJ)	70.8	70.8	70.8	70.8	70.8	70.8	70.8	70.785
<b>CO<sub>2</sub> emission (Gg)</b>	<b>295.6</b>	<b>87.1</b>	<b>124.8</b>	<b>161.8</b>	<b>155.9</b>	<b>140.6</b>	<b>159.0</b>	<b>172.6</b>
EF CH <sub>4</sub> – gasoline (kg/TJ)	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5
EF CH <sub>4</sub> – jet kerosene (kg/TJ)	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5
<b>CH<sub>4</sub> emission (Mg)</b>	<b>2.1</b>	<b>0.6</b>	<b>0.9</b>	<b>1.1</b>	<b>1.1</b>	<b>1.0</b>	<b>1.1</b>	<b>1.2</b>
EF N <sub>2</sub> O – gasoline (kg/TJ)	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0
EF N <sub>2</sub> O – jet kerosene (kg/TJ)	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0
<b>N<sub>2</sub>O emission (Mg)</b>	<b>8.4</b>	<b>2.5</b>	<b>3.5</b>	<b>4.6</b>	<b>4.4</b>	<b>4.0</b>	<b>4.5</b>	<b>4.9</b>

Table A2-13: The GHG emissions from National Navigation

	1990	1995	2000	2001	2002	2003	2004	2005
<b>Fuel consumption</b>								
Gasoline (1000 t)	0.1	0.6	0.3	0.4	0.3	0.3		
NCV for gasoline (MJ/kg)	44.6	44.6	44.6	44.6	44.6	44.6		
Diesel (1000 t)	40.3	24.7	25.7	25.6	27.9	28.6	29.1	31.8
NCV for diesel (MJ/kg)	42.7	42.7	42.7	42.7	42.7	42.7	42.7	42.7
Fuel oil (1000 t)	2.1	6.2	1.4	3.4	7.3	6.7		
NCV for fuel oil (MJ/kg)	40.2	40.2	40.2	40.2	40.2	40.2		
<b>Total fuel consumption (TJ)</b>	<b>1810.1</b>	<b>1330.9</b>	<b>1167.3</b>	<b>1247.9</b>	<b>1498.4</b>	<b>1504.2</b>	<b>1242.9</b>	<b>1358.2</b>
<b>Emissions</b>								
EF CO <sub>2</sub> – gasoline (t/TJ)	68.6	68.6	68.6	68.6	68.6	68.6	68.6	68.6
EF CO <sub>2</sub> – diesel (t/TJ)	73.3	73.3	73.3	73.3	73.3	73.3	73.3	73.3
EF CO <sub>2</sub> – fuel oil (t/TJ)	76.6	76.6	76.6	76.6	76.6	76.6	76.6	76.6
<b>CO<sub>2</sub> emission (Gg)</b>	<b>133.0</b>	<b>98.3</b>	<b>85.7</b>	<b>91.9</b>	<b>110.8</b>	<b>111.1</b>	<b>91.1</b>	<b>99.6</b>

Table A2-13: The GHG emissions from National Navigation (cont.)

	1990	1995	2000	2001	2002	2003	2004	2005
EF CH <sub>4</sub> – gasoline (kg/TJ)	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0
EF CH <sub>4</sub> – diesel (kg/TJ)	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0
EF CH <sub>4</sub> – fuel oil (kg/TJ)	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0
<b>CH<sub>4</sub> emission (Mg)</b>	<b>9.1</b>	<b>6.7</b>	<b>5.8</b>	<b>6.2</b>	<b>7.5</b>	<b>7.5</b>	<b>6.2</b>	<b>6.8</b>
EF N <sub>2</sub> O – gasoline (kg/TJ)	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
EF N <sub>2</sub> O – diesel (kg/TJ)	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
EF N <sub>2</sub> O – fuel oil (kg/TJ)	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
<b>N<sub>2</sub>O emission (Mg)</b>	<b>1.1</b>	<b>0.8</b>	<b>0.7</b>	<b>0.7</b>	<b>0.9</b>	<b>0.9</b>	<b>0.7</b>	<b>0.8</b>

Table A2-14: The GHG emissions from Railways

	1990	1995	2000	2001	2002	2003	2004	2005
<b>Fuel consumption</b>								
Gasoline (1000 t)			0.1					
NCV for gasoline (MJ/kg)			44.6					
Diesel (1000 t)	37.2	32.4	27.2	28.0	27.8	28.1	29.4	30.5
NCV for diesel (MJ/kg)	42.7	42.7	42.7	42.7	42.7	42.7	42.7	42.7
Fuel oil (1000 t)	0.2	1.5						
NCV for fuel oil (MJ/kg)	40.2	40.2						
Brown coal (1000 t)	10.0							
NCV for brown coal (MJ/kg)	16.7							
Lignite (1000 t)	4.3							
NCV for lignite (MJ/kg)	10.9							
<b>Total fuel consumption (TJ)</b>	<b>1811.1</b>	<b>1444.1</b>	<b>1166.2</b>	<b>1195.9</b>	<b>1187.3</b>	<b>1200.2</b>	<b>1255.7</b>	<b>1302.7</b>
<b>Emissions</b>								
EF CO <sub>2</sub> – gasoline (t/TJ)	68.6	68.6	68.6	68.6	68.6	68.6	68.6	68.6
EF CO <sub>2</sub> – diesel (t/TJ)	73.3	73.3	73.3	73.3	73.3	73.3	73.3	73.3
EF CO <sub>2</sub> – fuel oil (t/TJ)	76.6	76.6	76.6	76.6	76.6	76.6	76.6	76.6
EF CO <sub>2</sub> – brown coal (t/TJ)	94.1	94.1	94.1	94.1	94.1	94.1	94.1	94.1
EF CO <sub>2</sub> – lignite (t/TJ)	99.2	99.2	99.2	99.2	99.2	99.2	99.2	99.2
<b>CO<sub>2</sub> emission (Gg)</b>	<b>137.5</b>	<b>106.1</b>	<b>85.5</b>	<b>87.7</b>	<b>87.1</b>	<b>88.0</b>	<b>92.1</b>	<b>95.5</b>
EF CH <sub>4</sub> – gasoline (kg/TJ)	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0
EF CH <sub>4</sub> – diesel (kg/TJ)	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0
EF CH <sub>4</sub> – fuel oil (kg/TJ)	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0
EF CH <sub>4</sub> – brown coal (kg/TJ)	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0
EF CH <sub>4</sub> – lignite (kg/TJ)	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0
<b>CH<sub>4</sub> emission (Mg)</b>	<b>10.1</b>	<b>7.2</b>	<b>5.8</b>	<b>6.0</b>	<b>5.9</b>	<b>6.0</b>	<b>6.3</b>	<b>6.5</b>
EF N <sub>2</sub> O – gasoline (kg/TJ)	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
EF N <sub>2</sub> O – diesel (kg/TJ)	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
EF N <sub>2</sub> O – fuel oil (kg/TJ)	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
EF N <sub>2</sub> O – brown coal (kg/TJ)	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4
EF N <sub>2</sub> O – lignite (kg/TJ)	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4
<b>N<sub>2</sub>O emission (Mg)</b>	<b>1.3</b>	<b>0.9</b>	<b>0.7</b>	<b>0.7</b>	<b>0.7</b>	<b>0.7</b>	<b>0.8</b>	<b>0.8</b>

Table A2-15: The GHG emissions from Commercial/Institutional

	1990	1995	2000	2001	2002	2003	2004	2005
<b>Fuel consumption</b>								
Petroleum (1000 t)	0.1							
NCV for petroleum (MJ/kg)	44.0							
Extra light oil (1000 t)	89.5	101.4	120.6	130.7	141.4	144.5	143.8	131.6
NCV for extra light oil (MJ/kg)	42.7	42.7	42.7	42.7	42.7	42.7	42.7	42.7
Fuel oil (1000 t)	66.1		3.9	5	10.6	8.8	6.6	6.6
NCV for fuel oil (MJ/kg)	40.2		40.2	40.2	40.2	40.2	40.2	40.2
LPG (1000 t)	4.3	2.5	2.6	5.5	6.0	7.1	8.1	8.5
NCV for LPG (MJ/kg)	46.9	46.9	46.9	46.9	46.9	46.9	46.9	46.9
Anthracite (1000 t)				0.2				
NCV for anthracite (MJ/kg)				29.3				
Brown coal (1000 t)	24.5	12.7	9.5	5.6	8.4	10.9	4.8	0.2
NCV for brown coal (MJ/kg)	16.7	16.7	17.8	18.2	18.2	18.5	18.3	18.5
Lignite (1000 t)	40.0	1.6	1.2	2.4	2.6	2.2	0.6	0.6
NCV for lignite (MJ/kg)	10.9	10.9	12.0	12.2	12.2	12.3	12.2	12.1
Briquettes (1000 t)	2.9							
NCV for briquettes (MJ/kg)	16.7						16.7	16.7
Gas work gas (1000000 m <sup>3</sup> )	4.9	1.4	1.5	1.6	1.7	5.0	3.8	3.4
NCV for gas work gas (MJ/m <sup>3</sup> )	15.8	15.8	19.5	19.5	19.5	22.6	21.5	21.5
Natural gas (1000000 m <sup>3</sup> )	102.5	133.7	98.7	133.0	124.1	129.9	138.4	151.2
NCV for natural gas (MJ/m <sup>3</sup> )	34.0	34.0	34.0	34.0	34.0	34.0	34.0	34.0
<b>Total fuel consumption (TJ)</b>	<b>11142.3</b>	<b>9246.0</b>	<b>8998.0</b>	<b>10731.2</b>	<b>11183.6</b>	<b>11616.3</b>	<b>11669.6</b>	<b>11509.9</b>
<b>Emissions</b>								
EF CO <sub>2</sub> – petroleum (t/TJ)	71.1	71.1	71.1	71.1	71.1	71.1	71.1	71.1
EF CO <sub>2</sub> – diesel (t/TJ)	73.3	73.3	73.3	73.3	73.3	73.3	73.3	73.3
EF CO <sub>2</sub> – fuel oil (t/TJ)	76.6	76.6	76.6	76.6	76.6	76.6	76.6	76.6
EF CO <sub>2</sub> – LPG (t/TJ)	62.4	62.4	62.4	62.4	62.4	62.4	62.4	62.4
EF CO <sub>2</sub> – anthracite (t/TJ)	96.3	96.3	96.3	96.3	96.3	96.3	96.3	96.3
EF CO <sub>2</sub> – brown coal (t/TJ)	94.1	94.1	94.1	94.1	94.1	94.1	94.1	94.1
EF CO <sub>2</sub> – lignite (t/TJ)	99.2	99.2	99.2	99.2	99.2	99.2	99.2	99.2
EF CO <sub>2</sub> – briquettes (t/TJ)	94.1	94.1	94.1	94.1	94.1	94.1	94.1	94.1
EF CO <sub>2</sub> – gas work gas (t/TJ)	46.7	46.7	46.7	46.7	46.7	46.7	46.7	46.7
EF CO <sub>2</sub> – natural gas (t/TJ)	55.8	55.8	55.8	55.8	55.8	55.8	55.8	55.8
<b>CO<sub>2</sub> emission (Gg)</b>	<b>785.9</b>	<b>609.1</b>	<b>603.3</b>	<b>707.7</b>	<b>747.6</b>	<b>773.9</b>	<b>769.9</b>	<b>748.8</b>
EF CH <sub>4</sub> – petroleum (kg/TJ)	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0
EF CH <sub>4</sub> – diesel (kg/TJ)	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0
EF CH <sub>4</sub> – fuel oil (kg/TJ)	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0
EF CH <sub>4</sub> – LPG (kg/TJ)	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0
EF CH <sub>4</sub> – anthracite (kg/TJ)	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0
EF CH <sub>4</sub> – brown coal (kg/TJ)	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0
EF CH <sub>4</sub> – lignite (kg/TJ)	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0
EF CH <sub>4</sub> – briquettes (kg/TJ)	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0
EF CH <sub>4</sub> – gas work gas (kg/TJ)	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0
EF CH <sub>4</sub> – natural gas (kg/TJ)	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0
<b>CH<sub>4</sub> emission (Mg)</b>	<b>94.6</b>	<b>70.7</b>	<b>73.2</b>	<b>84.7</b>	<b>90.7</b>	<b>94.1</b>	<b>93.2</b>	<b>89.4</b>

Table A2-15: The GHG emissions from Commercial/Institutional (cont.)

	1990	1995	2000	2001	2002	2003	2004	2005
<b>Fuel consumption</b>								
EF N <sub>2</sub> O - petroleum (k/TJ)	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
EF N <sub>2</sub> O - diesel (kg/TJ)	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
EF N <sub>2</sub> O - fuel oil (kg/TJ)	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
EF N <sub>2</sub> O - LPG (kg/TJ)	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
EF N <sub>2</sub> O - anthracite (kg/TJ)	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4
EF N <sub>2</sub> O - brown coal (kg/TJ)	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4
EF N <sub>2</sub> O - lignite (kg/TJ)	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4
EF N <sub>2</sub> O - briquettes (kg/TJ)	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4
EF N <sub>2</sub> O - gas work gas (kg/TJ)	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4
EF N <sub>2</sub> O - natural gas (kg/TJ)	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
<b>N<sub>2</sub>O emission (Mg)</b>	<b>5.7</b>	<b>3.5</b>	<b>3.9</b>	<b>4.3</b>	<b>4.8</b>	<b>5.0</b>	<b>4.8</b>	<b>4.4</b>

Table A2-16: The GHG emissions from Residential sector

	1990	1995	2000	2001	2002	2003	2004	2005
<b>Fuel consumption</b>								
Petroleum (1000 t)		7.9	1.6	0.8	0.8	1.2	0.8	
NCV for petroleum (MJ/kg)		44.0	44.0	44.0	44.0	44.0	44.0	
Extra light oil (1000 t)	215.9	198.6	231.5	249.4	270.8	276.9	279.2	252.8
NCV for extra light oil (MJ/kg)	42.7	42.7	42.7	42.7	42.7	42.7	42.7	42.7
Fuel oil (1000 t)	48.7	6.5	8.1	11.6	24.5	21.5	15.3	15.4
NCV for fuel oil (MJ/kg)	40.2	40.2	40.2	40.2	40.2	40.2	40.2	40.2
LPG (1000 t)	97.9	57.3	51.9	53.5	55.9	57.4	61.3	60.9
NCV for LPG (MJ/kg)	46.9	46.9	46.9	46.9	46.9	46.9	46.9	46.9
Brown coal (1000 t)	123.1	11.1	12.0	1.8	2.7	7.0	5.2	14
NCV for brown coal (MJ/kg)	16.7	16.7	17.8	18.2	18.2	18.5	18.3	18.5
Lignite (1000 t)	207.3	10.8	15.0	12.3	18.0	19.6	12.0	11.7
NCV for lignite (MJ/kg)	10.9	10.9	12.0	12.2	12.2	12.3	12.2	12.1
Briquettes (1000 t)	6.1							
NCV for briquettes (MJ/kg)	16.7							
Gas work gas (1000000 m <sup>3</sup> )	24.4	11.8	9.9	9.9	10.8	10.8	10.6	10.24
NCV for gas work gas (MJ/m <sup>3</sup> )	15.8	15.8	19.5	19.5	19.5	22.6	21.5	21.5
Natural gas (1000000 m <sup>3</sup> )	230.0	381.3	496.6	561.5	548.7	633.1	629.5	687.8
NCV for natural gas (MJ/m <sup>3</sup> )	34.0	34.0	34.0	34.0	34.0	34.0	34.0	34
Biomass (TJ)	19080	11070	13410	10260	10370	13455	13140	13900
<b>Total fuel consumption (TJ)</b>	<b>47477.3</b>	<b>36301.9</b>	<b>43597.8</b>	<b>43388.0</b>	<b>44711.0</b>	<b>51028.3</b>	<b>50461.0</b>	<b>52177.2</b>
<b>Emissions</b>								
EF CO <sub>2</sub> - petroleum (t/TJ)	71.1	71.1	71.1	71.1	71.1	71.1	71.1	71.1
EF CO <sub>2</sub> - diesel (t/TJ)	73.3	73.3	73.3	73.3	73.3	73.3	73.3	73.3
EF CO <sub>2</sub> - fuel oil (t/TJ)	76.6	76.6	76.6	76.6	76.6	76.6	76.6	76.6
EF CO <sub>2</sub> - LPG (t/TJ)	62.4	62.4	62.4	62.4	62.4	62.4	62.4	62.4
EF CO <sub>2</sub> - brown coal (t/TJ)	94.1	94.1	94.1	94.1	94.1	94.1	94.1	94.1
EF CO <sub>2</sub> - lignite (t/TJ)	99.2	99.2	99.2	99.2	99.2	99.2	99.2	99.2
EF CO <sub>2</sub> - briquettes (t/TJ)	94.1	94.1	94.1	94.1	94.1	94.1	94.1	94.1
EF CO <sub>2</sub> - gas work gas (t/TJ)	46.7	46.7	46.7	46.7	46.7	46.7	46.7	46.7
EF CO <sub>2</sub> - natural gas (t/TJ)	55.8	55.8	55.8	55.8	55.8	55.8	55.8	55.8
EF CO <sub>2</sub> - biomass (t/TJ)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	107.4
<b>CO<sub>2</sub> Emission (Gg)</b>	<b>1994.8</b>	<b>1596.0</b>	<b>1896.3</b>	<b>2068.5</b>	<b>2167.2</b>	<b>2354.1</b>	<b>2332.2</b>	<b>3864.9</b>

Table A2-16: The GHG emissions from Residential sector (cont.)

	1990	1995	2000	2001	2002	2003	2004	2005
EF CH <sub>4</sub> - petroleum (k/TJ)	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0
EF CH <sub>4</sub> - diesel (kg/TJ)	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0
EF CH <sub>4</sub> - fuel oil (kg/TJ)	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0
EF CH <sub>4</sub> - LPG (kg/TJ)	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0
EF CH <sub>4</sub> - brown coal (kg/TJ)	300.0	300.0	300.0	300.0	300.0	300.0	300.0	300.0
EF CH <sub>4</sub> - lignite (kg/TJ)	300.0	300.0	300.0	300.0	300.0	300.0	300.0	300.0
EF CH <sub>4</sub> - briquettes (kg/TJ)	300.0	300.0	300.0	300.0	300.0	300.0	300.0	300.0
EF CH <sub>4</sub> - gas work gas (kg/TJ)	300.0	300.0	300.0	300.0	300.0	300.0	300.0	300.0
EF CH <sub>4</sub> - natural gas (kg/TJ)	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0
EF CH <sub>4</sub> - biomass (kg/TJ)	300.0	300.0	300.0	300.0	300.0	300.0	300.0	300.0
<b>CH<sub>4</sub> Emission (Mg)</b>	<b>7363.3</b>	<b>3650.7</b>	<b>4410.6</b>	<b>3422.6</b>	<b>3499.8</b>	<b>4482.4</b>	<b>4344.2</b>	<b>4615.8</b>
EF N <sub>2</sub> O - petroleum (k/TJ)	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
EF N <sub>2</sub> O - diesel (kg/TJ)	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
EF N <sub>2</sub> O - fuel oil (kg/TJ)	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
EF N <sub>2</sub> O - LPG (kg/TJ)	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
EF N <sub>2</sub> O - brown coal (kg/TJ)	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4
EF N <sub>2</sub> O - lignite (kg/TJ)	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4
EF N <sub>2</sub> O - briquettes (kg/TJ)	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4
EF N <sub>2</sub> O - gas work gas (kg/TJ)	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4
EF N <sub>2</sub> O - natural gas (kg/TJ)	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
EF N <sub>2</sub> O - biomass (kg/TJ)	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0
<b>N<sub>2</sub>O Emission (Mg)</b>	<b>93.3</b>	<b>53.3</b>	<b>63.8</b>	<b>51.7</b>	<b>53.1</b>	<b>66.1</b>	<b>64.6</b>	<b>67.4</b>

Table A2-17: The GHG emissions from Agriculture/Forestry/Fishing

	1990	1995	2000	2001	2002	2003	2004	2005
<b>Fuel consumption</b>								
Gasoline (1000 t)	4.0	7.8	12.1	10.5	8.2	8.1	7.2	8.1
NCV for gasoline (MJ/kg)	44.6	44.6	44.6	44.6	44.6	44.6	44.6	44.6
Other kerosene (1000 t)	0.1	0.1						
NCV for other kerosene (MJ/kg)	44.0	44.0						
Extra light oil (1000 t)	232.6	159.1	237.6	223.1	206	212.8	197.3	197.4
NCV for extra light oil (MJ/kg)	42.7	42.7	42.7	42.7	42.7	42.7	42.7	42.7
<b>Fuel consumption - mobile (TJ)</b>	<b>10117</b>	<b>7147</b>	<b>10687</b>	<b>9997</b>	<b>9164</b>	<b>9449</b>	<b>8748</b>	<b>8792.1</b>
Fuel oil (1000 t)	12.3	6.2	13.4	4.8	4.7	4.7	4.6	4.7
NCV for fuel oil (MJ/kg)	40.2	40.2	40.2	40.2	40.2	40.2	40.2	40.2
LPG (1000 t)	4.4	3.2	2.6	2.7	2.6	2.8	2.7	2.7
NCV for LPG (MJ/kg)	46.9	46.9	46.9	46.9	46.9	46.9	46.9	46.9
Gas work gas (1000000 m <sup>3</sup> )					0.1			
NCV for gas work gas (MJ/m <sup>3</sup> )					19.5			
Natural gas (1000000 m <sup>3</sup> )	25.0	15.5	14.5	23.6	24.3	19.9	19.4	23.2
NCV for natural gas (MJ/m <sup>3</sup> )	34.0	34.0	34.0	34.0	34.0	34.0	34.0	34.0
<b>Fuel consum. - stationary (TJ)</b>	<b>1550.7</b>	<b>926.2</b>	<b>1153.5</b>	<b>1121.9</b>	<b>1139.0</b>	<b>996.8</b>	<b>971.1</b>	<b>1104.3</b>
<b>Total fuel consumption (TJ)</b>	<b>11668</b>	<b>8074</b>	<b>11841</b>	<b>11119</b>	<b>10303</b>	<b>10447</b>	<b>9719</b>	<b>9896.4</b>



Table A2-17: The GHG emissions from Agriculture/Forestry/Fishing (cont.)

	1990	1995	2000	2001	2002	2003	2004	2005
<b>Emissions</b>								
EF CO <sub>2</sub> - gasoline (t/TJ)	68.6	68.6	68.6	68.6	68.6	68.6	68.6	68.6
EF CO <sub>2</sub> - other kerosene (t/TJ)	71.1	71.1	71.1	71.1	71.1	71.1	71.1	71.1
EF CO <sub>2</sub> - diesel (t/TJ)	73.3	73.3	73.3	73.3	73.3	73.3	73.3	73.3
<b>CO<sub>2</sub> emission (Gg) - mobile</b>	<b>741.0</b>	<b>522.4</b>	<b>781.1</b>	<b>730.8</b>	<b>670.2</b>	<b>691.2</b>	<b>639.9</b>	<b>643.0</b>
EF CO <sub>2</sub> - fuel oil (t/TJ)	76.6	76.6	76.6	76.6	76.6	76.6	76.6	76.6
EF CO <sub>2</sub> - LPG (t/TJ)	62.4	62.4	62.4	62.4	62.4	62.4	62.4	62.4
EF CO <sub>2</sub> - gas work gas (t/TJ)	106.0	106.0	106.0	106.0	106.0	106.0	106.0	106.0
EF CO <sub>2</sub> - natural gas (t/TJ)	55.8	55.8	55.8	55.8	55.8	55.8	55.8	55.8
<b>CO<sub>2</sub> emission (Gg) - stationary</b>	<b>98.2</b>	<b>57.9</b>	<b>76.4</b>	<b>67.5</b>	<b>68.4</b>	<b>60.4</b>	<b>58.9</b>	<b>66.4</b>
<b>Total CO<sub>2</sub> emission (Gg)</b>	<b>839.2</b>	<b>580.3</b>	<b>857.5</b>	<b>798.3</b>	<b>738.6</b>	<b>751.6</b>	<b>698.8</b>	<b>709.4</b>
EF CH <sub>4</sub> - gasoline (kg/TJ)	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0
EF CH <sub>4</sub> - other kerosene (kg/TJ)	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0
EF CH <sub>4</sub> - diesel (kg/TJ)	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0
<b>CH<sub>4</sub> emission (Mg) - mobile</b>	<b>50.6</b>	<b>35.7</b>	<b>53.4</b>	<b>50.0</b>	<b>45.8</b>	<b>47.2</b>	<b>43.7</b>	<b>44.0</b>
EF CH <sub>4</sub> - fuel oil (kg/TJ)	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0
EF CH <sub>4</sub> - LPG (kg/TJ)	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0
EF CH <sub>4</sub> - gas work gas (kg/TJ)	300.0	300.0	300.0	300.0	300.0	300.0	300.0	300.0
EF CH <sub>4</sub> - natural gas (kg/TJ)	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0
<b>CH<sub>4</sub> emission (Mg) - stationary</b>	<b>11.3</b>	<b>6.6</b>	<b>9.1</b>	<b>7.2</b>	<b>7.8</b>	<b>6.6</b>	<b>6.4</b>	<b>7.1</b>
<b>Total CH<sub>4</sub> emission (Mg)</b>	<b>61.8</b>	<b>42.4</b>	<b>62.5</b>	<b>57.2</b>	<b>53.6</b>	<b>53.8</b>	<b>50.2</b>	<b>51.1</b>
EF N <sub>2</sub> O - gasoline (kg/TJ)	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
EF N <sub>2</sub> O - other kerosene (kg/TJ)	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
EF N <sub>2</sub> O - diesel (kg/TJ)	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
<b>N<sub>2</sub>O emission (Mg) - mobile</b>	<b>6.1</b>	<b>4.3</b>	<b>6.4</b>	<b>6.0</b>	<b>5.5</b>	<b>5.7</b>	<b>5.2</b>	<b>5.3</b>
EF N <sub>2</sub> O - fuel oil (kg/TJ)	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
EF N <sub>2</sub> O - LPG (kg/TJ)	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
EF N <sub>2</sub> O - gas work gas (kg/TJ)	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4
EF N <sub>2</sub> O - natural gas (kg/TJ)	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
<b>N<sub>2</sub>O emission (Mg) - stationary</b>	<b>0.5</b>	<b>0.3</b>	<b>0.4</b>	<b>0.3</b>	<b>0.3</b>	<b>0.3</b>	<b>0.3</b>	<b>0.3</b>
<b>Total N<sub>2</sub>O emission (Mg)</b>	<b>6.6</b>	<b>4.6</b>	<b>6.9</b>	<b>6.3</b>	<b>5.8</b>	<b>5.9</b>	<b>5.5</b>	<b>5.5</b>

Table A2-18: Methane emissions from Coal Mining and Handling from 1990 to 1999

Source and Sink Categories		Activity Data Production (PJ)	Emission Estimates CH <sub>4</sub> (Gg)	Emission Factor kgCH <sub>4</sub> /t	Emission Factor m <sup>3</sup> CH <sub>4</sub> /t
<b>Year 1990</b>					
1B 1a Underground mines			<b>2.32</b>		
	Mining	0.174	2.04	5.86	17.50
	Post-Mining	0.174	0.29	0.82	2.45
<b>Year 1991</b>					
1B 1a Underground mines			<b>2.07</b>		
	Mining	0.155	1.82	5.86	17.50
	Post-Mining	0.155	0.25	0.82	2.45
<b>Year 1992</b>					
1B 1a Underground mines			<b>1.61</b>		
	Mining	0.120	1.41	5.86	17.50
	Post-Mining	0.120	0.20	0.82	2.45
<b>Year 1993</b>					
1B 1a Underground mines			<b>1.54</b>		
	Mining	0.115	1.35	5.86	17.50
	Post-Mining	0.115	0.19	0.82	2.45
<b>Year 1994</b>					
1B 1a Underground mines			<b>1.38</b>		
	Mining	0.103	1.21	5.86	17.50
	Post-Mining	0.103	0.17	0.82	2.45
<b>Year 1995</b>					
1B 1a Underground mines			<b>1.10</b>		
	Mining	0.082	0.96	5.86	17.50
	Post-Mining	0.082	0.13	0.82	2.45
<b>Year 1996</b>					
1B 1a Underground Mines			<b>0.89</b>		
	Mining	0.066	0.78	5.86	17.50
	Post-Mining	0.066	0.11	0.82	2.45
<b>Year 1997</b>					
1B 1a Underground Mines			<b>0.65</b>		
	Mining	0.049	0.57	5.86	17.50
	Post-Mining	0.049	0.08	0.82	2.45
<b>Year 1998</b>					
1B 1a Underground Mines			<b>0.68</b>		
	Mining	0.051	0.60	5.86	17.50
	Post-Mining	0.051	0.08	0.82	2.45
<b>Year 1999</b>					
1B 1a Underground Mines			<b>0.20</b>		
	Mining	0.015	0.18	5.86	17.50
	Post-Mining	0.015	0.03	0.82	2.45

\* - 0.67 kg/m<sup>3</sup> – Methane density at 20 °C and pressure 1 atm.

Table A2-19: Methane emissions from Oil and Gas Activities, years 1990, 1995, 2000, 2005

Source and sink categories		Activity data Fuel Quantity PJ	Emission Estimates CH <sub>4</sub> Gg	Emission Factor kgCH <sub>4</sub> /PJ
<b>Year 1990</b>				
1B 2a Oil			<b>0.68</b>	
	Production	112.9	0.30	2650
	Transport	174.1	0.13	745
	Refining	287.3	0.21	135
	Storage	287.3	0.04	135
1B 2b Natural gas			<b>54.59</b>	
	Prod./Process./Trans./Distrib.	67.4	30.87 <sup>1)</sup>	458000
	Other Leakage (non-residential)	78.4	21.93 <sup>2)</sup>	279500
	Other Leakage (residential)	12.9	1.80 <sup>3)</sup>	139500
1B 2c Venting and flaring			<b>1.21</b>	
	Gas	67.4	1.21	18000
<b>Year 1995</b>				
1B 2a Oil			<b>0.49</b>	
	Production	62.8	0.17	2650
	Transport	159.3	0.12	745
	Refining	227.6	0.17	135
	Storage	227.6	0.03	135
1B 2b Natural gas			<b>50.60</b>	
	Prod./Process./Trans./Distrib.	66.9	30.62 <sup>1)</sup>	458000
	Other Leakage (non-residential)	62.5	17.47 <sup>2)</sup>	279500
	Other Leakage (residential)	18.0	2.51 <sup>3)</sup>	139500
1B 2c Venting and flaring			<b>1.20</b>	
	Gas	66.9	1.20	18000
<b>Year 2000</b>				
1B 2a Liquid Fossil Fuel			<b>0.45</b>	
	Production	51.4	0.14	2650
	Transport	165.6	0.12	745
	Refining	218.4	0.16	135
	Storage	218.4	0.03	135
1B 2b Natural Gas			<b>51.39</b>	
	Prod./Process./Trans./Distrib.	59.4	27.21 <sup>1)</sup>	458000
	Other Leakage(non-residential)	78.1	21.83 <sup>2)</sup>	279500
	Other Leakage (residential)	16.9	2.35 <sup>3)</sup>	139500
1B 2c Venting and Flaring			<b>1.07</b>	
	Gas	59.4	1.07	18000
<b>Year 2005</b>				
1B 2a Liquid Fossil Fuel			<b>0.414</b>	
	Production	40.110	0.106	2650
	Transport	169.540	0.126	745
	Refining	206.700	0.154	135
	Storage	206.7	0.028	135
1B 2b Natural Gas			<b>61.670</b>	
	Prod./Process./Trans./Distrib.	79.761	36.531	458000
	Other Leakage(non-residential)	78.270	21.876	279500
	Other Leakage (residential)	23.390	3.263	139500
1B 2c Venting and Flaring			<b>1.4</b>	
	Gas	79.761	1.4	18000

<sup>1)</sup> – Methane emissions from Processing, Transmission and Distribution<sup>2)</sup> – Other Leakage at Industrial Plants and Power Stations<sup>3)</sup> – Other Leakage in Residential and Commercial Sectors

## **ANNEX 3**

### **CO<sub>2</sub> REFERENCE APPROACH AND COMPARISON WITH SECTORAL APPROACH, AND RELEVANT INFORMATION ON THE NATIONAL ENERGY BALANCE**

Table A3-1: Fuel combustion CO<sub>2</sub> emissions (Reference and Sectoral Approach)

YEAR	FUEL TYPES	Reference approach		Sectoral approach		Difference	
		Energy Consumpt. excluding non-energy (PJ)	CO <sub>2</sub> emissions (Gg)	Energy consumption (PJ)	CO <sub>2</sub> emissions (Gg)	Energy consumption (%)	CO <sub>2</sub> emissions (%)
1990	Liquid Fuels	173.760	12792.36	181.020	13265.56	-4.011	-3.567
	Solid Fuels	32.152	3102.87	32.152	3161.89	0.000	-1.867
	Gaseous Fuels	83.086	4946.66	74.021	4101.27	12.246	20.613
	<b>Total</b>	<b>288.998</b>	<b>20841.91</b>	<b>287.193</b>	<b>20528.73</b>	<b>0.628</b>	<b>1.526</b>
1991	Liquid Fuels	122.573	8988.23	129.608	9494.29	-5.428	-5.330
	Solid Fuels	21.044	1850.52	21.044	1945.44		-4.879
	Gaseous Fuels	76.735	4576.39	64.811	3583.66	18.398	27.702
	<b>Total</b>	<b>220.352</b>	<b>15415.14</b>	<b>215.463</b>	<b>15023.39</b>	<b>2.269</b>	<b>2.608</b>
1992	Liquid Fuels	115.571	8462.96	122.618	9029.98	-5.747	-6.279
	Solid Fuels	16.340	1433.76	16.340	1405.74		1.993
	Gaseous Fuels	77.494	4706.59	67.974	3756.04	14.005	25.307
	<b>Total</b>	<b>209.404</b>	<b>14603.31</b>	<b>206.931</b>	<b>14191.76</b>	<b>1.195</b>	<b>2.900</b>
1993	Liquid Fuels	114.699	8394.74	126.727	9316.05	-9.491	-9.889
	Solid Fuels	13.717	1176.38	13.717	1165.77		0.910
	Gaseous Fuels	92.369	5458.96	78.670	4372.20	17.413	24.856
	<b>Total</b>	<b>220.785</b>	<b>15030.08</b>	<b>219.113</b>	<b>14854.02</b>	<b>0.763</b>	<b>1.185</b>
1994	Liquid Fuels	121.697	9050.62	125.802	9225.41	-3.263	-1.895
	Solid Fuels	8.422	753.01	8.422	720.825		4.465
	Gaseous Fuels	78.958	4712.19	73.545	4085.46	7.360	15.341
	<b>Total</b>	<b>209.076</b>	<b>14515.82</b>	<b>207.769</b>	<b>14031.70</b>	<b>0.629</b>	<b>3.450</b>
1995	Liquid Fuels	137.496	10095.79	142.567	10467.89	-3.557	-3.555
	Solid Fuels	7.653	696.283	7.653	713.279		-2.383
	Gaseous Fuels	72.594	4348.17	65.975	3661.58	10.034	18.751
	<b>Total</b>	<b>217.744</b>	<b>15140.24</b>	<b>216.195</b>	<b>14842.75</b>	<b>0.716</b>	<b>2.004</b>
1996	Liquid Fuels	141.592	10601.74	145.549	10682.59	-2.718	-0.757
	Solid Fuels	6.593	581.76	6.593	610.025	0.000	-4.633
	Gaseous Fuels	81.576	4876.64	74.874	4159.03	8.950	17.254
	<b>Total</b>	<b>229.761</b>	<b>16060.14</b>	<b>227.016</b>	<b>15451.65</b>	<b>1.209</b>	<b>3.938</b>
1997	Liquid Fuels	147.704	10608.77	150.030	10976.47	-1.550	-3.350
	Solid Fuels	10.552	948.590	10.552	977.738	0.000	-2.981
	Gaseous Fuels	84.088	5046.39	78.829	4380.25	6.672	15.208
	<b>Total</b>	<b>242.344</b>	<b>16603.76</b>	<b>239.410</b>	<b>16334.46</b>	<b>1.226</b>	<b>1.649</b>
1998	Liquid Fuels	158.342	11790.93	163.747	12038.57	-3.301	-2.057
	Solid Fuels	10.290	920.050	10.290	948.906		-3.041
	Gaseous Fuels	89.906	4888.27	78.184	4349.16	14.993	12.396
	<b>Total</b>	<b>258.539</b>	<b>17599.25</b>	<b>252.221</b>	<b>17336.63</b>	<b>2.505</b>	<b>1.515</b>
1999	Liquid Fuels	171.561	12695.27	174.360	12846.34	-1.605	-1.176
	Solid Fuels	9.061	803.387	9.061	830.609		-3.277
	Gaseous Fuels	82.213	4923.22	77.535	4307.51	6.034	14.294
	<b>Total</b>	<b>262.836</b>	<b>18421.87</b>	<b>260.956</b>	<b>17984.46</b>	<b>0.720</b>	<b>2.432</b>
2000	Liquid Fuels	148.689	11039.96	153.326	11222.38	-3.024	-1.625
	Solid Fuels	19.027	1732.77	19.027	1765.04		-1.828
	Gaseous Fuels	82.904	4966.47	78.678	4267.38	5.371	16.382
	<b>Total</b>	<b>250.621</b>	<b>17739.21</b>	<b>251.032</b>	<b>17254.80</b>	<b>-0.164</b>	<b>2.807</b>
2001	Liquid Fuels	154.151	11456.23	158.819	11661.00	-2.939	-1.756
	Solid Fuels	20.062	1842.48	20.062	1854.62	0.000	-0.654
	Gaseous Fuels	89.072	5244.62	82.940	4610.82	7.393	13.746
	<b>Total</b>	<b>263.285</b>	<b>18543.33</b>	<b>261.822</b>	<b>18126.44</b>	<b>0.559</b>	<b>2.300</b>

Table A3-1: Fuel combustion CO<sub>2</sub> emissions (Reference and Sectoral Approach) - cont.

YEAR	FUEL TYPES	Reference approach		Sectoral approach		Difference	
		Energy Consumpt. excluding non-energy (PJ)	CO <sub>2</sub> emissions (Gg)	Energy consumption (PJ)	CO <sub>2</sub> emissions (Gg)	Energy consumption (%)	CO <sub>2</sub> emissions (%)
2002	Liquid Fuels	164.311	12194.47	165.655	12167.18	-0.811	0.224
	Solid Fuels	24.896	2224.09	24.896	2300.42	0.000	-3.318
	Gaseous Fuels	92.098	5386.32	86.407	4805.84	6.586	12.079
	<b>Total</b>	<b>281.304</b>	<b>19804.88</b>	<b>276.957</b>	<b>19273.44</b>	<b>1.570</b>	<b>2.757</b>
2003	Liquid Fuels	178.899	13299.37	181.697	13355.41	-1.540	-0.420
	Solid Fuels	27.558	2532.94	27.558	2545.21		-0.482
	Gaseous Fuels	90.696	5338.38	85.468	4750.53	6.117	12.374
	<b>Total</b>	<b>297.154</b>	<b>21170.69</b>	<b>294.723</b>	<b>20651.15</b>	<b>0.825</b>	<b>2.516</b>
2004	Liquid Fuels	165.498	12324.05	167.575	12312.60	-1.239	0.093
	Solid Fuels	28.797	2687.52	28.797	2663.00		0.921
	Gaseous Fuels	93.328	5545.67	91.241	5072.16	2.288	9.335
	<b>Total</b>	<b>165.498</b>	<b>12324.05</b>	<b>167.575</b>	<b>12312.59</b>	<b>-1.239</b>	<b>0.093</b>
2005	Liquid Fuels	172.273	12893.25	173.136	12825.11	-0.498	0.531
	Solid Fuels	28.939	2667.30	28.939	2681.01		-0.511
	Gaseous Fuels	90.130	5360.38	88.319	4910.28	2.051	9.166
	<b>Total</b>	<b>291.342</b>	<b>20920.93</b>	<b>290.394</b>	<b>20416.41</b>	<b>0.327</b>	<b>2.471</b>

\* - Excluding international bunkers

Table A3-2: Net calorific values for different fossil fuels from 1990 to 2004

			Net calorific values 1990- 2005 MJ/kg(m <sup>3</sup> )
Liquid Fossil	Primary Fuel	Crude Oil	41.87-42.4
	Secondary Fuel	Motor Gasoline	44.59
		Jet Kerosene	43.96
		Gas/Diesel Oil	42.71
		Residual Fuel Oil	40.19
		LPG	46.89
		Naphtha	44.59
		Bitumen	33.5
		Lubricants	33.5
		Refinery Gas	48.57
		Petroleum Coke	29.31-31
		Ethane	47.31
Solid Fossil	Primary Fuel	Anthracite	29.29-29.31
		Other Bituminous Coal	24.3-26.9
		Sub Bituminous Coal	16.74-18.73
		Lignite	10.52-12.15
	Secondary Fuel	Gas Work Gas	15.82-22.63
		Coke Oven Coke	29.31
			TJ/Mm <sup>3</sup>
Natural Gas		Natural Gas	34
Biomass		Solid Biomass Fuel Wood	9

Table A3-3: National energy balance for 2005

	Anthracite		Hard coal		Brown coal		Lignite	
	10 <sup>3</sup> t	PJ	10 <sup>3</sup> t	PJ	10 <sup>3</sup> t	PJ	10 <sup>3</sup> t	PJ
Production	-	-	-	-	-	-	-	-
Import	0.3	0.01	957.6	24.04	70.1	1.30	12.5	0.15
Export	-	-	-	-	-	-	-	-
Stock change	-	-	99.2	2.49	-	-	-	-
International marine bunkers	-	-	-	-	-	-	-	-
<b>Energy supplied</b>	<b>0.3</b>	<b>0.01</b>	<b>1056.8</b>	<b>26.53</b>	<b>70.1</b>	<b>1.30</b>	<b>12.5</b>	<b>0.15</b>
<b>Energy sector own use</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>
-oil and gas extraction	-	-	-	-	-	-	-	-
-electric energy supply industry	-	-	-	-	-	-	-	-
-oil refineries	-	-	-	-	-	-	-	-
-NGL plant	-	-	-	-	-	-	-	-
<b>Total transformation sector</b>	<b>0.0</b>	<b>0.0</b>	<b>887.5</b>	<b>22.28</b>	<b>46.7</b>	<b>0.86</b>	<b>0.0</b>	<b>0.0</b>
-hydro power plants	-	-	-	-	-	-	-	-
-thermal power plants	-	-	887.5	22.28	-	-	-	-
-public cogeneration plants	-	-	-	-	-	-	-	-
-public heating plants	-	-	-	-	-	-	-	-
-indust. cogeneration plants	-	-	-	-	46.7	0.86	-	-
-industrial heating plants	-	-	-	-	-	-	-	-
-petroleum refineries	-	-	-	-	-	-	-	-
-NGL-plant	-	-	-	-	-	-	-	-
-gas works	-	-	-	-	-	-	-	-
<b>Non energy use</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>
<b>Losses</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>
<b>Final energy demand</b>	<b>0.3</b>	<b>0.01</b>	<b>169.3</b>	<b>4.25</b>	<b>23.4</b>	<b>0.43</b>	<b>12.5</b>	<b>0.15</b>
<b>Industry</b>	<b>0.3</b>	<b>0.01</b>	<b>169.3</b>	<b>4.25</b>	<b>9.2</b>	<b>0.17</b>	<b>0.2</b>	<b>0.00</b>
-iron and steel	-	-	1.0	0.03	-	-	-	-
-non-ferrous metals	-	-	-	-	-	-	-	-
-non-metallic minerals	0.1	0.00	-	-	-	-	-	-
-chemical	0.2	0.01	-	-	-	-	-	-
-construction materials	-	-	168.3	4.22	5.0	0.09	-	-
-pulp and paper	-	-	-	-	-	-	-	-
-food production	-	-	-	-	-	-	-	-
-not elsewhere specified	-	-	-	-	4.2	0.08	0.2	0.00
<b>Other sectors</b>	<b>-</b>	<b>-</b>	<b>-</b>	<b>-</b>	<b>14.2</b>	<b>0.26</b>	<b>12.3</b>	<b>0.15</b>
-households	-	-	-	-	14.0	0.26	11.7	0.14
-services	-	-	-	-	0.2	0.00	0.6	0.01
-agriculture	-	-	-	-	-	-	-	-
-construction	-	-	-	-	-	-	-	-
<b>Transport</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>
-rail	-	-	-	-	-	-	-	-
-road	-	-	-	-	-	-	-	-
-air	-	-	-	-	-	-	-	-
-sea and river	-	-	-	-	-	-	-	-
-public city	-	-	-	-	-	-	-	-

Table A3-3: National energy balance for 2005 (continue)

	Crude Oil		Natural gas		Fuel wood		Industrial waste
	10 <sup>3</sup> t	PJ	10 <sup>6</sup> m <sup>3</sup>	PJ	10 <sup>3</sup> m <sup>3</sup>	PJ	TJ
Production	946.0	40.11	2283.4	79.761	-	-	2362.5
Import	3998.7	169.54	1134.1	38.559	1390.0	12.51	-
Export	-	-	446.6	15.184	-	-	-
Stock change	46.6	1.97	-61.0	2.074	-	-	-
International marine bunkers	-	-	-	-	-	-	-
<b>Energy supplied</b>	<b>4991.1</b>	<b>211.62</b>	<b>2909.9</b>	<b>101.06</b>	<b>1390.0</b>	<b>12.51</b>	<b>2362.5</b>
<b>Energy sector own use</b>	<b>0.0</b>	<b>0.0</b>	<b>107.9</b>	<b>3.36</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>
-oil and gas extraction	-	-	98.7	1.91	-	-	-
-electric energy supply industry	-	-	-	-	-	-	-
-oil refineries	-	-	1.2	0.04	-	-	-
-NGL plant	-	-	8.0	0.27	-	-	-
<b>Total transformation sector</b>	<b>4991.1</b>	<b>211.62</b>	<b>988.2</b>	<b>35.72</b>	<b>0.0</b>	<b>0.0</b>	<b>2362.5</b>
-hydro power plants	-	-	-	-	-	-	-
-thermal power plants	-	-	36.3	1.23	-	-	-
-public cogeneration plants	-	-	479.0	16.29	-	-	-
-public heating plants	-	-	71.3	2.42	-	-	-
-industrial cogeneration plants	-	-	286.1	9.73	-	-	105.6
-industrial heating plants	-	-	83.3	2.83	-	-	2151.6
-petroleum refineries	4875.0	206.7	-	-	-	-	-
-NGL-plant	116.1	4.92	32.2	3.22	-	-	-
-gas works	-	-	-	-	-	-	-
<b>Non energy use</b>	<b>0.0</b>	<b>0.0</b>	<b>259.0</b>	<b>8.86</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>
<b>Losses</b>	<b>0.0</b>	<b>0.0</b>	<b>61.9</b>	<b>2.10</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>
<b>Final energy demand</b>	<b>0.0</b>	<b>0.0</b>	<b>1751.9</b>	<b>43.31</b>	<b>1390.0</b>	<b>12.51</b>	<b>-</b>
<b>Industry</b>	<b>0.0</b>	<b>0.0</b>	<b>604.4</b>	<b>21.61</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>
-iron and steel	-	-	20.1	0.68	-	-	-
-non-ferrous metals	-	-	1.0	0.03	-	-	-
-non-metallic minerals	-	-	70.9	2.41	-	-	-
-chemical	-	-	298.2	10.23	-	-	-
-construction materials	-	-	124.2	4.22	-	-	-
-pulp and paper	-	-	2.1	0.07	-	-	-
-food production	-	-	62.6	2.13	-	-	-
-not elsewhere specified	-	-	51.6	1.75	-	-	-
<b>Other sectors</b>	<b>0.0</b>	<b>0.0</b>	<b>862.2</b>	<b>29.31</b>	<b>1390.0</b>	<b>12.51</b>	<b>0.0</b>
-households	-	-	687.8	23.39	1390.0	12.51	-
-services	-	-	151.2	5.14	-	-	-
-agriculture	-	-	23.2	0.79	-	-	-
-construction	-	-	-	-	-	-	-
<b>Transport</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>
-rail	-	-	-	-	-	-	-
-road	-	-	-	-	-	-	-
-air	-	-	-	-	-	-	-
-sea and river	-	-	-	-	-	-	-
-public city	-	-	-	-	-	-	-



Table A3-3: National energy balance for 2005 (continue)

	Coke oven coke		Liquefied petroleum gases		Unleaded motor gasoline		Standard motor gasoline	
	10 <sup>3</sup> t	PJ	10 <sup>3</sup> t	PJ	10 <sup>3</sup> t	PJ	10 <sup>3</sup> t	PJ
Production	-	-	364.2	17.08	886.9	39.55	280.7	12.52
Import	22.9	0.67	3.4	0.16	255.8	11.41	1.3	0.06
Export	0.3	0.01	226.6	10.63	460.0	20.51	246.2	10.98
Stock change	-	-	-2.9	-0.14	13.6	0.21	-13.6	-0.61
International marine bunkers	-	-	-	-	-	-	-	-
<b>Energy supplied</b>	<b>22.6</b>	<b>0.66</b>	<b>138.1</b>	<b>10.60</b>	<b>687.4</b>	<b>30.65</b>	<b>22.2</b>	<b>0.99</b>
<b>Energy sector own use</b>	<b>0.0</b>	<b>0.0</b>	<b>6.6</b>	<b>0.31</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>
-oil and gas extraction	-	-	-	-	-	-	-	-
-electric energy supply industry	-	-	-	-	-	-	-	-
-oil refineries	-	-	6.6	0.31	-	-	-	-
-NGL plant	-	-	-	-	-	-	-	-
<b>Total transformation sector</b>	<b>0.0</b>	<b>0.0</b>	<b>16.0</b>	<b>0.75</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>
-hydro power plants	-	-	-	-	-	-	-	-
-thermal power plants	-	-	-	-	-	-	-	-
-public cogeneration plants	-	-	-	-	-	-	-	-
-public heating plants	-	-	-	-	-	-	-	-
-industrial cogeneration plants	-	-	2.9	0.14	-	-	-	-
-industrial heating plants	-	-	1.5	0.07	-	-	-	-
-petroleum refineries	-	-	-	-	-	-	-	-
-NGL-plant	-	-	-	-	-	-	-	-
-gas works	-	-	11.6	0.54	-	-	-	-
<b>Non energy use</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>
<b>Losses</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>
<b>Final energy demand</b>	<b>22.6</b>	<b>0.66</b>	<b>115.5</b>	<b>5.42</b>	<b>687.4</b>	<b>39.55</b>	<b>22.2</b>	<b>0.99</b>
<b>Industry</b>	<b>22.6</b>	<b>0.66</b>	<b>18.7</b>	<b>0.88</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>
-iron and steel	4.3	0.13	4.2	0.20	-	-	-	-
-non-ferrous metals	-	-	2.1	0.10	-	-	-	-
-non-metallic minerals	7.7	0.23	2.2	0.10	-	-	-	-
-chemical	-	-	-	-	-	-	-	-
-construction materials	-	-	4.6	0.22	-	-	-	-
-pulp and paper	-	-	0.1	0.00	-	-	-	-
-food production	9.6	0.28	0.7	0.03	-	-	-	-
-not elsewhere specified	1.0	0.03	4.8	0.23	-	-	-	-
<b>Other sectors</b>	<b>0.0</b>	<b>0.0</b>	<b>74.4</b>	<b>3.50</b>	<b>13.0</b>	<b>0.58</b>	<b>2.0</b>	<b>0.09</b>
-households	-	-	60.9	2.86	-	-	-	-
-services	-	-	8.5	0.40	-	-	-	-
-agriculture	-	-	2.7	0.13	8.1	0.36	-	-
-construction	-	-	2.6	0.12	4.9	0.22	2.0	0.09
<b>Transport</b>	<b>0.0</b>	<b>0.0</b>	<b>22.1</b>	<b>1.04</b>	<b>674.4</b>	<b>30.07</b>	<b>22.2</b>	<b>0.90</b>
-rail	-	-	-	-	-	-	-	-
-road	-	-	22.1	1.04	674.4	30.07	19.1	0.85
-air	-	-	-	-	-	-	1.1	0.05
-sea and river	-	-	-	-	-	-	-	-
-public city	-	-	-	-	-	-	-	-

Table A3-3: National energy balance for 2005 (continue)

	Petroleum		Jet fuel		Diesel oil		Light heating oil	
	10 <sup>3</sup> t	PJ	10 <sup>3</sup> t	PJ	10 <sup>3</sup> t	PJ	10 <sup>3</sup> t	PJ
Production	0.7	0.03	98.6	4.33	1080.9	46.17	522.2	22.30
Import	0.9	0.04	2.7	0.12	502.9	21.48	39.8	1.70
Export	0.5	0.02	1.7	0.07	253.9	10.84	95.0	4.06
Stock change	-0.1	-	-6.6	-0.29	-9.3	-0.40	-0.5	-0.02
International marine bunkers	-	-	-	-	9.1	0.39	-	-
<b>Energy supplied</b>	<b>1.0</b>	<b>0.04</b>	<b>93.0</b>	<b>4.09</b>	<b>1311.5</b>	<b>56.01</b>	<b>466.5</b>	<b>19.92</b>
<b>Energy sector own use</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>5.5</b>	<b>0.23</b>
-oil and gas extraction	-	-	-	-	-	-	5.5	0.23
-electric energy supply industry	-	-	-	-	-	-	-	-
-oil refineries	-	-	-	-	-	-	-	-
-NGL plant	-	-	-	-	-	-	-	-
<b>Total transformation sector</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>18.0</b>	<b>0.77</b>
-hydro power plants	-	-	-	-	-	-	-	-
-thermal power plants	-	-	-	-	-	-	3.0	0.3
-public cogeneration plants	-	-	-	-	-	-	1.5	0.06
-public heating plants	-	-	-	-	-	-	6.7	0.29
-industrial cogeneration plants	-	-	-	-	-	-	-	-
-industrial heating plants	-	-	-	-	-	-	6.8	0.29
-petroleum refineries	-	-	-	-	-	-	-	-
-NGL-plant	-	-	-	-	-	-	-	-
-gas works	-	-	-	-	-	-	-	-
<b>Non energy use</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>
<b>Losses</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>
<b>Final energy demand</b>	<b>1.0</b>	<b>0.04</b>	<b>93.0</b>	<b>4.09</b>	<b>1311.5</b>	<b>56.01</b>	<b>443.0</b>	<b>18.92</b>
<b>Industry</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>31.3</b>	<b>1.34</b>
-iron and steel	-	-	-	-	-	-	2.5	0.11
-non-ferrous metals	-	-	-	-	-	-	0.2	0.01
-non-metallic minerals	-	-	-	-	-	-	2.7	0.12
-chemical	-	-	-	-	-	-	0.5	0.02
-construction materials	-	-	-	-	-	-	6.9	0.29
-pulp and paper	-	-	-	-	-	-	0.1	0.00
-food production	-	-	-	-	-	-	9.5	0.41
-not elsewhere specified	-	-	-	-	-	-	8.9	0.38
<b>Other sectors</b>	<b>1.0</b>	<b>0.04</b>	<b>0.0</b>	<b>0.0</b>	<b>293.6</b>	<b>12.54</b>	<b>411.7</b>	<b>17.58</b>
-households	1.0	0.04	-	-	-	-	252.8	10.80
-services	-	-	-	-	-	-	131.6	5.62
-agriculture	-	-	-	-	183.0	7.82	14.4	0.62
-construction	-	-	-	-	110.6	4.72	12.9	0.55
<b>Transport</b>	<b>0.0</b>	<b>0.0</b>	<b>93.0</b>	<b>4.09</b>	<b>1017.9</b>	<b>43.47</b>	<b>0.0</b>	<b>0.0</b>
-rail	-	-	-	-	30.5	1.30	-	-
-road	-	-	-	-	927.3	39.60	-	-
-air	-	-	93.0	4.09	-	-	-	-
-sea and river	-	-	-	-	31.8	1.36	-	-
-public city	-	-	-	-	28.3	1.21	-	-

Table A3-3: National energy balance for 2005 (continue)

	Low sulphur fuel oil		Standard fuel oil		Naphta		White spirit	
	10 <sup>3</sup> t	PJ	10 <sup>3</sup> t	PJ	10 <sup>3</sup> t	PJ	10 <sup>3</sup> t	PJ
Production	-	-	1159.6	46.60	211.0	9.41	-	-
Import	84.9	3.41	0.1	0.00	-	-	5.0	0.17
Export	-	-	223.6	9.39	160.6	7.16	-	-
Stock change	-	-	-30.3	-1.22	-50.4	-2.25	-	-
International marine bunkers	-	-	16.4	0.66	-	-	-	-
<b>Energy supplied</b>	<b>84.9</b>	<b>3.41</b>	<b>879.4</b>	<b>35.33</b>	<b>0.0</b>	<b>0.0</b>	<b>5.0</b>	<b>0.17</b>
<b>Energy sector own use</b>	<b>0.0</b>	<b>0.0</b>	<b>70.8</b>	<b>2.85</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>
-oil and gas extraction	-	-	-	-	-	-	-	-
-electric energy supply industry	-	-	-	-	-	-	-	-
-oil refineries	-	-	70.8	2.85	-	-	-	-
-NGL plant	-	-	-	-	-	-	-	-
<b>Total transformation sector</b>	<b>71.7</b>	<b>2.88</b>	<b>706.1</b>	<b>35.34</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>
-hydro power plants	-	-	-	-	-	-	-	-
-thermal power plants	41.8	1.68	242.2	9.73	-	-	-	-
-public cogeneration plants	29.9	1.20	132.1	5.31	-	-	-	-
-public heating plants	-	-	39.0	1.57	-	-	-	-
-industrial cogeneration plants	-	-	254.5	10.23	-	-	-	-
-industrial heating plants	-	-	38.3	1.54	-	-	-	-
-petroleum refineries	-	-	-	-	-	-	-	-
-NGL-plant	-	-	-	-	-	-	-	-
-gas works	-	-	-	-	-	-	-	-
<b>Non energy use</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>-</b>	<b>0.0</b>	<b>0.0</b>	<b>5.0</b>	<b>0.17</b>
<b>Losses</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>
<b>Final energy demand</b>	<b>13.2</b>	<b>0.58</b>	<b>102.5</b>	<b>4.12</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>
<b>Industry</b>	<b>13.2</b>	<b>0.58</b>	<b>75.8</b>	<b>3.05</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>
-iron and steel	-	-	1.6	0.06	-	-	-	-
-non-ferrous metals	-	-	4.0	0.16	-	-	-	-
-non-metallic minerals	-	-	3.8	0.15	-	-	-	-
-chemical	2.2	0.09	-	-	-	-	-	-
-construction materials	0.2	0.01	52.3	2.10	-	-	-	-
-pulp and paper	-	-	2.6	0.10	-	-	-	-
-food production	10.8	0.43	-	-	-	-	-	-
-not elsewhere specified	-	-	11.5	0.46	-	-	-	-
<b>Other sectors</b>	<b>0.0</b>	<b>0.0</b>	<b>26.7</b>	<b>1.07</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>
-households	-	-	15.4	0.62	-	-	-	-
-services	-	-	6.6	0.27	-	-	-	-
-agriculture	-	-	4.7	0.19	-	-	-	-
-construction	-	-	-	-	-	-	-	-
<b>Transport</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>
-rail	-	-	-	-	-	-	-	-
-road	-	-	-	-	-	-	-	-
-air	-	-	-	-	-	-	-	-
-sea and river	-	-	-	-	-	-	-	-
-public city	-	-	-	-	-	-	-	-

Table A3-3: National energy balance for 2005 (continue)

	Bitumen		Lubricants		Paraffin and wax		Petroleum coke	
	10 <sup>3</sup> t	PJ	10 <sup>3</sup> t	PJ	10 <sup>3</sup> t	PJ	10 <sup>3</sup> t	PJ
Production	180.7	6.05	60.7	2.03	7.4	0.25	104.2	3.23
Import	105.5	3.53	25.3	0.85	5.6	0.19	172.5	5.35
Export	86.7	2.9	54.7	1.83	2.6	0.09	34.5	1.07
Stock change	4.5	0.15	4.1	0.14	0.6	0.02	0.8	0.02
International marine bunkers	-	-	-	-	-	-	-	-
<b>Energy supplied</b>	<b>204.0</b>	<b>6.83</b>	<b>35.4</b>	<b>1.19</b>	<b>11.0</b>	<b>0.37</b>	<b>243.0</b>	<b>7.53</b>
<b>Energy sector own use</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>64.2</b>	<b>1.99</b>
-oil and gas extraction	-	-	-	-	-	-	-	-
-electric energy supply industry	-	-	-	-	-	-	-	-
-oil refineries	-	-	-	-	-	-	64.2	1.99
-NGL plant	-	-	-	-	-	-	-	-
<b>Total transformation sector</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>6.5</b>	<b>0.20</b>
-hydro power plants	-	-	-	-	-	-	-	-
-thermal power plants	-	-	-	-	-	-	-	-
-public cogeneration plants	-	-	-	-	-	-	-	-
-public heating plants	-	-	-	-	-	-	-	-
-industrial cogeneration plants	-	-	-	-	-	-	6.5	0.20
-industrial heating plants	-	-	-	-	-	-	-	-
-petroleum refineries	-	-	-	-	-	-	-	-
-NGL-plant	-	-	-	-	-	-	-	-
-gas works	-	-	-	-	-	-	-	-
<b>Non energy use</b>	<b>204.0</b>	<b>6.83</b>	<b>35.4</b>	<b>1.19</b>	<b>11.0</b>	<b>0.37</b>	<b>0.0</b>	<b>0.0</b>
<b>Losses</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>
<b>Final energy demand</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>172.3</b>	<b>5.34</b>
<b>Industry</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>172.3</b>	<b>5.34</b>
-iron and steel	-	-	-	-	-	-	-	-
-non-ferrous metals	-	-	-	-	-	-	-	-
-non-metallic minerals	-	-	-	-	-	-	-	-
-chemical	-	-	-	-	-	-	0.7	0.02
-construction materials	-	-	-	-	-	-	171.6	5.32
-pulp and paper	-	-	-	-	-	-	-	-
-food production	-	-	-	-	-	-	-	-
-not elsewhere specified	-	-	-	-	-	-	-	-
<b>Other sectors</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>
-households	-	-	-	-	-	-	-	-
-services	-	-	-	-	-	-	-	-
-agriculture	-	-	-	-	-	-	-	-
-construction	-	-	-	-	-	-	-	-
<b>Transport</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>
-rail	-	-	-	-	-	-	-	-
-road	-	-	-	-	-	-	-	-
-air	-	-	-	-	-	-	-	-
-sea and river	-	-	-	-	-	-	-	-
-public city	-	-	-	-	-	-	-	-

Table A3-3: National energy balance for 2005 (continue)

	Ethane		Refinery gas		Refinery semiproducts		Aditives	
	10 <sup>3</sup> t	PJ	10 <sup>3</sup> t	PJ	10 <sup>3</sup> t	PJ	10 <sup>3</sup> t	PJ
Production	66.1	3.13	263.8	11.71	-	-	-	-
Import	-	-	-	-	287.0	12.17	20.2	0.86
Export	-	-	-	-	-	-	-	-
Stock change	-	-	-	-	-5.0	-0.21	-0.5	-0.02
International marine bunkers	-	-	-	-	-	-	-	-
<b>Energy supplied</b>	<b>66.1</b>	<b>3.13</b>	<b>263.8</b>	<b>11.71</b>	<b>282.0</b>	<b>11.96</b>	<b>19.7</b>	<b>0.84</b>
<b>Energy sector own use</b>	<b>0.0</b>	<b>0.0</b>	<b>241.7</b>	<b>10.78</b>	<b>282.0</b>	<b>11.96</b>	<b>0.0</b>	<b>0.0</b>
-oil and gas extraction	-	-	-	-	-	-	-	-
-electric energy supply industry	-	-	-	-	-	-	-	-
-oil refineries	-	-	241.7	10.78	282.0	11.96	-	-
-NGL plant	-	-	-	-	-	-	-	-
<b>Total transformation sector</b>	<b>0.0</b>	<b>0.0</b>	<b>22.1</b>	<b>0.93</b>	<b>0.00</b>	<b>0.00</b>	<b>19.7</b>	<b>0.84</b>
-hydro power plants	-	-	-	-	-	-	-	-
-thermal power plants	-	-	-	-	-	-	-	-
-public cogeneration plants	-	-	-	-	-	-	-	-
-public heating plants	-	-	-	-	-	-	-	-
-industrial cogeneration plants	-	-	22.1	0.93	-	-	-	-
-industrial heating plants	-	-	-	-	-	-	-	-
-petroleum refineries	-	-	-	-	-	-	19.7	0.84
-NGL-plant	-	-	-	-	-	-	-	-
-gas works	-	-	-	-	-	-	-	-
<b>Non energy use</b>	<b>66.1</b>	<b>3.13</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>
<b>Losses</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>
<b>Final energy demand</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>
<b>Industry</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>
-iron and steel	-	-	-	-	-	-	-	-
-non-ferrous metals	-	-	-	-	-	-	-	-
-non-metallic minerals	-	-	-	-	-	-	-	-
-chemical	-	-	-	-	-	-	-	-
-construction materials	-	-	-	-	-	-	-	-
-pulp and paper	-	-	-	-	-	-	-	-
-food production	-	-	-	-	-	-	-	-
-not elsewhere specified	-	-	-	-	-	-	-	-
<b>Other sectors</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>
-households	-	-	-	-	-	-	-	-
-services	-	-	-	-	-	-	-	-
-agriculture	-	-	-	-	-	-	-	-
-construction	-	-	-	-	-	-	-	-
<b>Transport</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>
-rail	-	-	-	-	-	-	-	-
-road	-	-	-	-	-	-	-	-
-air	-	-	-	-	-	-	-	-
-sea and river	-	-	-	-	-	-	-	-
-public city	-	-	-	-	-	-	-	-

Table A3-3: National energy balance for 2005 (continue)

	Gas works gas		Other products		Electricity		Steam and hot water
	10 <sup>3</sup> m <sup>3</sup>	PJ	10 <sup>3</sup> t	PJ	GWh	PJ	TJ
Production	19547.0	0.42	48.2	1.94	12458.9	44.85	34417.9
Import	-	-	5.7	0.23	8746.4	31.49	-
Export	-	-	12.8	0.51	3633.5	13.08	-
Stock change	-	-	-35.4	-1.42	-	-	-
International marine bunkers	-	-	-	-	-	-	-
<b>Energy supplied</b>	<b>19547.0</b>	<b>0.42</b>	<b>5.7</b>	<b>0.23</b>	<b>17571.8</b>	<b>63.26</b>	<b>34417.9</b>
<b>Energy sector own use</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>1036.1</b>	<b>3.73</b>	<b>9136.3</b>
-oil and gas extraction	-	-	-	-	112.3	0.40	1359.0
-electric energy supply industry	-	-	-	-	32.2	0.12	-
-hydro power plants	-	-	-	-	182.7	0.66	-
-thermal power plants	-	-	-	-	297.0	1.07	-
-public cogeneration plants	-	-	-	-	101.6	0.37	870.8
-oil refineries	-	-	-	-	295.4	1.06	6906.5
-NGL plant	-	-	-	-	13.9	0.05	-
<b>Total transform. production</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>
-hydro power plants	-	-	-	-	-	-	-
-thermal power plants	-	-	-	-	-	-	-
-public cogeneration plants	-	-	-	-	-	-	-
-public heating plants	-	-	-	-	-	-	-
-industrial cogeneration plants	-	-	-	-	-	-	-
-industrial heating plants	-	-	-	-	-	-	-
-gas works	-	-	-	-	-	-	-
-public heating plants	1460.0	0.03	-	-	-	-	-
<b>Total transformation sector</b>	<b>1460.0</b>	<b>0.03</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>
<b>Non energy use</b>	<b>0.0</b>	<b>0.0</b>	<b>5.7</b>	<b>0.23</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>
<b>Losses</b>	<b>862.1</b>	<b>0.02</b>	<b>0.0</b>	<b>0.0</b>	<b>2130.9</b>	<b>7.67</b>	<b>1673.1</b>
<b>Final energy demand</b>	<b>17224.9</b>	<b>0.37</b>	<b>0.0</b>	<b>0.0</b>	<b>14404.8</b>	<b>51.86</b>	<b>23608.5</b>
<b>Industry</b>	<b>3550.9</b>	<b>0.08</b>	<b>0.0</b>	<b>0.0</b>	<b>3270.5</b>	<b>11.77</b>	<b>15072.2</b>
-iron and steel	31.0	0.00	-	-	249.7	0.90	142.4
-non-ferrous metals	-	-	-	-	93.2	0.34	2.0
-non-metallic minerals	923.0	0.02	-	-	130.3	0.47	70.2
-chemical	-	-	-	-	484.3	1.74	4893.3
-construction materials	-	-	-	-	611.0	2.20	29.9
-pulp and paper	31.0	0.00	-	-	249.5	0.90	2278.1
-food production	109.9	0.00	-	-	522.0	1.88	4548.1
-not elsewhere specified	2456.0	0.05	-	-	930.5	3.35	3108.2
<b>Other sectors</b>	<b>13674.0</b>	<b>0.29</b>	<b>0.0</b>	<b>0.0</b>	<b>10830.2</b>	<b>38.99</b>	<b>8536.3</b>
-households	10240.0	0.22	-	-	6333.2	22.80	6875.7
-services	3434.0	0.07	-	-	4182.5	15.06	1660.6
-agriculture	-	-	-	-	66.5	0.24	-
-construction	-	-	-	-	248.0	0.89	-
<b>Transport</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>304.1</b>	<b>1.09</b>	<b>0.0</b>
-rail	-	-	-	-	174.7	0.63	-
-road	-	-	-	-	-	-	-
-air	-	-	-	-	18.8	0.07	-
-sea and river	-	-	-	-	24.2	0.07	-
-public city	-	-	-	-	57.1	0.21	-
-other	-	-	-	-	29.3	0.11	-

## **ANNEX 4**

### **ASSESSMENT OF COMPLETENESS AND (POTENTIAL) SOURCES AND SINKS OF GREENHOUSE GAS EMISSIONS AND REMOVALS EXCLUDED**

Table A4-1 shows source/sink categories of GHGs that are not estimated in the Croatian GHG inventory, and the explanations for those categories being omitted. This table is taken from the CRF Table9s1.

*Table A4-1 GHGs and source/sink categories not considered in the Croatian GHG inventory*

GHG	Sector	Source/sink category	Explanation
CO <sub>2</sub>	2. Industrial Processes	2.A.5 Asphalt Roofing	The IPCC Guidelines do not provide methodologies for calculation of CO <sub>2</sub> emission
CO <sub>2</sub>	2. Industrial Processes	2.A.6 Road Paving with Asphalt	The IPCC Guidelines do not provide methodologies for calculation of CO <sub>2</sub> emission
CO <sub>2</sub>	2. Industrial Processes	2.A.7.1 Glass Production	The IPCC Guidelines do not provide methodologies for calculation of CO <sub>2</sub> emission
CO <sub>2</sub>	2. Industrial Processes	2.B.5.2 Ethylene	The IPCC Guidelines do not provide methodologies for calculation of CO <sub>2</sub> emission
CO <sub>2</sub>	2. Industrial Processes	2.D.2 Food and Drink	CO <sub>2</sub> from Food and Drink Production (e.g. gasification of water) can be of biogenic or non-biogenic origin. Only information on CO <sub>2</sub> emissions of non-biogenic origin should be reported.
CO <sub>2</sub>	2. Industrial Processes	Propylene	The IPCC Guidelines do not provide methodologies for calculation of CO <sub>2</sub> emission.
CO <sub>2</sub>	2. Industrial Processes	Polyvinylchloride	The IPCC Guidelines do not provide methodologies for calculation of CO <sub>2</sub> emission.
CO <sub>2</sub>	2. Industrial Processes	Polystyrene	The IPCC Guidelines do not provide methodologies for calculation of CO <sub>2</sub> emission
CO <sub>2</sub>	2. Industrial Processes	Sulphuric acid production	The IPCC Guidelines do not provide methodologies for calculation of CO <sub>2</sub> emission.
CO <sub>2</sub>	2. Industrial Processes	Polyethene low density	The IPCC Guidelines do not provide methodologies for calculation of CO <sub>2</sub> emission.
CO <sub>2</sub>	5. LULUCF	5.A.1 Forest Land remaining Forest Land	Dificulies in collecting adequate activity data.
CO <sub>2</sub>	5. LULUCF	5.B.1 Cropland remaining Cropland	Dificulties in collecting adequate activity data.
CO <sub>2</sub>	5. LULUCF	5.C.1 Grassland remaining Grassland	Dificulties in collecting adequate activity data.
CO <sub>2</sub>	5. LULUCF	5.D.1 Wetlands remaining Wetlands	Dificulties in collecting adequate activity data.
CO <sub>2</sub>	5. LULUCF	Harvested Wood Products	Dificulties in collecting adequate activity data.
CO <sub>2</sub>	6. Waste	6.A.1 Managed Waste Disposal on Land	IPCC Guidelines do not provide methodology for the calculation of CO <sub>2</sub> emissions from Solid Waste Disposal on Land.
CH <sub>4</sub>	1. Energy	1.B.2.A.1 Exploration	Activity data and emission factors were not available.
CH <sub>4</sub>	1. Energy	1.B.2.B.1 Exploration	Activity data and emission factors were not available.
CH <sub>4</sub>	2. Industrial Processes	2.A.7.1 Glass Production	The IPCC Guidelines do not provide methodologies for calculation of CH <sub>4</sub> emission.
CH <sub>4</sub>	2. Industrial Processes	2.B.1 Ammonia Production	The IPCC Guidelines do not provide methodologies for calculation of CH <sub>4</sub> emission.
CH <sub>4</sub>	2. Industrial Processes	2.C.1.1 Steel	The IPCC Guidelines do not provide methodologies for calculation of CH <sub>4</sub> emission.



*Table A4-1 GHGs and source/sink categories not considered in the Croatian GHG inventory (cont.)*

GHG	Sector	Source/sink category	Explanation
CH <sub>4</sub>	2. Industrial Processes	Propylene	The IPCC Guidelines do not provide methodologies for calculation of CH <sub>4</sub> emission.
CH <sub>4</sub>	2. Industrial Processes	Polyvinylchloride	The IPCC Guidelines do not provide methodologies for calculation of CH <sub>4</sub> emission.
CH <sub>4</sub>	2. Industrial Processes	Polystyrene	The IPCC Guidelines do not provide methodologies for calculation of CH <sub>4</sub> emission.
CH <sub>4</sub>	2. Industrial Processes	Sulphuric acid production	The IPCC Guidelines do not provide methodologies for calculation of CH <sub>4</sub> emission.
CH <sub>4</sub>	2. Industrial Processes	Polyethene low density	The IPCC Guidelines do not provide methodologies for calculation of CH <sub>4</sub> emission.
CH <sub>4</sub>	4. Agriculture	4.A Enteric Fermentation	No data available.
CH <sub>4</sub>	5. LULUCF	5.A.1 Forest Land remaining Forest Land	Difficulties in collecting adequate activity data.
CH <sub>4</sub>	5. LULUCF	5.B.1 Cropland remaining Cropland	Difficulties in collecting adequate activity data.
CH <sub>4</sub>	5. LULUCF	5.C.1 Grassland remaining Grassland	Difficulties in collecting adequate activity data.
CH <sub>4</sub>	5. LULUCF	5.D.1 Wetlands remaining Wetlands	Difficulties in collecting adequate activity data.
CH <sub>4</sub>	5. LULUCF	5.E.1 Settlements remaining Settlements	Difficulties in collecting adequate activity data.
CH <sub>4</sub>	5. LULUCF	5.E.2 Land converted to Settlements	Difficulties in collecting adequate activity data.
CH <sub>4</sub>	5. LULUCF	5.F.1 Other Land remaining Other Land	Difficulties in collecting adequate activity data.
CH <sub>4</sub>	5. LULUCF	Harvested Wood Products	Difficulties in collecting adequate activity data.
CH <sub>4</sub>	6. Waste	6.B.2.1 Domestic and Commercial (w/o human sewage)	CH <sub>4</sub> emissions from Wastewater Handling (Domestic and Commercial Wastewater) are not estimated because activity data are not available.
CH <sub>4</sub>	6. Waste	Incineration of hospital wastes	IPCC Guidelines do not provide default emission factor for CH <sub>4</sub> emission calculation from Incineration of clinical waste. There is no national information on these data. Information about type of incineration/technology is lacking.
N <sub>2</sub> O	2. Industrial Processes	2.A.7.1 Glass Production	The IPCC Guidelines do not provide methodologies for calculation of N <sub>2</sub> O emission
N <sub>2</sub> O	2. Industrial Processes	2.B.1 Ammonia Production	The IPCC Guidelines do not provide methodologies for calculation of N <sub>2</sub> O emission
N <sub>2</sub> O	2. Industrial Processes	2.B.5.2 Ethylene	The IPCC Guidelines do not provide methodologies for calculation of N <sub>2</sub> O emission
N <sub>2</sub> O	2. Industrial Processes	Propylene	The IPCC Guidelines do not provide methodologies for calculation of N <sub>2</sub> O emission
N <sub>2</sub> O	2. Industrial Processes	Polyvinylchloride	The IPCC Guidelines do not provide methodologies for calculation of N <sub>2</sub> O emission
N <sub>2</sub> O	2. Industrial Processes	Polystyrene	The IPCC Guidelines do not provide methodologies for calculation of N <sub>2</sub> O emission
N <sub>2</sub> O	2. Industrial Processes	Sulphuric acid production	The IPCC Guidelines do not provide methodologies for calculation of N <sub>2</sub> O emission
N <sub>2</sub> O	2. Industrial Processes	Polyethene low density	The IPCC Guidelines do not provide methodologies for calculation of N <sub>2</sub> O emission

**Table A4-1 GHGs and source/sink categories not considered in the Croatian GHG inventory (cont.)**

GHG	Sector	Source/sink category	Explanation
N <sub>2</sub> O	3. Solvent and Other Product Use	3.B Degreasing and Dry Cleaning	N <sub>2</sub> O emissions from medical uses and other possible sources are not estimated because activity data are not available.
N <sub>2</sub> O	3. Solvent and Other Product Use	3.D.1 Use of N <sub>2</sub> O for Anaesthesia	N <sub>2</sub> O emissions from medical uses and other possible sources are not estimated because activity data are not available.
N <sub>2</sub> O	3. Solvent and Other Product Use	3.D.2 Fire Extinguishers	N <sub>2</sub> O emissions from medical uses and other possible sources are not estimated because activity data are not available.
N <sub>2</sub> O	3. Solvent and Other Product Use	3.D.3 N <sub>2</sub> O from Aerosol Cans	N <sub>2</sub> O emissions from medical uses and other possible sources are not estimated because activity data are not available.
N <sub>2</sub> O	3. Solvent and Other Product Use	3.D.4 Other Use of N <sub>2</sub> O	N <sub>2</sub> O emissions from medical uses and other possible sources are not estimated because activity data are not available.
N <sub>2</sub> O	5. LULUCF	5.A.1 Forest Land remaining Forest Land	Difficulties in collecting adequate activity data.
N <sub>2</sub> O	5. LULUCF	5.A.2 Land converted to Forest Land	Difficulties in collecting adequate activity data.
N <sub>2</sub> O	5. LULUCF	5.B.1 Cropland remaining Cropland	Difficulties in collecting adequate activity data.
N <sub>2</sub> O	5. LULUCF	5.B.2.1 Forest Land converted to Cropland	Difficulties in collecting adequate activity data.
N <sub>2</sub> O	5. LULUCF	5.B.2.2 Grassland converted to Cropland	Difficulties in collecting adequate activity data.
N <sub>2</sub> O	5. LULUCF	5.B.2.3 Wetlands converted to Cropland	Difficulties in collecting adequate activity data.
N <sub>2</sub> O	5. LULUCF	5.B.2.5 Other Land converted to Cropland	Difficulties in collecting adequate activity data.
N <sub>2</sub> O	5. LULUCF	5.C.1 Grassland remaining Grassland	Difficulties in collecting adequate activity data.
N <sub>2</sub> O	5. LULUCF	5.D.1 Wetlands remaining Wetlands	Difficulties in collecting adequate activity data.
N <sub>2</sub> O	5. LULUCF	5.E.1 Settlements remaining Settlements	Difficulties in collecting adequate activity data.
N <sub>2</sub> O	5. LULUCF	5.E.2 Land converted to Settlements	Difficulties in collecting adequate activity data.
N <sub>2</sub> O	5. LULUCF	Harvested Wood Products	Difficulties in collecting adequate activity data.
N <sub>2</sub> O	6. Waste	6.B.2.1 Domestic and Commercial (w/o human sewage)	Activity data are not available.
N <sub>2</sub> O	6. Waste	6.B.2.1 Domestic and Commercial (w/o human sewage)	IPCC Guidelines do not provide methodology for the calculation of N <sub>2</sub> O emission from Domestic Sludge.
N <sub>2</sub> O	6. Waste	Incineration of hospital wastes	IPCC Guidelines do not provide default emission factor for N <sub>2</sub> O emission calculation from Incineration of clinical waste. There is no national information on these data. Information about type of incineration/technology is lacking.
HFCs	2. Industrial Processes	2.F.2 Foam Blowing	Total potential emissions were not estimated because input data for those emissions were not available.
HFCs	2. Industrial Processes	2.F.3 Fire Extinguishers	Total potential emissions were not estimated because input data for those emissions were not available.
HFCs	2. Industrial Processes	2.F.4 Aerosols/ Metered Dose Inhalers	Total potential emissions were not estimated because input data for those emissions were not available.

*Table A4-1 GHGs and source/sink categories not considered in the Croatian GHG inventory (cont.)*

GHG	Sector	Source/sink category	Explanation
HFCs	2. Industrial Processes	2.F.5 Solvents	Total potential emissions were not estimated because input data for those emissions were not available.
SF <sub>6</sub>	2. Industrial Processes	2.F.8 Electrical Equipment	Total potential emissions of SF <sub>6</sub> are not estimated because data are not available.
SF <sub>6</sub>	2. Industrial Processes	2.F.8 Electrical Equipment	Croatia does not yet have the necessary data to calculate actual emissions. According to requirement of Regulation of the greenhouse gases emissions monitoring in the Republic of Croatia (Official gazette 1/07), which will go into force in 2008, each source of SF <sub>6</sub> emissions should report required activity data for more accurate emissions estimation.
SF <sub>6</sub>	2. Industrial Processes	2.F.P2.1 In bulk	The potential emissions of SF <sub>6</sub> are not estimated because data are not available.
SF <sub>6</sub>	2. Industrial Processes	2.F.P3.1 In bulk	The potential emissions of SF <sub>6</sub> are not estimated because data are not available.
Carbon	5. LULUCF	5.A.2.2 Grassland converted to Forest Land	Difficulties in collecting adequate activity data.
Carbon	5. LULUCF	5.A.2.5 Other Land converted to Forest Land	Difficulties in collecting adequate activity data.
Carbon	5. LULUCF	5.B.1 Cropland remaining Cropland	Difficulties in collecting adequate activity data.
Carbon	5. LULUCF	5.B.2.1 Forest Land converted to Cropland	Difficulties in collecting adequate activity data.
Carbon	5. LULUCF	5.B.2.2 Grassland converted to Cropland	Difficulties in collecting adequate activity data.
Carbon	5. LULUCF	5.B.2.3 Wetlands converted to Cropland	Difficulties in collecting adequate activity data.
Carbon	5. LULUCF	5.B.2.5 Other Land converted to Cropland	Difficulties in collecting adequate activity data.
Carbon	5. LULUCF	5.C.1 Grassland remaining Grassland	Difficulties in collecting adequate activity data.
Carbon	5. LULUCF	5.C.2.1 Forest Land converted to Grassland	Difficulties in collecting adequate activity data.
Carbon	5. LULUCF	5.C.2.2 Cropland converted to Grassland	Difficulties in collecting adequate activity data.
Carbon	5. LULUCF	5.C.2.3 Wetlands converted to Grassland	Difficulties in collecting adequate activity data.
Carbon	5. LULUCF	5.C.2.5 Other Land converted to Grassland	Difficulties in collecting adequate activity data.
Carbon	5. LULUCF	5.D.1 Wetlands remaining Wetlands	Difficulties in collecting adequate activity data.
Carbon	5. LULUCF	5.E.1 Settlements remaining Settlements	Difficulties in collecting adequate activity data.
Carbon	5. LULUCF	5.E.2.1 Forest Land converted to Settlements	Difficulties in collecting adequate activity data.
Carbon	5. LULUCF	5.E.2.2 Cropland converted to Settlements	Difficulties in collecting adequate activity data.
Carbon	5. LULUCF	5.E.2.3 Grassland converted to Settlements	Difficulties in collecting adequate activity data.
Carbon	5. LULUCF	5.E.2.4 Wetlands converted to Settlements	Difficulties in collecting adequate activity data.

*Table A4-1 GHGs and source/sink categories not considered in the Croatian GHG inventory (cont.)*

GHG	Sector	Source/sink category	Explanation
Carbon	5. LULUCF	5.E.2.5 Other Land converted to Settlements	Dificulties in collecting adequate activity data.
Carbon	5. LULUCF	5.F.2.1 Forest Land converted to Other Land	Dificulties in collecting adequate activity data.
Carbon	5. LULUCF	5.F.2.2 Cropland converted to Other Land	Dificulties in collecting adequate activity data.
Carbon	5. LULUCF	5.F.2.3 Grassland converted to Other Land	Dificulties in collecting adequate activity data.
Carbon	5. LULUCF	5.F.2.4 Wetlands converted to Other Land	Dificulties in collecting adequate activity data.
Carbon	5. LULUCF	5.F.2.5 Settlements converted to Other Land	Dificulties in collecting adequate activity data.
Carbon	5. LULUCF	5.F.2.1 Forest Land converted to Other Land	Dificulties in collecting adequate activity data.

## **ANNEX 5**

### **TABLE 6.1 OF THE IPCC GOOD PRACTICE GUIDANCE**

Table A5-1: Tier 1 Uncertainty Calculation and Reporting – excluding LULUCF (Table 6.1 – IPCC Good Practice Guidance)

A	B	C	D	E	F	G	H	I	J	K	L	M
	GHG	Base year emissions 1990	Year t emissions 2005	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor uncertainty	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty introduced into the trend in total national emissions
		Gg CO <sub>2</sub> equivalent	Gg CO <sub>2</sub> equivalent	%	%	%	%	%	%	%	%	%
1A	CO <sub>2</sub> Emissions from Stationary Combustion : Coal	3141.5	2681.0	5	5	7.07	0.62	-0.01	0.08	-0.06	0.42	0.43
1A	CO <sub>2</sub> Emissions from Stationary Combustion: Oil	8424.1	6505.1	5	5	7.07	1.51	-0.05	0.21	-0.26	1.03	1.06
1A	CO <sub>2</sub> Emissions from Stationary Combustion: Gas	4053.8	4866.2	5	5	7.07	1.13	0.03	0.15	0.15	0.77	0.79
1A	Mobile Combustion : Road Vehicles	3504.0	5286.9	5	5	7.07	1.23	0.06	0.17	0.30	0.84	0.89
1A	Mobile Combustion: Water-borne Navigation	133.0	99.6	5	5	7.07	0.02	0.00	0.00	0.00	0.02	0.02
1A	Mobile Combustion: Aircraft	295.6	172.6	5	5	7.07	0.04	-0.00358	0.01	-0.02	0.03	0.03
1A	Mobile Combustion: Railways	137.5	95.5	5	5	7.07	0.02	-0.00118	0.003	-0.01	0.02	0.02
1A	Mobile Combustion : Agriculture/Forestry/Fishing	839.2	709.4	5	5	7.07	0.16	-0.00321	0.02	-0.02	0.11	0.11
1B	CO <sub>2</sub> Emissions from Natural Gas Scrubbing	415.9	691.0	10	3	10.44	0.24	0.00916	0.02	0.03	0.22	0.22
2A	CO <sub>2</sub> Emissions from Cement Production	1047.5	1495.0	3	3	4.24	0.21	0.02	0.05	0.05	0.14	0.15
2A	CO <sub>2</sub> Emissions from Lime Production	159.8	196.4	7.5	15	16.77	0.11	0.00	0.01	0.02	0.05	0.05
2A	CO <sub>2</sub> Emissions from Limestone and Dolomite Use	43.2	12.1	7.5	30	30.92	0.01	0.00	0.00	-0.03	0.00	0.03
2A	CO <sub>2</sub> Emissions from Soda Ash Production and Use	25.7	17.2	7.5	30	30.92	0.02	-0.0002	0.0005	-0.0073	0.0041	0.0083
2B	CO <sub>2</sub> Emissions from Ammonia Production	491.6	511.2	3	5	5.83	0.10	0.0012	0.0162	0.0058	0.0486	0.0489
2C	CO <sub>2</sub> Emissions from Iron and Steel Production	0.9	0.3	7.5	30	30.92	0.0003	0.0000	0.0000	-0.0005	0.0001	0.0005
2C	CO <sub>2</sub> Emissions from Ferroalloys Production	194.5	-	7.5	30	30.92	0.0000	-0.0060	0.0000	-0.1787	0.0000	0.1787
2C	CO <sub>2</sub> Emissions from Aluminium Production	111.4	-	3	30	30.15	0.0000	-0.0034	0.0000	-0.1023	0.0000	0.1023
2G	CO <sub>2</sub> Emissions from Other non-specified NEU	208.1	72.2	5	50	50.25	0.12	-0.004082	0.002288	-0.204087	0.011440	0.204407
3	CO <sub>2</sub> Emissions from Solvent and Other Product Use	80.2	155.1	50	50	70.71	0.36	0.002458	0.004914	0.122910	0.245707	0.274734
6C	CO <sub>2</sub> Emissions from Waste Incineration	-	0.0	10	30	31.62	0.000035	0.000001	0.000001	0.000032	0.000011	0.000034
	<b>CO<sub>2</sub> Total</b>	<b>23307.5</b>	<b>23566.9</b>									
1A	CH <sub>4</sub> Emissions from Fuel Combustion: Stationary Sources	172.2	110.0	5	20	20.62	0.07	-0.0018	0.0035	-0.0357	0.0174	0.0397
1A	Mobile Combustion : Road Vehicles	29.9	32.6	5	40	40.31	0.04	0.0001	0.0010	0.0048	0.0052	0.0070
1A	Mobile Combustion: Water-borne Navigation	0.2	0.1	5	40	40.31	0.0002	0.0000	0.0000	-0.0001	0.0000	0.0001
1A	Mobile Combustion: Aircraft	0.0	0.0	5	40	40.31	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000

Table A5-1: Tier 1 Uncertainty Calculation and Reporting – excluding LULUCF (Table 6.1 – IPCC Good Practice Guidance) (cont.)

A	B	C	D	E	F	G	H	I	J	K	L	M
IPCC Source Category												
	GHG	Base year emissions 1990	Year t emissions 2005	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor uncertainty	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty introduced into the trend in total national emissions
		Gg CO <sub>2</sub> equivalent	Gg CO <sub>2</sub> equivalent	%	%	%	%	%	%	%	%	%
1A	Mobile Combustion: Railways	0.2	0.1	5	40	40.31	0.0002	0.0000	0.0000	-0.0001	0.0000	0.0001
1A	Mobile Combustion: Agriculture/Forestry/Fishing	1.3	1.1	5	40	40.31	0.0014	0.0000	0.0000	-0.0002	0.0002	0.0003
1B	Fugitive Emissions from Coal Mining and Handling	48.8	-	5	250	250.05	0.0000	-0.0015	0.0000	-0.3732	0.0000	0.3732
1B	Fugitive Emissions from Oil and Gas Operations	1186.3	1333.9	5	300	300.04	13.1307	0.0060	0.0423	1.7863	0.2114	1.7988
2B	Production of Other Chemicals	15.8	5.3	7.5	30	30.92	0.0054	-0.0003	0.0002	-0.0094	0.0013	0.0095
4A	CH <sub>4</sub> Emissions from Enteric Fermentation in Domestic Livestock	1343.9	822.4	30	40	50.00	1.3491	-0.0151	0.0261	-0.6029	0.7820	0.9874
4B	CH <sub>4</sub> Emissions from Manure Management	227.4	157.1	30	40	50.00	0.2577	-0.0020	0.0050	-0.0794	0.1494	0.1691
6A	CH <sub>4</sub> Emissions from Solid Waste Disposal Sites	221.2	489.5	50	50	70.71	1.0892	0.0081	0.0149	0.4053	0.7440	0.8473
6B	CH <sub>4</sub> Emissions from Waste Water Handling	-	24.0	10	30	31.62	0.0334	0.0009	0.0009	0.0284	0.0095	0.0300
	<b>CH<sub>4</sub> Total</b>	<b>3247.1</b>	<b>2956.2</b>									
1A	N <sub>2</sub> O Emissions from Fuel Combustion : Stationary Sources	65.5	51.0	5	200	200.06	0.3350	-0.0004	0.0016	-0.0773	0.0081	0.0777
1A	Mobile Combustion : Road Vehicles	49.2	165.6	5	200	200.06	1.0871	0.0037	0.0052	0.7488	0.0262	0.7493
1A	Mobile Combustion: Water-borne Navigation	0.3	0.3	5	200	200.06	0.0017	0.0000	0.0000	-0.0005	0.0000	0.0005
1A	Mobile Combustion: Aircraft	2.6	1.5	5	200	200.06	0.0099	0.0000	0.0000	-0.0063	0.0002	0.0063
1A	Mobile Combustion: Railways	0.4	0.2	5	200	200.06	0.0016	0.0000	0.0000	-0.0009	0.0000	0.0009
1A	Mobile Combustion: Agriculture/Forestry/Fishing	2.0	1.7	5	200	200.06	0.0113	0.0000	0.0001	-0.0016	0.0003	0.0016
2B	N <sub>2</sub> O Emissions from Nitric Acid Production	927.6	783.3	3	30	30.15	0.7748	-0.0036	0.0248	-0.1072	0.0745	0.1305
4B	N <sub>2</sub> O Emissions from Manure Management	376.7	236.7	30	60	67.08	0.5209	-0.0040	0.0075	-0.2420	0.2250	0.3304
4B	Direct N <sub>2</sub> O Emissions from Agricultural Soils	1347.6	1238.3	30	40	50.00	2.0312	-0.0020	0.0392	-0.0806	1.1773	1.1801
4D	N <sub>2</sub> O Emissions from Pasture, Range and Paddock Manure	251.8	223.5	30	40	50.00	0.3666	-0.0006	0.0071	-0.0251	0.2125	0.2139
4F	Indirect N <sub>2</sub> O Emissions from Nitrogen Used in Agriculture	916.6	816.9	30	60	67.08	1.7979	-0.0022	0.0259	-0.1304	0.7767	0.7876
6B	N <sub>2</sub> O Emissions from Waste Water Handling	77.1	89.3	10	30	31.62	0.0927	0.0005	0.0028	0.0141	0.0283	0.0316
	<b>N<sub>2</sub>O Total</b>	<b>4017.4</b>	<b>3608.3</b>									

Table A5-1: Tier 1 Uncertainty Calculation and Reporting – excluding LULUCF (Table 6.1 – IPCC Good Practice Guidance) (cont.)

	A	B	C	D	E	F	G	H	I	J	K	L	M
	IPCC Source Category	GHG	Base year emissions 1990	Year t emissions 2005	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor uncertainty	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty introduced into the trend in total national emissions
			Gg CO <sub>2</sub> equivalent	Gg CO <sub>2</sub> equivalent	%	%	%	%	%	%	%	%	%
2C	PFC Emissions from Aluminium production	PFC	936.6		30	50	58.31	0.0000	-0.0287	0.0000	-1.4333	0.0000	1.4333
2F	HFC Emissions from Consumption of HFCs	HFC	43.4	349.1	70	70	98.99	1.1337	0.0097	0.0111	0.6815	0.7745	1.0316
		HFC/PFC/SF <sub>6</sub> Total	979.9	349.1									
	Total GHG Emissions	CO <sub>2</sub> -eq	31551.9	30480.6									
	Total Uncertainties (Level/Trend)							13.66					3.34



Table A5-2: Tier 1 Uncertainty Calculation and Reporting – including LULUCF (Table 6.1 – IPCC Good Practice Guidance)

A	B	C	D	E	F	G	H	I	J	K	L	M
	GHG	Base year emissions 1990	Year t emissions 2005	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor uncertainty	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty introduced into the trend in total national emissions
		Gg CO <sub>2</sub> equivalent	Gg CO <sub>2</sub> equivalent	%	%	%	%	%	%	%	%	%
1A	CO <sub>2</sub> Emissions from Stationary Combustion: Coal	3141.5	2681.0	5	5	7.07	0.50	-0.01	0.07	-0.07	0.35	0.36
1A	CO <sub>2</sub> Emissions from Stationary Combustion: Oil	8424.1	6505.1	5	5	7.07	1.20	-0.05	0.17	-0.27	0.86	0.90
1A	CO <sub>2</sub> Emissions from Stationary Combustion: Gas	4053.8	4866.2	5	5	7.07	0.90	0.02	0.13	0.10	0.64	0.65
1A	Mobile Combustion: Road Vehicles	3504.0	5286.9	5	5	7.07	0.98	0.05	0.14	0.23	0.70	0.74
1A	Mobile Combustion: Water-borne Navigation	133.0	99.6	5	5	7.07	0.02	0.00	0.00	0.00	0.01	0.01
1A	Mobile Combustion: Aircraft	295.6	172.6	5	5	7.07	0.03	0.00	0.00	-0.02	0.02	0.03
1A	Mobile Combustion: Railways	137.5	95.5	5	5	7.07	0.02	0.00	0.00	-0.01	0.01	0.01
1A	Mobile Combustion: Agriculture/Forestry/Fishing	839.2	709.4	5	5	7.07	0.13	0.00	0.02	-0.02	0.09	0.10
1B	CO <sub>2</sub> Emissions from Natural Gas Scrubbing	415.9	691.0	10	3	10.44	0.19	0.01	0.02	0.02	0.18	0.18
2A	CO <sub>2</sub> Emissions from Cement Production	1047.5	1495.0	3	3	4.24	0.17	0.01	0.04	0.03	0.12	0.12
2A	CO <sub>2</sub> Emissions from Lime Production	159.8	196.4	7.5	15	16.77	0.09	0.00	0.01	0.01	0.04	0.04
2A	CO <sub>2</sub> Emissions from Limestone and Dolomite Use	43.2	12.1	7.5	30	30.92	0.01	0.00	0.00	-0.03	0.00	0.03
2A	CO <sub>2</sub> Emissions from Soda Ash Production and Use	25.7	17.2	7.5	30	30.92	0.0139	-0.0002	0.0005	-0.0070	0.0034	0.0078
2C	CO <sub>2</sub> Emissions from Iron and Steel Production	0.9	0.3	7.5	30	30.92	0.0003	0.0000	0.0000	-0.0004	0.0001	0.0004
2B	CO <sub>2</sub> Emissions from Ammonia Production	491.6	511.2	3	5	5.83	0.0779	0.0004	0.0135	0.0019	0.0405	0.0406
2C	CO <sub>2</sub> Emissions from Ferroalloys Production	194.5	-	7.5	30	30.92	0.0000	-0.0052	0.0000	-0.1560	0.0000	0.1560
2C	CO <sub>2</sub> Emissions from Aluminium Production	111.4	-	3	30	30.15	0.0000	-0.0030	0.0000	-0.0893	0.0000	0.0893
2G	CO <sub>2</sub> Emissions from Other non-specified NEU	208.1	72.2	5	50	50.25	0.094814	-0.003653	0.001908	-0.182640	0.009541	0.182889
3	CO <sub>2</sub> Emissions from Solvent and Other Product Use	80.2	155.1	50	50	70.71	0.286562	0.001954	0.004098	0.097710	0.204914	0.227017
5A	Forest land remaining forest land	-6281.2	-7779.0	45	30	54.08	10.9963	0.0377	0.2056	1.1296	9.2526	9.3213
6C	Emissions from Waste Incineration	-	0.0	10	30	31.62	0.000028	0.000001	0.000001	0.000027	0.000009	0.000028
	<b>CO<sub>2</sub> Total</b>	<b>17026.2</b>	<b>15788.0</b>									
1A	CH <sub>4</sub> Emissions from Fuel Combustion: Stationary Sources	172.2	110.0	5	20	20.62	0.0999	-0.0018	0.0044	-0.0354	0.0218	0.0415
1A	Mobile Combustion: Road Vehicles	29.9	32.6	5	40	40.31	0.0579	0.0002	0.0013	0.0091	0.0065	0.0112
1A	Mobile Combustion: Water-borne Navigation	0.2	0.1	5	40	40.31	0.0003	0.0000	0.0000	0.0000	0.0000	0.0001

Table A5-2: Tier 1 Uncertainty Calculation and Reporting – including LULUCF (Table 6.1 – IPCC Good Practice Guidance) (cont.)

A	B	C	D	E	F	G	H	I	J	K	L	M
	GHG	Base year emissions 1990	Year t emissions 2005	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor uncertainty	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty introduced into the trend in total national emissions
		Gg CO <sub>2</sub> equivalent	Gg CO <sub>2</sub> equivalent	%	%	%	%	%	%	%	%	%
IPCC Source Category												
1A	Mobile Combustion: Aircraft	0.0	0.0	5	40	40.31	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
1A	Mobile Combustion: Railways	0.2	0.1	5	40	40.31	0.0002	0.0000	0.0000	-0.0001	0.0000	0.0001
1A	Mobile Combustion: Agriculture/Forestry/Fishing	1.3	1.1	5	40	40.31	0.0019	0.0000	0.0000	-0.0001	0.0002	0.0003
1B	Fugitive Emissions from Coal Mining and Handling	48.8	-	5	250	250.05	0.0000	-0.0017	0.0000	-0.4333	0.0000	0.4333
1B	Fugitive Emissions from Oil and Gas Operations	1186.3	1333.9	5	300	300.04	17.6301	0.0106	0.0528	3.1832	0.2639	3.1941
2B	Production of Other Chemicals	15.8	5.3	7.5	30	30.92	0.0073	-0.0004	0.0002	-0.0105	0.0016	0.0106
4A	CH <sub>4</sub> Emissions from Enteric Fermentation in Domestic Livestock	1343.9	822.4	30	40	50.00	1.8114	-0.0152	0.0325	-0.6088	0.9763	1.1506
4B	CH <sub>4</sub> Emissions from Manure Management	227.4	157.1	30	40	50.00	0.3460	-0.0019	0.0062	-0.0747	0.1865	0.2009
6A	CH <sub>4</sub> Emissions from Solid Waste Disposal Sites	221.2	469.5	50	50	70.71	1.4824	0.0107	0.0186	0.5357	0.9289	1.0723
6B	CH <sub>4</sub> Emissions from Waste Water Handling	-	24.0	10	30	31.62	0.0334	0.0009	0.0009	0.0284	0.0095	0.0300
	<b>CH<sub>4</sub> Total</b>	<b>3247.1</b>	<b>2956.2</b>									
1A	N <sub>2</sub> O Emissions from Fuel Combustion : Stationary Sources	65.5	51.0	5	200	200.06	0.4499	-0.0003	0.0020	-0.0615	0.0101	0.0623
1A	Mobile Combustion - Road Vehicles	49.2	165.6	5	200	200.06	1.4596	0.0048	0.0066	0.9613	0.0328	0.9619
1A	Mobile Combustion: Water-borne Navigation	0.3	0.3	5	200	200.06	0.0022	0.0000	0.0000	-0.0004	0.0000	0.0004
1A	Mobile Combustion: Aircraft	2.6	1.5	5	200	200.06	0.0133	0.0000	0.0001	-0.0064	0.0003	0.0064
1A	Mobile Combustion: Railways	0.4	0.2	5	200	200.06	0.0021	0.0000	0.0000	-0.0009	0.0000	0.0009
1A	Mobile Combustion - Agriculture/Forestry/Fishing	2.0	1.7	5	200	200.06	0.0151	0.0000	0.0001	-0.0009	0.0003	0.0010
2B	N <sub>2</sub> O Emissions from Nitric Acid Production	927.6	783.3	3	30	30.15	1.0403	-0.0020	0.0310	-0.0593	0.0930	0.1103
4B	N <sub>2</sub> O Emissions from Manure Management	376.7	236.7	30	60	67.08	0.6993	-0.0040	0.0094	-0.2415	0.2810	0.3705
4B	Direct N <sub>2</sub> O Emissions from Agricultural Soils	1347.6	1238.3	30	40	50.00	2.7272	0.0011	0.0490	0.0437	1.4700	1.4706
4D	N <sub>2</sub> O Emissions from Pasture, Range and Paddock Manure	251.8	223.5	30	40	50.00	0.4922	-0.0001	0.0088	-0.0043	0.2653	0.2653
4F	Indirect N <sub>2</sub> O Emissions from Nitrogen Used in Agriculture	916.6	816.9	30	60	67.08	2.4140	-0.0003	0.0323	-0.0155	0.9698	0.9699
6B	N <sub>2</sub> O Emissions from Waste Water Handling	77.1	89.3	10	30	31.62	0.1244	0.0008	0.0035	0.0238	0.0353	0.0426
	<b>N<sub>2</sub>O Total</b>	<b>4017.4</b>	<b>3608.3</b>									

Table A5-2: Tier 1 Uncertainty Calculation and Reporting – including LULUCF (Table 6.1 – IPCC Good Practice Guidance) (cont.)

A	B	C	D	E	F	G	H	I	J	K	L	M
	GHG	Base year emissions 1990	Year t emissions 2005	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor uncertainty	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty introduced into the trend in total national emissions
		Gg CO <sub>2</sub> equivalent	Gg CO <sub>2</sub> equivalent	%	%	%	%	%	%	%	%	%
2C	PFO Emissions from Aluminium production	936.6	-	30	50	58.31	0.0000	-0.0333	0.0000	-1.6641	0.0000	1.6641
2F	HFC Emissions from Consumption of HFCs	43.4	349.1	70	70	98.99	1.5222	0.0123	0.0138	0.8590	0.9670	1.2934
	HFC/PFC/SF <sub>6</sub> Total	979.9	349.1									
	CO <sub>2</sub> eq	25270.7	22701.6									
	Total GHG Emissions											
	Total Uncertainties (Level/Trend)						15.47					9.68

## **ANNEX 6**

### **QA/QC FRAMEWORK PLAN**

Table A6-1: The framework of the Croatian QA/QC Plan

Item	Check/Review			Corrective actions		Comments
	Individual (first initial, last name)	Delivery date	Date of performance	Individual (first initial, last name)	Final date	
A. DATA GATHERING, INPUT, AND HANDLING ACTIVITIES: QUALITY CHECKS						
1. Check a sample of input data for transcription errors (Energy)						
2. Check a sample of input data for transcription errors (Industry+Solvents)						
3. Check a sample of input data for transcription errors (Agriculture+LULUCF)						
4. Check a sample of input data for transcription errors (Waste)						
5. Check a sample of input data for transcription errors (All sources)						
6. Identify spreadsheet modifications that could provide additional controls or checks on quality (Energy-Fugitive emission)						
7. Identify spreadsheet modifications that could provide additional controls or checks on quality (Energy)						
B. DATA DOCUMENTATION: QUALITY CHECKS						
8 Check project file for completeness						
9. Confirm that bibliographical data references are included (in spreadsheet) for every primary data element						
10. Check that all appropriate citations from the spreadsheets appear in the Inventory Report						
11. Check that all citations in spreadsheets and inventory are complete (i.e.. include all relevant information)						
12. Check that assumptions and criteria for selection of activity data and emission factors are documented (Waste)						
13. Check that changes in data or methodology are documented (Energy-Fugitive)						
14. Check that changes in data or methodology are documented (Industries)						
15. Check that changes in data or methodology are documented (Waste)						
C. CALCULATING EMISSIONS AND CHECKING CALCULATIONS						
16. Check that all emission calculations are included (i.e.. emissions are not hard-wired)						
17. Check whether emission units, parameters, and conversion factors are inappropriately hardwired						
18. Check if units are properly labeled and correctly carried through from beginning to end of calculation						
19. Check that conversion factors are correct						
20. Check that temporal and spatial adjustment factors are used correctly						
21. Check the data relationships (comparability) and data processing steps (e.g.. equations) in the spreadsheets						
22. Check a representative sample of calculations, by hand or electronically						
23. Check some calculations with abbreviated calculations						
24. Check the aggregation of data within a source category						
25. When methods or data have changed, check consistency of time series inputs and calculations						
26. Check for consistency with IPCC inventory guidelines and good practices, particularly if changes occur						

Table A6-2: An example of Inventory Data Record Sheet for 2005 in Waste

**INVENTORY DATA RECORD SHEET****Year: 2005**

<b>MODULE:</b> WASTE	
<b>SUBMODULE:</b> METHANE EMISSIONS FROM SOLID WASTE DISPOSAL SITES	
<b>WORKSHEET:</b> 6-1	<b>SHEET:</b> 1 OF 1 CH <sub>4</sub> EMISSIONS
<b>STEP:</b> 1 TO 4	<b>PAGE:</b> 1 of 2
<b>DIRECT DATA SOURCE:</b> <b>A. ACTIVITY DATA:</b> Extrapolation of inappropriate activity data on quantities of MSW disposed to different types of SWDs. Quantities of MSW disposed to SWDSs: Managed: 100.96 Gg Unmanaged – deep: 969.83 Gg Unmanaged – shallow: 215.35 Gg Country-specific methane correction factor (MCF): 0.749 Country-specific fraction of degradable organic carbon (DOC): 0.16 Recovered methane: 0 Gg  <b>B. METHODOLOGY/EMISSION FACTOR:</b> Publications: IPCC/UNEP/OECD/IEA (1997), <i>Greenhouse Gas Inventory Workbook</i> , Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories, Volume 2. IPCC/UNEP/OECD/IEA (1997), <i>Greenhouse Gas Inventory Reference Manual</i> , Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories, Volume 3. IPCC (2000), <i>Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories</i> Methodology: First Order Decay method (Tier 2) Methane generation rate constant k=0.05 Fraction of DOC which really degrades: 0.55 (0.5-0.6) Fraction of carbon released as methane: 0.5	
<b>ORIGINAL DATA SOURCE:</b> <b>A. ACTIVITY DATA:</b> Fundurulja, D., Mužinić, M. (2000) <i>Estimation of the Quantities of Municipal Solid Waste in the Republic of Croatia in the period 1990 – 1998 and 1998 – 2010</i> . Ministry of Environmental Protection, Physical Planning and Construction (2007) <i>Waste Management Plan in the Republic of Croatia (2007-2015)</i>	
<b>METHOD:</b> bottom-up (see publications in original data source)	
<b>ADDITIONAL INTERCALCULATION:</b> Evaluation and compiling of data coming from original data source and adjusting to recommended Intergovernmental Panel on Climate Change (IPCC) methodology.	
<b>DATA ARCHIVATION:</b> Publications: Potočnik, V. (2000), Report: The basis for methane emission estimation in Croatia 1990-1998, B. Data on Municipal Solid Waste in Croatia 1990-1998 Fundurulja, D., Mužinić, M. (2000) <i>Estimation of the Quantities of Municipal Solid Waste in the Republic of Croatia in the period 1990 – 1998 and 1998 – 2010</i> . Schaller, A. (2000), Republic of Croatia: First National Communication, Waste Management Review – Waste Disposal Sites.	
<b>DATA GAPS:</b> Quantities on MSW were in most cases gained by test weighing in order to estimate average volumes of waste delivered by vehicles and density of MSW.	

<b>MODULE:</b> WASTE	
<b>SUBMODULE:</b> METHANE EMISSIONS FROM SOLID WASTE DISPOSAL SITES	
<b>WORKSHEET:</b> 6-1	<b>SHEET:</b> 1 OF 1 CH <sub>4</sub> EMISSIONS
<b>STEP:</b> 1 TO 4	<b>PAGE:</b> 2 of 2
<b>SUGGESTION FOR THE FUTURE:</b> <ul style="list-style-type: none"> <li>▪ Equipping the major landfills with automatic weigh-bridges in order to accurately estimate the quantities of delivered MSW</li> <li>▪ Providing methodology to determine country-specific MSW composition</li> <li>▪ Periodic analysis of waste composition at major landfills according to provided methodology</li> <li>▪ Modification of Environmental Emissions Inventory (KEO) Reporting Forms regarding to MSW with additional information on waste quantities and composition</li> <li>▪ Adjustment of country-specific to IPCC SWDSs classification, in order to accurately MCF estimation.</li> </ul>	
<b>NOTES:</b> -	
<b>RESPONSIBILITY:</b> Andrea Hublin, M.Sc. EKONERG address: Koranska 5, 10000 Zagreb tel.: +385 1 6000 134 fax.: +385 1 6171 560 e-mail: <a href="mailto:andrea.hublin@ekonerg.hr">andrea.hublin@ekonerg.hr</a>	